arXiv:1303.1307v3 [math.DG] 20 May 2013

THE GAP PHENOMENON IN PARABOLIC GEOMETRIES

BORIS KRUGLIKOV AND DENNIS THE

Abstract. The infinitesimal symmetry algebra of any Cartan geometry has maximum dimension realized by the flat model, but often this dimension drops significantly when considering geometries which are not locally flat, so a gap phenomenon arises. (Here, "not locally flat" means only that the curvature function does not vanish in at least one point.) For general (regular, normal) parabolic geometries of type (G, P), we derive a universal upper bound on the submaximal symmetry dimension using Tanaka theory and Kostant's version of the Bott–Borel–Weil theorem. In almost all complex or split-real cases, we show that this bound is in fact sharp by exhibiting (abstract) models. We give a complete classification of all submaximal symmetry dimensions when G is any complex or split-real simple Lie group.

Contents

1. Introduction 2
2. Background 6
2.1. Tanaka theory in a nutshell 6
2.2. Parabolic subalgebras 8
2.3. Infinitesimal flag structures and prolongation 10
2.4. Regular, normal parabolic geometries 11
2.5. Kostant's theorem 11
2.6. Regularity and Yamaguchi's rigidity theorem 12
2.7. Correspondence and twistor spaces 13
2.8. Annihilators 14
3. Prolongation analysis 16
3.1. Maximizing the Tanaka prolongation 16
3.2. A Dynkin diagram recipe 17
3.3. Prolongation-rigidity 19
3.4. Classical flag varieties 20
3.5. Nested parabolic subalgebras 22
3.6. Prolongation height 23
4. Submaximal symmetry dimensions 24
4.1. Filtration and embedding of the symmetry algebra 24
4.2. A universal upper bound 25
4.3. Establishing submaximal symmetry dimensions 27
4.4. Deformations 31
4.5. Exceptional cases and local homogeneity 32
5. Results and local models for specific geometries 35
5.1. Conformal geometry 35
5.2. Scalar 3rd order ODE 39
5.3. Systems of 2nd order ODE 39

Date: May 21, 2013.
2010 Mathematics Subject Classification. Primary 58J70, Secondary 53B99, 22E46, 17B70.
Key words and phrases. Submaximal symmetry, parabolic geometry, harmonic curvature, Tanaka theory.

5.4. Projective structures 43
5.5. Bracket-generating distributions 44
Appendix A. Representation-theoretic data for complex simple Lie algebras 48
Appendix B. Submaximal symmetry dimensions 48
Appendix C. Direct proof of prolongation-rigidity for conformal geometry 52
References 53

1. Introduction

Let M be a connected manifold. It is well-known that for any Cartan geometry $(\mathcal{G} \to M, \omega)$ of a given type (G, K) (henceforth, a "G/K geometry"), the Lie algebra of (infinitesimal) symmetries $\mathfrak{inf}(\mathcal{G}, \omega)$ (see Definition 2.1.4) is always finite-dimensional. Indeed, we always have $\dim(\mathfrak{inf}(\mathcal{G}, \omega)) \leq \dim(G)$ and equality is realized by the flat model $(G \to G/K, \omega_{MC})$, where ω_{MC} is the Maurer–Cartan form on G. Hence, a natural question is: *Among all G/K geometries $(\mathcal{G} \to M, \omega)$ with $\dim(\mathfrak{inf}(\mathcal{G}, \omega)) < \dim(G)$, what is the maximum of $\dim(\mathfrak{inf}(\mathcal{G}, \omega))$?* Often there is a significant *gap* between this number and $\dim(G)$, i.e. there are forbidden dimensions, so the above question is sometimes referred to as the *gap problem.* However, as stated, the problem is rather too general. We will instead concentrate on classes of Cartan geometries which admit an equivalent description as an underlying geometric structure on M. The gap problem for various geometric structures has been studied since the time of Sophus Lie by many authors, including Fubini, Cartan, Yano, Wakakuwa, Vranceanu, Egorov, Obata, and Kobayashi – see [23, 21, 27] and references therein.

The prototypical example is Riemannian geometry, which in dimension n corresponds to an $\mathrm{E}(n)/\mathrm{O}(n)$ geometry, where $\mathrm{E}(n)$ is the Euclidean group and $\mathrm{O}(n)$ is the orthogonal group. Maximal symmetry occurs (non-uniquely) for the flat model $\mathrm{E}(n)/\mathrm{O}(n) \cong \mathbb{R}^n$, and all other constant curvature spaces. Submaximal symmetry dimensions are given in Table 1 and were studied by Wang [44] and Egorov [18] for $n \geq 3$, and by Darboux and Koenigs in the $n = 2$ case [15].

n	Max	Submax
2	3	1
3	6	4
4	10	8
≥ 5	$\binom{n+1}{2}$	$\binom{n}{2} + 1$

Table 1. Maximal / submaximal symmetry dimensions for Riemannian geometry

A *parabolic geometry* is a G/P geometry, where G is a real or complex semisimple Lie group and $P \subset G$ is a parabolic subgroup. Many well-known geometric structures such as conformal, projective, CR, systems of 2nd order ODE, scalar 3rd order ODE, and many bracket-generating distributions, etc. are equivalently described as parabolic geometries. Indeed, there is an equivalence of categories between *regular, normal* parabolic geometries and underlying structures [10]. For such geometries, maximal symmetry occurs *uniquely* for the (locally) flat model. The gap problem we will focus on is: *Among all* **regular, normal** *G/P geometries $(\mathcal{G} \to M, \omega)$ which are* **not locally flat***, what is the maximum of $\dim(\mathfrak{inf}(\mathcal{G}, \omega))$?* We denote this *submaximal symmetry dimension* by $\mathfrak{S}$. (Equivalently, if we let $\mathcal{S}$ be the symmetry algebra of the underlying structure on M, then $\mathfrak{S}$ maximizes $\dim(\mathcal{S})$ among all structures which are not locally flat.)

In the context of parabolic geometries, previously known results are given in Table 2, but this list comes with some caveats. For example, consider the geometry of generic rank 2 distributions on 5-manifolds [11], also known as $(2, 3, 5)$-distributions, which is modelled on G_2/P_1. Using his method of equivalence, Cartan identified a fundamental binary quartic invariant, whose root type gives a pointwise algebraic classification similar to the Petrov classification of the Weyl tensor in 4-dimensional Lorentzian (conformal) geometry. Cartan then identified the maximal amount of symmetry admitted *within each class with constant root type*, and in many cases exhibited local models either explicitly or via (closed) structure equations. The flat model has G_2 symmetry, which is 14-dimensional, and occurs precisely when the binary quartic vanishes everywhere. Submaximally symmetric models, having 7-dimensional symmetry, all occur within the class where the binary quartic has a single root of multiplicity 4. However, since our only assumption is that the geometry is not locally flat, our result that $\mathfrak{S} = 7$ is a sharpening of Cartan's result. For any (regular, normal) parabolic geometry, there is a fundamental tensorial invariant called *harmonic curvature* κ_H, valued in the Lie algebra cohomology group $H^2(\mathfrak{g}_-, \mathfrak{g})$, whose vanishing everywhere characterizes local flatness. (The Cartan quartic and the Weyl tensor are instances of κ_H for their respective geometries.) For us, "submaximally symmetric" assumes only that κ_H does not vanish *at some point* in the neighbourhood under consideration.[1]

Geometry	Model	Max	Submax	Citation
Scalar 2nd order ODE mod point transformations	$\mathrm{SL}_3(\mathbb{R})/P_{1,2}$	8	3	Tresse (1896)
2-dim. projective structures	$\mathrm{SL}_3(\mathbb{R})/P_1$	8	3	Tresse (1896)
$(2,3,5)$-distributions	G_2/P_1	14	7	Cartan (1910)
Projective structures (dim. $\ell \geq 3$)	$\mathrm{SL}_{\ell+1}(\mathbb{R})/P_1$	$\ell^2+2\ell$	$(\ell-1)^2+4$	Egorov (1951)
Scalar 3rd order ODE mod contact transformations	$\mathrm{Sp}_4(\mathbb{R})/P_{1,2}$	10	5	Wafo Soh, Mahomed, Qu (2002)
Pairs of 2nd order ODE	$\mathrm{SL}_4(\mathbb{R})/P_{1,2}$	15	9	Casey, Dunajski, Tod (2012)

Table 2. Previously known submaximal symmetry dimensions for parabolic geometries

The main results of this article are:

- For any complex or real regular, normal G/P geometry we give a universal upper bound $\mathfrak{S} \leq \mathfrak{U}$ (Theorem 4.2.5), where $\mathfrak{U}$ is algebraically determined – see (3.1).
- In complex or *split-real*[2] cases, we:
    - exhibit models with $\dim(\mathcal{S}) = \mathfrak{U}$ in almost all cases (Theorem 4.3.5). Thus, $\mathfrak{S} = \mathfrak{U}$ almost always (Theorem 4.3.8). Exceptions are also studied (Section 4.5).
    - give a Dynkin diagram recipe to efficiently compute $\mathfrak{U}$ (Section 3.2).
    - establish local homogeneity of all submaximally symmetric models[3] near non-flat *regular* points (Theorem 4.5.2); the set of all such points is open and dense in M (Lemma 4.2.4).
- We recover all results in Table 2; some sample new results are given in Table 3. Our complete classification when G is (complex or split-real) *simple* is presented in Appendix B.

The study of the gap problem is much more subtle for general real forms and we do not attempt to complete the picture here. However, in the conformal case, we exhibit local models in all non-Riemannian and non-Lorentzian signatures which realize the upper bound coming from complexified considerations, so we establish $\mathfrak{S}$ for these cases as well. The problem for Riemannian and Lorentzian conformal structures has been recently settled by B. Doubrov and D. The [16].

---

[1] The global problem of studying symmetry dimensions which are "less than maximal" is quite different. For example, take the flat model $G \to G/P$ and remove a point on the base and the fibre over it. This yields a (locally flat) G/P geometry with *global* automorphism group isomorphic to P. In the G_2/P_1 case, $\dim(P_1) = 9$.

[2] We refer to $\mathfrak{g}$ as split-real if it is a split real form of its complexification, e.g. $\mathfrak{sl}(n, \mathbb{R})$, but not $\mathfrak{su}(n)$.

[3] This is not universally true outside the parabolic context, e.g. Killing fields for metrics on surfaces (see Table 1).

Geometry	Range	Model	Max	Submax
Sig. (p,q) conformal geometry in dim. $n=p+q$	$p,q \geq 2$	$\mathrm{SO}_{p+1,q+1}/P_1$	$\binom{n+2}{2}$	$\binom{n-1}{2}+6$
Systems of 2nd order ODE in m dependent variables	$m \geq 2$	$\mathrm{SL}_{m+2}(\mathbb{R})/P_{1,2}$	$(m+2)^2-1$	m^2+5
Generic rank ℓ distributions on $\frac{1}{2}\ell(\ell+1)$-dim. manifolds	$\ell \geq 3$	$\mathrm{SO}_{\ell,\ell+1}/P_\ell$	$\binom{2\ell+1}{2}$	$\begin{cases} \frac{\ell(3\ell-7)}{2}+10, & \ell \geq 4; \\ 11, & \ell=3 \end{cases}$
Lagrangean contact structures	$\ell \geq 3$	$\mathrm{SL}_{\ell+1}(\mathbb{R})/P_{1,\ell}$	$\ell^2+2\ell$	$(\ell-1)^2+4$
Contact projective structures	$\ell \geq 2$	$\mathrm{Sp}_{2\ell}(\mathbb{R})/P_1$	$\ell(2\ell+1)$	$\begin{cases} 2\ell^2-5\ell+8, & \ell \geq 3; \\ 5, & \ell=2 \end{cases}$
Contact path geometries	$\ell \geq 3$	$\mathrm{Sp}_{2\ell}(\mathbb{R})/P_{1,2}$	$\ell(2\ell+1)$	$2\ell^2-5\ell+9$
Exotic parabolic contact structure of type E_8	-	E_8/P_8	248	147

Table 3. Sample new results of submaximal symmetry dimensions for parabolic geometries

For arbitrary parabolic geometries, Čap & Neusser [9] gave a general algebraic strategy for finding upper bounds on $\mathfrak{S}$ using Kostant's version of the Bott–Borel–Weil theorem [28]. However, the implementation of their strategy must be carried out on a case-by-case basis; moreover, their upper bounds are in general not sharp. (See also Remark 4.1.1.) For structures determined by a bracket-generating distribution (not necessarily underlying parabolic geometries), another approach based on an elaboration of Tanaka theory [40, 41] was proposed in the works [29] and [30] of the first author; the latter reference contains also a review of some results on submaximal symmetry dimensions.

The main idea behind our approach is to combine Tanaka theory with the Čap–Neusser approach based on Kostant theory. This yields a *uniform algebraic approach* to the gap problem which is rooted in the structure theory of semisimple Lie algebras. In contrast, earlier (sharp) results were obtained using a variety of techniques, e.g. computation of the algebra of all differential invariants of a pseudogroup (scalar 2nd order ODE), Cartan's method of equivalence ($(2,3,5)$-distributions), classification of Lie algebras of contact vector fields in the plane (3rd order ODE), or studying integrability conditions for the equations characterizing symmetries (projective structures). Some of these relied heavily on the low-dimensional setup.

In Section 2, we review background from Tanaka theory, representation theory, and parabolic geometry. We discuss the Yamaguchi prolongation and rigidity theorems, Kostant's theorem, as well as correspondence and twistor space constructions. Given a subalgebra $\mathfrak{a}_0$ of the reductive part $\mathfrak{g}_0$ of $\mathfrak{p}$, we introduce in Definition 2.3.3 a slight variant $\mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{a}_0)$ of Tanaka's original prolongation algebra $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{a}_0)$.

Section 3 is representation-theoretic and focuses on the following questions: Letting $\phi \in H^2(\mathfrak{g}_-, \mathfrak{g})$ be **nonzero**, $\mathfrak{ann}(\phi) \subset \mathfrak{g}_0$ its annihilator, and $\mathfrak{a}^\phi := \mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{ann}(\phi))$,

- *What is the maximum of* $\dim(\mathfrak{a}^\phi)$*?*
- *In the maximal case, how can one describe the Lie algebra structure of* $\mathfrak{a}^\phi$*?*
- *What is the maximal height of the grading on* $\mathfrak{a}^\phi$*?*

Over $\mathbb{C}$, we show in Lemma 3.1.1 that $\dim(\mathfrak{a}^\phi)$ must be maximized on a lowest weight vector ϕ_0 of some $\mathfrak{g}_0$-irreducible submodule $\mathbb{V}_\mu \subseteq H^2(\mathfrak{g}_-, \mathfrak{g})$. We then prove a new Dynkin diagram recipe in Section 3.2 which gives a way to understand the structure of $\mathfrak{a}^{\phi_0}$. For example, the E_8/P_8 case (Example 2.8.2 and 3.2.7) becomes a trivial exercise. When $\mathfrak{g}$ is *simple*, regularity forces the grading height $\tilde{\nu}$ of $\mathfrak{a}^\phi$ (for any $\phi \neq 0$) to be highly constrained: $0 \leq \tilde{\nu} \leq 2$ always – see Section 3.6.

Section 4 is the core technical part of the paper. Let $(\mathcal{G} \xrightarrow{\pi} M, \omega)$ be a regular, normal parabolic geometry and let $u \in \mathcal{G}$. Any $\xi \in \mathfrak{inf}(\mathcal{G}, \omega)$ is uniquely determined by its value at u, so the isomorphism $\omega_u : T_u\mathcal{G} \to \mathfrak{g}$ embeds $\mathfrak{inf}(\mathcal{G}, \omega)$ into $\mathfrak{g}$ as a vector subspace. Its image $\mathfrak{f}(u)$ becomes a filtered Lie algebra by restriction of the canonical P-invariant filtration on $\mathfrak{g}$, and the associated-graded $\mathfrak{s}(u) := \mathrm{gr}(\mathfrak{f}(u))$ is a graded subalgebra of $\mathfrak{g}$. If $x = \pi(u)$ is a *regular point* (Definition 4.2.1), then we prove the crucial fact that $\mathfrak{s}(u) \subseteq \mathfrak{a}^{\kappa_H(u)}$ (Proposition 4.2.2), which establishes a bridge to Tanaka theory. Since the set of regular points is open and dense in M (Lemma 4.2.4), any neighbourhood of a non-flat point contains a non-flat regular point, and this leads to our upper bound $\mathfrak{S} \leq \mathfrak{U}$ (Theorem 4.2.5). In complex or split-real cases, realizability of $\mathfrak{U}$ (or more precisely, $\mathfrak{U}_\mu$, when restricting curvature type $\mathrm{im}(\kappa_H) \subseteq \mathbb{V}_\mu \subseteq H^2_+(\mathfrak{g}_-, \mathfrak{g})$) is then addressed: for almost all cases, a model can be constructed by deforming the Lie algebra structure on $\mathfrak{a}^{\phi_0}$ by ϕ_0 (Theorem 4.3.5). We then investigate exceptional cases in Section 4.5.

Section 5 contains finer analysis of specific geometries, and in particular we exhibit some submaximally symmetric models in local coordinates. We mention a few highlights.

- *Conformal geometry:* In addition to investigating the gap problem in general signatures, we find the maximum conformal symmetry dimensions for each Petrov type in the 4-dimensional Lorentzian case. In doing so, we discovered that a Petrov type II metric with four (conformal) Killing vectors was not known in the literature. We exhibit such a metric in Section 5.1.2.
- *Systems of 2nd order ODE:* We discuss the Fels invariants [22] and their connection to harmonic curvature. There are two (degenerate) branches: the projective branch and the Segré branch.[4] Submaximally symmetric models arise in the latter. We exhibit such a model and illustrate the twistor correspondence by exhibiting the corresponding Segré geometry.
- *Projective structures:* We recover Egorov's result [17] by our algebraic method.
- $(2,3,5)$*-distributions:* We establish $\mathfrak{S} = 7$, and then investigate maximal symmetry dimensions for each root type of the binary quartic invariant.
- *Generic rank ℓ distributions on $\binom{\ell+1}{2}$-manifolds:* We obtain $\mathfrak{S}$ in general, and give a submaximally symmetric $(3,6)$-distribution along with its 11 symmetry generators.

Conventions: We assume throughout that M is a connected manifold. When working with real and complex parabolic geometries, our results are formulated in the smooth and holomorphic categories, respectively. We use *left* cosets and *right* principal bundles. For a Lie group G, we identify its Lie algebra $\mathfrak{g} := \mathrm{Lie}(G)$ with the *left*-invariant vector fields on G. We write $A_\ell, B_\ell, C_\ell, D_\ell, G_2, F_4, E_6, E_7, E_8$ for the complex simple Lie algebras (see Appendix A), **or** any complex Lie groups having these as their Lie algebras. (In Section 5, we abuse this notation further by letting it refer to real forms, and specify the precise real form as necessary.) Parabolic subalgebras will be denoted $\mathfrak{p}$, $\mathfrak{q}$, and corresponding parabolic subgroups are P, Q.

We always assume that G acts on G/P *infinitesimally effectively*, i.e. the kernel of the G-action on G/P is at most discrete. Equivalently, simple ideals of $\mathfrak{g}$ are not contained in $\mathfrak{g}_0$. (This avoids situations like $G = G' \times G''$ and $P = P' \times G''$, where the G''-action is not visible on G/P.)

For complex semisimple Lie algebras, we draw our Dynkin diagrams with open white circles. This is the same notation as the Satake diagram for the corresponding split real form, and serves to emphasize that all our results are the same in both settings. We use the "minus lowest weight" Dynkin diagram convention (see Section 2.5) when referring to irreducible $\mathfrak{g}_0$-modules. If $\mathfrak{g}$ is simple, we use $\lambda_{\mathfrak{g}}$ to denote its highest weight (root).

Acknowledgements: We are grateful for many helpful discussions with Boris Doubrov, Mike Eastwood, Katharina Neusser, Katja Sagerschnig, and Travis Willse. Much progress on this paper was made during the conference "The Interaction of Geometry and Representation Theory:

---

[4] Equations in the Segré branch correspond to the *torsion-free* path geometries in the sense of Grossman [25].

Exploring New Frontiers" devoted to Mike Eastwood's 60th birthday, and held in Vienna in September 2012 at the Erwin Schrödinger Institute. Boris Doubrov gave some key insights during this conference which led to the proof of Lemma 3.1.1.

The representation theory software `LiE`, as well as Ian Anderson's `DifferentialGeometry` package in `Maple` were invaluable tools for facilitating the analysis in this paper.

B.K. was supported by the University of Tromsø while visiting the Australian National University, where this work was initiated. The hospitality of ANU is gratefully acknowledged. D.T. was partially supported by an ARC Research Fellowship.

2. BACKGROUND

We review some necessary background from Tanaka theory [40, 41, 47], representation theory and parabolic geometry [10], including Kostant's version of the Bott–Borel–Weil theorem [28, 1, 10].

2.1. Tanaka theory in a nutshell.

The aim of Tanaka theory is to study the equivalence problem for geometric structures. The given geometric data is a manifold M with a (vector) distribution $D \subseteq TM$ endowed possibly with some additional structure on it, e.g. a sub-Riemannian metric or a conformal structure. For many interesting geometric structures, one can canonically associate a Cartan geometry $(\mathcal{G} \to M, \omega)$ of some type (G, K). We give an outline of these ideas.

Iterating Lie brackets of sections of D, we form the weak derived flag

$$D =: D^{-1} \subset D^{-2} \subset D^{-3} \subset \dots\ , \quad \text{where} \quad \Gamma(D^{i-1}) := [\Gamma(D^i), \Gamma(D^{-1})].$$

We assume that all D^i have constant rank, and D is bracket-generating in TM, i.e. $D^{-\nu} = TM$ for some minimal $\nu \geq 1$ (called the *depth*). By construction, the Lie bracket satisfies $\Gamma(D^i) \times \Gamma(D^j) \to \Gamma(D^{i+j})$, so M is a *filtered manifold*. On the associated graded of the above filtration

$$\mathfrak{m}(x) = \mathfrak{g}_-(x) = \bigoplus_{i<0} \mathfrak{g}_i(x), \qquad \mathfrak{g}_i(x) = D^i(x)/D^{i+1}(x),$$

the Lie bracket induces a tensorial bracket on $\mathfrak{m}(x)$ called the Levi bracket, which turns $\mathfrak{m}(x)$ into a graded nilpotent Lie algebra (GNLA) called the *symbol algebra at* x. We further assume that D is of *constant type*, i.e. there is a fixed GNLA $\mathfrak{m} = \mathfrak{g}_-$ such that $\mathfrak{m}(x) \cong \mathfrak{m}$, $\forall x \in M$.

Example 2.1.1. *Any* $(2,3,5)$*-distribution* D *is locally specified by an underdetermined ODE* $z' = F(x, y, y', y'', z)$ *(Monge equation) satisfying* $F_{y''y''} \neq 0$. *On a 5-manifold* M *with coordinates* (x, y, p, q, z), $D = D_F$ *is spanned by* ∂_q *and* $\frac{d}{dx} := \partial_x + p\partial_y + q\partial_p + F\partial_z$. *We have*

$$\left[\partial_q, \frac{d}{dx}\right] = \partial_p + F_q\partial_z, \quad [\partial_q, \partial_p + F_q\partial_z] = F_{qq}\partial_z, \quad \left[\frac{d}{dx}, \partial_p + F_q\partial_z\right] = -\partial_y + H\partial_z,$$

for some differential function H *of* F. *Assuming* $F_{qq} \neq 0$, *we get a filtration* $D = D^{-1} \subset D^{-2} \subset D^{-3} = TM$. *Its associated graded is isomorphic to* $\mathfrak{g}_- = \mathfrak{g}_{-1} \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-3}$ *with relations*

$$[e^1_{-1}, e^2_{-1}] = e_{-2}, \quad [e^1_{-1}, e_{-2}] = e^1_{-3}, \quad [e^2_{-1}, e_{-2}] = e^2_{-3}. \tag{2.1}$$

Since D has constant type, consider the graded frame bundle $\mathcal{F}_{gr}(M) \xrightarrow{\pi} M$ whose total space consists of all GNLA isomorphisms $u : \mathfrak{m} \to \mathfrak{m}(x)$, where $x = \pi(u)$. Indeed, $\mathcal{F}_{gr}(M) \to M$ is a (right) principal bundle with structure group $\mathrm{Aut}_{gr}(\mathfrak{m})$, the group of graded automorphisms of $\mathfrak{m}$. If D is endowed with additional structure, we can reduce to a principal subbundle $\mathcal{G}_0 \to M$ with structure group $G_0 \subseteq \mathrm{Aut}_{gr}(\mathfrak{m})$. For example, in conformal geometry, $D = TM$, and $\mathcal{G}_0 \to M$ is the conformal frame bundle with $G_0 = \mathrm{CO}(\mathfrak{m}_{-1})$; in Example 2.1.1, $G_0 \cong \mathrm{Aut}_{gr}(\mathfrak{m}) \cong \mathrm{GL}(\mathfrak{g}_{-1}) \cong \mathrm{GL}_2$. From here, one constructs the *geometric prolongation* of the given structure, namely a tower of adapted bundles $\dots \to \mathcal{G}_2 \to \mathcal{G}_1 \to \mathcal{G}_0 \to M$. We refer to [47] for details on this procedure. This geometric prolongation is controlled by an *algebraic prolongation* which we describe below.

At the Lie algebra level, $\mathfrak{g}_0 \subseteq \mathfrak{der}_{gr}(\mathfrak{m})$, where the latter is the algebra of graded derivations of $\mathfrak{m}$. Given $(\mathfrak{m}, \mathfrak{g}_0)$, the axioms for Tanaka's algebraic prolongation $\mathrm{pr}(\mathfrak{m}, \mathfrak{g}_0) := \bigoplus_{i\in\mathbb{Z}} \mathfrak{g}_i(\mathfrak{m}, \mathfrak{g}_0)$ are:

(T.1) $\mathfrak{g}_{\leq 0}(\mathfrak{m}, \mathfrak{g}_0) = \mathfrak{m} \oplus \mathfrak{g}_0$;

(T.2) If $X \in \mathfrak{g}_i(\mathfrak{m}, \mathfrak{g}_0)$ for $i \geq 0$ satisfies $[X, \mathfrak{g}_{-1}] = 0$, then $X = 0$;

(T.3) $\mathrm{pr}(\mathfrak{m}, \mathfrak{g}_0)$ is the maximal graded Lie algebra satisfying (T.1) and (T.2).

Write $\mathrm{pr}(\mathfrak{m}) := \mathrm{pr}(\mathfrak{m}, \mathfrak{der}_{gr}(\mathfrak{m}))$. Up to isomorphism, there is a unique graded Lie algebra satisfying (T.1)–(T.3). In fact, Tanaka gives an explicit inductive realization of $\mathfrak{g} := \mathrm{pr}(\mathfrak{m}, \mathfrak{g}_0)$; for $i > 0$,

$$\mathfrak{g}_i = \left\{ f \in \bigoplus_{j<0} \mathfrak{g}_j^* \otimes \mathfrak{g}_{j+i} \mid f([v_1, v_2]) = [f(v_1), v_2] + [v_1, f(v_2)],\ \forall v_1, v_2 \in \mathfrak{m} \right\}. \tag{2.2}$$

The brackets on $\mathfrak{g}$ are: (i) The given brackets on $\mathfrak{m}$ and $\mathfrak{g}_0$; (ii) If $f \in \mathfrak{g}_i$, $i \geq 0$, and $v \in \mathfrak{m}$, then define $[f, v] = -[v, f] := f(v)$; (iii) Brackets on the non-negative part are defined inductively: If $f_1 \in \mathfrak{g}_i$ and $f_2 \in \mathfrak{g}_j$, for $i, j \geq 0$, then define $[f_1, f_2] \in \mathfrak{g}_{i+j}$ by $[f_1, f_2](v) := [f_1(v), f_2] + [f_1, f_2(v)]$, $v \in \mathfrak{m}$. (Note $f_1(v) \in \mathfrak{g}_{<i}$ and $f_2(v) \in \mathfrak{g}_{<j}$, so the brackets on the right side are known by induction.) Verification of the Jacobi identity is left to the interested reader.

Remark 2.1.2. Since $\mathfrak{g}_-$ is generated by $\mathfrak{g}_{-1}$, any $X \in \mathfrak{g}_i$, $i \geq 0$, is determined by its action on $\mathfrak{g}_{-1}$. By (T.2) the Lie bracket on $\mathfrak{g}$ induces $\mathfrak{g}_i \hookrightarrow \mathfrak{g}_{-1}^* \otimes \mathfrak{g}_{i-1}$, so $\mathfrak{g}_i \hookrightarrow \mathrm{Hom}(\otimes^i \mathfrak{g}_{-1}, \mathfrak{g}_0) \hookrightarrow \mathrm{Hom}(\otimes^{i+1} \mathfrak{g}_{-1}, \mathfrak{g}_{-1})$.

Note that by (T.2), if $\mathfrak{g}_r = 0$ for some $r \geq 0$, then $\mathfrak{g}_i = 0$ for all $i > r$. This case of finite termination is particularly important.

Theorem 2.1.3 (Tanaka). *Let $\mathcal{G}_0 \to M$ be a structure of constant type $(\mathfrak{m}, \mathfrak{g}_0)$ and $\mathfrak{g} = \mathrm{pr}(\mathfrak{m}, \mathfrak{g}_0)$. Suppose $r \geq 0$ is minimal such that $\mathfrak{g}_{r+1} = 0$. Then there exists a canonical frame on the r-th geometric prolongation $\mathcal{G}_r$ of $\mathcal{G}_0$. Moreover, any such structure can be described as a Cartan geometry of some type (G, K), where $\mathfrak{g} = \mathrm{Lie}(G)$ and $\mathfrak{g}_{\geq 0} =: \mathfrak{k} = \mathrm{Lie}(K)$.*

We recall some definitions from Cartan geometry:

Definition 2.1.4. *A Cartan geometry $(\mathcal{G} \to M, \omega)$ of type (G, K) (or a "G/K geometry") is a principal K-bundle $\mathcal{G} \to M$ endowed with a Cartan connection $\omega \in \Omega^1(\mathcal{G}, \mathfrak{g})$ with defining properties: (i) ω is K-equivariant; (ii) $\omega(\zeta_Y) = Y$ for any $Y \in \mathfrak{k}$, where $\zeta_Y(u) = \frac{d}{dt}\Big|_{t=0} u \cdot \exp(tY)$; (iii) $\omega_u : T_u\mathcal{G} \to \mathfrak{g}$ is a linear isomorphism for any $u \in \mathcal{G}$. The curvature of ω is $d\omega + \frac{1}{2}[\omega, \omega] \in \Omega^2(\mathcal{G}, \mathfrak{g})$. Evaluation on $\omega^{-1}(X)$, $X \in \mathfrak{g}$, yields the curvature function $\kappa : \mathcal{G} \to \bigwedge^2 \mathfrak{g}^* \otimes \mathfrak{g}$, which is horizontal, so $\kappa : \mathcal{G} \to \bigwedge^2(\mathfrak{g}/\mathfrak{k})^* \otimes \mathfrak{g}$. The infinitesimal symmetries are $\mathfrak{inf}(\mathcal{G}, \omega) = \{\xi \in \mathfrak{X}(\mathcal{G})^K \mid \mathcal{L}_\xi \omega = 0\}$.*

While calculating the Tanaka prolongation from given data $(\mathfrak{m}, \mathfrak{g}_0)$ is algorithmic, a naive application of (2.2) generally leads to a computationally intensive exercise in linear algebra. Moreover, even if this task is completed, there still remains the general problem of understanding the structure of the resulting Tanaka algebra. However, in the context of parabolic geometries, these aforementioned problems are resolved by Yamaguchi's prolongation theorem [45]; see Theorem 2.3.1.

A general situation in which the computation of the Tanaka prolongation is much simpler is when one already knows $\mathfrak{g} = \mathrm{pr}(\mathfrak{m}, \mathfrak{g}_0)$, and one is interested in $\mathfrak{a} = \mathrm{pr}(\mathfrak{m}, \mathfrak{a}_0)$ for some subalgebra $\mathfrak{a}_0 \subseteq \mathfrak{g}_0$. By Remark 2.1.2, for $k > 0$, we have $\mathfrak{a}_k \hookrightarrow \mathfrak{g}_{-1}^* \otimes \mathfrak{a}_{k-1} \hookrightarrow \mathfrak{g}_{-1}^* \otimes \mathfrak{g}_{k-1}$, and $\mathfrak{a}_k \hookrightarrow \mathrm{Hom}(\otimes^k \mathfrak{g}_{-1}, \mathfrak{a}_0)$. The inclusion $\mathfrak{a}_0 \hookrightarrow \mathfrak{g}_0$ induces inclusions $\mathfrak{a}_k \hookrightarrow \mathfrak{g}_k$. Hence,

Lemma 2.1.5. *Suppose $\mathfrak{m} = \mathfrak{g}_-$ is generated by $\mathfrak{g}_{-1}$. If $\mathfrak{a}_0 \subseteq \mathfrak{g}_0$ is any subalgebra, then $\mathfrak{a} := \mathrm{pr}(\mathfrak{m}, \mathfrak{a}_0) \hookrightarrow \mathfrak{g} := \mathrm{pr}(\mathfrak{m}, \mathfrak{g}_0)$. Indeed for $k > 0$,*

$$\mathfrak{a}_k = \{X \in \mathfrak{g}_k \mid [X, \mathfrak{g}_{-1}] \subset \mathfrak{a}_{k-1}\} = \{X \in \mathfrak{g}_k \mid \mathrm{ad}_{\mathfrak{g}_{-1}}^k(X) \subset \mathfrak{a}_0\}.$$

More precisely, $\mathfrak{a}_k = \{X \in \mathfrak{g}_k \mid \mathrm{ad}_{u_1} \circ \ldots \circ \mathrm{ad}_{u_k}(X) \in \mathfrak{a}_0,\ \forall u_i \in \mathfrak{g}_{-1}\}$.

2.2. **Parabolic subalgebras.** Given a real or complex semisimple Lie algebra $\mathfrak{g}$, a $\mathbb{Z}$*-grading* with *depth* $\nu \in \mathbb{Z}_{\geq 0}$ (or a ν*-grading*) is a vector space decomposition

$$\mathfrak{g} = \overbrace{\mathfrak{g}_{-\nu} \oplus \dots \oplus \mathfrak{g}_{-1}}^{\mathfrak{g}_-} \oplus \overbrace{\mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_\nu}^{\mathfrak{p}}, \tag{2.3}$$

with: (i) $\mathfrak{g}_{\pm\nu} \neq 0$; (ii) $[\mathfrak{g}_i, \mathfrak{g}_j] \subseteq \mathfrak{g}_{i+j}$, $\forall i, j$; (iii) $[\mathfrak{g}_j, \mathfrak{g}_{-1}] = \mathfrak{g}_{j-1}$ for $j < 0$, i.e. $\mathfrak{g}_{-1}$ is *bracket-generating* in $\mathfrak{g}_-$. The subalgebra $\mathfrak{g}_0$ is reductive with $\mathfrak{g}_0 = \mathfrak{z}(\mathfrak{g}_0) \oplus \mathfrak{g}_0^{ss}$, and each $\mathfrak{g}_j$ is a $\mathfrak{g}_0$-module. (Here, $\mathfrak{z}(\mathfrak{g}_0)$ is the centralizer of $\mathfrak{g}_0$ in $\mathfrak{g}$, and $\mathfrak{g}_0^{ss}$ is the semisimple part of $\mathfrak{g}_0$.) Defining $\mathfrak{g}^i := \bigoplus_{j \geq i} \mathfrak{g}_j$, we have $[\mathfrak{g}^i, \mathfrak{g}^j] \subseteq \mathfrak{g}^{i+j}$, so $\mathfrak{g}$ is canonically a filtered Lie algebra. A subalgebra $\mathfrak{p} \subset \mathfrak{g}$ is *parabolic* if $\mathfrak{p} = \mathfrak{g}^0 = \mathfrak{g}_{\geq 0}$ for some ν-grading of $\mathfrak{g}$. (We call $\mathfrak{p}^{\rm op} = \mathfrak{g}_{\leq 0}$ the *opposite* parabolic.) Each filtrand $\mathfrak{g}^i$ is $\mathfrak{p}$-invariant, and the quotient $\mathfrak{g}/\mathfrak{p}$ is naturally filtered. The Killing form $B(X, Y) = \mathrm{tr}(\mathrm{ad}_X \circ \mathrm{ad}_Y)$ is compatible with the grading and filtration on $\mathfrak{g}$, so induces the dualities $\mathfrak{g}_{-i} \cong (\mathfrak{g}_i)^*$ (as $\mathfrak{g}_0$-modules) and $\mathfrak{g}^i = (\mathfrak{g}^{-i+1})^\perp$ (annihilator with respect to B).

Let G be a semisimple Lie group with Lie algebra $\mathfrak{g}$ and parabolic subalgebra $\mathfrak{p} \subset \mathfrak{g}$. A subgroup $P \subset G$ is *parabolic* if it lies between the normalizer $N_G(\mathfrak{p})$ and its connected component of the identity. Under the adjoint action, P preserves the filtration on $\mathfrak{g}$. Define $G_0 \subset P$ to be the subgroup which preserves the grading on $\mathfrak{g}$; its Lie algebra is $\mathfrak{g}_0$. There is a decomposition $P = G_0 \ltimes P_+$ for some closed normal subgroup $P_+ \subset P$ with Lie algebra $\mathfrak{p}_+ := \mathfrak{g}_+$.

We focus now on the complex case and recall some representation theory. Let $\mathfrak{b} \subset \mathfrak{g}$ be a Borel subalgebra. This is equivalent to a choice of Cartan subalgebra $\mathfrak{h} \subset \mathfrak{b}$, a root system $\Delta \subset \mathfrak{h}^*$, and a basis of simple roots $\Delta^0 = \{\alpha_1, \dots, \alpha_\ell\}$, where $\ell = \dim(\mathfrak{h}) = \mathrm{rank}(\mathfrak{g})$. (We use the Bourbaki ordering – see Appendix A.) For any $\alpha \in \Delta$, write $\alpha = \sum_{i=1}^\ell m_i(\alpha)\alpha_i$, where $m_i(\alpha) \in \mathbb{Z}$ are all non-negative or all non-positive, and $\mathfrak{g}_\alpha$ is the corresponding root space, with $\dim(\mathfrak{g}_\alpha) = 1$. If $\alpha, \beta \in \Delta$, we have: (i) $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] = \mathfrak{g}_{\alpha+\beta}$ if $\alpha + \beta \in \Delta$; (ii) $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] = 0$ if $\alpha + \beta \notin \Delta$ and $\beta \neq -\alpha$; (iii) $[\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] \subset \mathfrak{h}$. If a subspace $\mathfrak{k} \subset \mathfrak{g}$ is a direct sum of root spaces and possibly some subspace of $\mathfrak{h}$, let $\Delta(\mathfrak{k})$ denote the corresponding collection of roots. Thus, $\mathfrak{b} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$ and $\Delta(\mathfrak{b}) = \Delta^+$. If $\mathfrak{g}$ is simple, there is a unique highest root, which we denote by $\lambda_{\mathfrak{g}}$.

The Killing form B induces a symmetric pairing $\langle \cdot, \cdot \rangle$ on $\mathfrak{h}^*$. This determines the Cartan matrix $c_{ij} = \langle \alpha_i, \alpha_j^\vee \rangle \in \mathbb{Z}$, where $\alpha^\vee = \frac{2\alpha}{\langle \alpha, \alpha \rangle}$. We have $c_{ij} = 2$ iff $i = j$; otherwise, $-3 \leq c_{ij} \leq 0$. The Dynkin diagram $\mathcal{D}(\mathfrak{g})$ of $\mathfrak{g}$ is the graph whose i-th node corresponds to α_i, and nodes i, j are connected by an edge of multiplicity $c_{ij}c_{ji} \leq 3$ (or disconnected if $c_{ij}c_{ji} = 0$). Moreover, if $c_{ij}c_{ji} = 2$ or 3, this edge is directed towards the shorter root (i.e. if $c_{ij} = -1$, then it is directed from j to i).

Index sets $I \subset \{1, \dots, \ell\}$ correspond to (standard) parabolic subalgebras $\mathfrak{p} \supset \mathfrak{b}$:

$$\mathfrak{p} \mapsto I_{\mathfrak{p}} := \{i \mid \mathfrak{g}_{-\alpha_i} \not\subseteq \mathfrak{p}\}, \qquad I \mapsto \mathfrak{p}_I := \mathfrak{b} \oplus \bigoplus_{\substack{\alpha \in \Delta^+ \\ m_i(\alpha) = 0,\, \forall i \in I}} \mathfrak{g}_{-\alpha}.$$

We encode $\mathfrak{p}$ on $\mathcal{D}(\mathfrak{g})$ by putting a cross over all nodes corresponding to $I_{\mathfrak{p}}$. This marked Dynkin diagram will be denoted $\mathcal{D}(\mathfrak{g}, \mathfrak{p})$. Thus, $I_{\mathfrak{b}} = \{1, \dots, \ell\}$ and all nodes are crossed.

Let $\{Z_1, \dots, Z_\ell\}$ be the basis of $\mathfrak{h}$ dual to $\Delta^0 \subset \mathfrak{h}^*$. Fixing a parabolic $\mathfrak{p}$ determines the *grading element* $Z = Z_I = \sum_{i \in I_{\mathfrak{p}}} Z_i$, which induces a $\mathbb{Z}$-grading (2.3) via $\mathfrak{g}_j = \{x \in \mathfrak{g} \mid [Z, x] = jx\}$, and $\mathfrak{z}(\mathfrak{g}_0) \subset \mathfrak{g}_0$ is spanned by $\{Z_i\}_{i \in I_{\mathfrak{p}}}$. The grading has depth $\nu = \max\{Z(\lambda_{\mathfrak{g}'}) \mid \mathfrak{g}' \subset \mathfrak{g} \text{ simple ideal}\}$.

Recipe 1. *Deleting the* $I_{\mathfrak{p}}$ *nodes from* $\mathcal{D}(\mathfrak{g}, \mathfrak{p})$ *yields* $\mathcal{D}(\mathfrak{g}_0^{ss})$, *and* $\dim(\mathfrak{z}(\mathfrak{g}_0)) = |I_{\mathfrak{p}}|$. *Also,* $\dim(\mathfrak{g}_-) = \frac{1}{2}(\dim(\mathfrak{g}) - \dim(\mathfrak{g}_0))$ *and* $\dim(\mathfrak{p}) = \frac{1}{2}(\dim(\mathfrak{g}) + \dim(\mathfrak{g}_0))$.

Given a weight $\lambda \in \mathfrak{h}^*$, we refer to $Z(\lambda)$ as the *homogeneity* of λ. Let $\{\lambda_1, \dots, \lambda_\ell\} \subset \mathfrak{h}^*$ denote the fundamental weights of $\mathfrak{g}$, defined by $\langle \lambda_i, \alpha_j^\vee \rangle = \delta_{ij}$. Encode $\lambda = \sum_{i=1}^\ell r_i(\lambda)\lambda_i$, where $r_i(\lambda) :=$

$\langle \lambda, \alpha_i^\vee \rangle$, on a $\mathcal{D}(\mathfrak{g})$ or $\mathcal{D}(\mathfrak{g}, \mathfrak{p})$ by inscribing $r_i(\lambda)$ over the i-th node. If $r_i(\lambda)$ is nonnegative [integral] for all i, then λ is $\mathfrak{g}$-dominant [integral]. If these statements hold only for $i \in \{1, ..., \ell\} \backslash I_\mathfrak{p}$, then λ is $\mathfrak{p}$-dominant [integral]. The Cartan matrix is the transition matrix between $\{\alpha_i\}_{i=1}^\ell$ and $\{\lambda_i\}_{i=1}^\ell$,

$$\lambda = \sum_{i=1}^{\ell} m_i \alpha_i = \sum_{i=1}^{\ell} r_i \lambda_i \quad \Rightarrow \quad \sum_{i=1}^{\ell} m_i c_{ij} = r_j. \tag{2.4}$$

Some distinguished weights are $\rho = \sum_{i=1}^\ell \lambda_i$ and $\rho^\mathfrak{p} = \sum_{i \in I_\mathfrak{p}} \lambda_i$, as well as the highest weight (root) $\lambda_\mathfrak{g}$ of any simple Lie algebra $\mathfrak{g}$ (see Appendix A).

Example 2.2.1. *G_2 has $\lambda_\mathfrak{g} = \lambda_2 = 3\alpha_1 + 2\alpha_2$. Also, $\rho = \lambda_1 + \lambda_2$. If $I = \{1\}$, then $Z = Z_1$, so $\lambda_\mathfrak{g}$ has homogeneity $+3$. Also, $\mathfrak{p} = \mathfrak{p}_I = \mathfrak{g}_{\geq 0} = \mathfrak{b} \oplus \mathfrak{g}_{-\alpha_2}$, and $\rho^\mathfrak{p} = \lambda_1$. We write*

$$\rho = \overset{1}{\circ}\!\Lleftarrow\!\overset{1}{\circ}\ , \qquad \lambda_\mathfrak{g} = \overset{0}{\circ}\!\Lleftarrow\!\overset{1}{\circ}\ , \qquad \mathfrak{p} = \times\!\Lleftarrow\!\circ\ , \qquad \rho^\mathfrak{p} = \overset{1}{\times}\!\Lleftarrow\!\overset{0}{\circ}\ .$$

Given any $\alpha \in \Delta$, the reflection $\sigma_\alpha \in O(\mathfrak{h}^*)$ is defined by $\sigma_\alpha(\beta) = \beta - \langle \beta, \alpha^\vee \rangle \alpha$. The Weyl group $W \subset O(\mathfrak{h}^*)$ is the group generated by all simple reflections $\sigma_i := \sigma_{\alpha_i}$. Representing w as a product of simple reflections, let $|w|$ denote its length, i.e. the minimal length of a representative. We will use the notation $(ij) := \sigma_i \sigma_j$, which means that σ_j acts first, followed by σ_i. There is a simple Dynkin diagram recipe [1] for the (standard) W-action of a simple reflection on weights:

> **Recipe 2.** *To compute $\sigma_i(\lambda)$, add $c = \langle \lambda, \alpha_i^\vee \rangle$ (i.e. i-th coefficient) to adjacent coefficients, with multiplicity if there is a multiple edge directed towards the adjacent node. Then replace c by $-c$.*

We will also use the affine action of W, which is defined by

$$w \cdot \lambda := w(\lambda + \rho) - \rho. \tag{2.5}$$

Defining the inversion set $\Phi_w = w(\Delta^-) \cap \Delta^+$, we have [10, Proposition 3.2.14]:

$$|\Phi_w| = |w|, \qquad \sum_{\alpha \in \Phi_w} \alpha = -w \cdot 0. \tag{2.6}$$

The *Hasse diagram* is $W^\mathfrak{p} = \{w \in W \mid \Phi_w \subset \Delta(\mathfrak{g}_+)\}$ with length r elements $W^\mathfrak{p}(r)$. (Equivalently, $w \in W^\mathfrak{p}$ sends $\mathfrak{g}$-dominant weights to $\mathfrak{p}$-dominant weights.) By a lemma of Kostant [1],

> **Recipe 3.** *$W^\mathfrak{p} \overset{1:1}{\longleftrightarrow}$ W-orbit through $\rho^\mathfrak{p}$ under the right action $(\rho^\mathfrak{p}, w) \mapsto w^{-1}\rho^\mathfrak{p}$. In particular, $w = (jk) \in W^\mathfrak{p}(2)$ iff $j \in I_\mathfrak{p}$ and $k \in (I_\mathfrak{p} \cup \mathcal{N}(j)) \backslash \{j\}$, where $\mathcal{N}(j) = \{i \mid c_{ij} \leq -1\}$.*

The *right* action forces a *reversal* of the order of reflections from their order of application above.

Example 2.2.2 (G_2/P_1)**.** *For $\times\!\Lleftarrow\!\circ$, $\mathfrak{z}(\mathfrak{g}_0) = \mathrm{span}\{Z_1\}$, $\mathfrak{g}_0^{ss} \cong \mathfrak{sl}_2(\mathbb{C})$, and $W^\mathfrak{p}(2) = \{(12)\}$ since*

$$\rho^\mathfrak{p} = \overset{1}{\times}\!\Lleftarrow\!\overset{0}{\circ} \xrightarrow{\sigma_1} \overset{-1}{\times}\!\Lleftarrow\!\overset{1}{\circ} \xrightarrow{\sigma_2} \overset{2}{\times}\!\Lleftarrow\!\overset{-1}{\circ}\ .$$

(This also follows from the second statement in Recipe 3.) Using $w = (12)$ and $\lambda_\mathfrak{g} = \lambda_2 =$ $\overset{0}{\times}\!\Lleftarrow\!\overset{1}{\circ}$,

$$w \cdot \lambda_\mathfrak{g} = (12) \cdot \left(\overset{0}{\times}\!\Lleftarrow\!\overset{1}{\circ} \right) = (12) \left(\overset{1}{\times}\!\Lleftarrow\!\overset{2}{\circ} \right) - \rho = (1) \left(\overset{7}{\times}\!\Lleftarrow\!\overset{-2}{\circ} \right) - \rho = \overset{-8}{\times}\!\Lleftarrow\!\overset{4}{\circ}\ ,$$

and since $Z = Z_1$, then $-w \cdot \lambda_\mathfrak{g} = 8\lambda_1 - 4\lambda_2 = +4\alpha_1$ has homogeneity $+4$.

2.3. **Infinitesimal flag structures and prolongation.** Let $(\mathcal{G} \to M, \omega)$ be a G/P geometry with curvature function $\kappa : \mathcal{G} \to \bigwedge^2(\mathfrak{g}/\mathfrak{p})^* \otimes \mathfrak{g}$. Since ω trivializes $T\mathcal{G} \cong \mathcal{G} \times \mathfrak{g}$, the filtration $\{\mathfrak{g}^i\}_{i=-\nu}^{\nu}$ on $\mathfrak{g}$ induces a (P-invariant) filtration $\{T^i\mathcal{G}\}_{i=-\nu}^{\nu}$ of $T\mathcal{G}$. This projects to a filtration $\{T^iM\}_{i=-\nu}^{-1}$ of $TM = \mathcal{G} \times_P (\mathfrak{g}/\mathfrak{p})$, and defining the principal G_0-bundle $\mathcal{G}_0 := \mathcal{G}/P_+ \to M$, we have $\mathrm{gr}(TM) \cong \mathcal{G}_0 \times_{G_0} \mathfrak{g}_-$. Then M becomes a filtered manifold iff $\kappa(\mathfrak{g}_i, \mathfrak{g}_j) \subset \mathfrak{g}^{i+j}$ for all $i, j < 0$ [10, Corollary 3.1.8]. In this case, the parabolic geometry is *regular* if the algebraic bracket on $\mathrm{gr}(TM)$ induced from $\mathfrak{g}_-$ is the same as the Levi bracket. Indeed, regularity is equivalent to $\kappa(\mathfrak{g}_i, \mathfrak{g}_j) \subset \mathfrak{g}^{i+j+1}$. In the latter case, we get a *regular infinitesimal flag structure* of type (G, P) on M, i.e. a filtered manifold M generated by $D = T^{-1}M$ together with a reduction of structure group of the frame bundle of $\mathrm{gr}(TM)$ to $\mathcal{G}_0$ corresponding to $\mathrm{Ad} : G_0 \to \mathrm{Aut}_{gr}(\mathfrak{g}_-)$.

Conversely, a regular infinitesimal flag structure of type (G, P) determines a regular parabolic geometry. There is an algebraic statement underlying this fact and motivated by the following question: Given $(\mathfrak{g}, \mathfrak{p})$ and the resulting $\mathbb{Z}$-grading (2.3), what are $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0)$ or $\mathrm{pr}(\mathfrak{g}_-)$?[5]

Theorem 2.3.1 (Yamaguchi's prolongation theorem [45])**.** *Let $\mathfrak{g} = \bigoplus_{k\in\mathbb{Z}} \mathfrak{g}_k$ be a simple graded Lie algebra over $\mathbb{C}$ such that $\mathfrak{g}_{-1}$ generates $\mathfrak{g}_-$. Then $\mathfrak{g} \cong \mathrm{pr}(\mathfrak{g}_-)$ except for:*

(a) 1-gradings: A_ℓ/P_k, B_ℓ/P_1, C_ℓ/P_ℓ, D_ℓ/P_1, D_ℓ/P_ℓ, E_6/P_6, E_7/P_7.

(b) contact gradings, i.e. 2-gradings with $\dim(\mathfrak{g}_{-2}) = 1$, and non-degenerate bracket $\bigwedge^2 \mathfrak{g}_{-1} \to \mathfrak{g}_{-2}$:

$$A_\ell/P_{1,\ell}, \quad B_\ell/P_2, \quad C_\ell/P_1, \quad D_\ell/P_2, \quad G_2/P_2, \quad F_4/P_1, \quad E_6/P_2, \quad E_7/P_1, \quad E_8/P_8.$$

(c) $(\mathfrak{g}, \mathfrak{p}) \cong (A_\ell, P_{1,i})$ with $\ell \geq 3$ and $i \neq 1, \ell$, or $(C_\ell, P_{1,\ell})$ with $\ell \geq 2$.

Moreover, we always have $\mathfrak{g} \cong \mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0)$ except when $(\mathfrak{g}, \mathfrak{p}) \cong (A_\ell, P_1)$ or (C_ℓ, P_1).

If $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0) \cong \mathfrak{g}$, then there is a regular parabolic geometry (e.g. the *normal* one – see next section) which is completely determined by its underlying regular infinitesimal flag structure.

Example 2.3.2. *In Example 2.1.1, the symbol algebra $\mathfrak{g}_-$ is isomorphic to the $\mathfrak{g}_-$ in* (2.3) *arising from $\mathfrak{p}_1 \subset \mathrm{Lie}(G_2) =: \mathfrak{g}$. Namely, we can choose root vectors such that*

$$e^1_{-1} = e_{-\alpha_1}, \quad e^2_{-1} = e_{-\alpha_1-\alpha_2}, \quad e_{-2} = e_{-2\alpha_1-\alpha_2}, \quad e^1_{-3} = e_{-3\alpha_1-\alpha_2}, \quad e^2_{-3} = e_{-3\alpha_1-2\alpha_2}$$

satisfy the bracket relations (2.1)*. By Theorem 2.3.1, $\mathrm{pr}(\mathfrak{g}_-) \cong \mathfrak{g}$, and so by Theorem 2.1.3,* $(2, 3, 5)$*-distributions can be described by a Cartan geometry of type (G_2, P_1).*

Generally, geometries with $\mathrm{pr}(\mathfrak{g}_-) \cong \mathfrak{g}$ have underlying structure a bracket-generating distribution. For 1-gradings, the filtration on M is trivial, so the geometry is determined by a reduction of structure group alone. These include conformal structures $(B_\ell/P_1, D_\ell/P_1)$, Segré (almost Grassmannian) structures (A_ℓ/P_k for $k \neq 1, \ell$), and almost spinorial structures (D_ℓ/P_ℓ). Parabolic geometries with contact gradings are determined by a non-trivial filtration as well as a reduction of structure group, e.g. CR and Lagrangean contact $(A_\ell/P_{1,\ell})$, and Lie contact structures $(B_\ell/P_2, D_\ell/P_2)$. In the two exceptional cases, A_ℓ/P_1 and C_ℓ/P_1, $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0)$ is infinite-dimensional. Algebraically, one needs to appropriately constrain the grading one component of $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0)$ through additional structure, namely projective and contact projective structures, respectively. We refer to the regular infinitesimal flag structure, or the structure reduction of $\mathrm{pr}_1(\mathfrak{g}_-, \mathfrak{g}_0)$ to $\mathfrak{g}_1$ in exceptional cases, as the *underlying structure* for a regular parabolic geometry.

Motivated by Lemma 2.1.5 and Theorem 2.3.1, we define a variant of Tanaka prolongation:

Definition 2.3.3. *Let $\mathfrak{g}$ be a $\mathbb{Z}$-graded Lie algebra, and $\mathfrak{a}_0 \subset \mathfrak{g}_0$ a subalgebra. Define the graded subalgebra $\mathfrak{a} \subset \mathfrak{g}$ by: (i) $\mathfrak{a}_{\leq 0} := \mathfrak{g}_{\leq 0}$; (ii) $\mathfrak{a}_k = \{X \in \mathfrak{g}_k \mid [X, \mathfrak{g}_{-1}] \subset \mathfrak{a}_{k-1}\}$ for $k > 0$. We will denote $\mathfrak{a} = \bigoplus_k \mathfrak{a}_k$ by $\mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{a}_0)$. (In particular, $\mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{g}_0) = \mathfrak{g}$.)*

When $\mathfrak{g}$ is simple and $\mathfrak{g}/\mathfrak{p}$ is not A_ℓ/P_1 or C_ℓ/P_1, then $\mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{a}_0)$ is the same as $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{a}_0)$.

---

[5] In the semisimple case, any simple ideal $\mathfrak{g}' \subset \mathfrak{g}$ contained in $\mathfrak{g}_0$ satisfies $[\mathfrak{g}', \mathfrak{g}_{-1}] = 0$. This violates Tanaka's axiom (T.2), so by convention, we exclude this case from all our considerations in this article.

2.4. **Regular, normal parabolic geometries.** There is an equivalence of categories between regular, normal parabolic geometries and underlying structures. We articulate normality below.

Using the P-equivariant isomorphism $(\mathfrak{g}/\mathfrak{p})^* \cong \mathfrak{p}_+$, consider the chain spaces $\bigwedge^k \mathfrak{p}_+ \otimes \mathfrak{g}$. The ($P$-equivariant) Kostant codifferential ∂^*, which is minus the Lie algebra homology differential, turns these chain spaces into a complex. A regular parabolic geometry is *normal* if the curvature function $\kappa : \mathcal{G} \to \bigwedge^2 (\mathfrak{g}/\mathfrak{p})^* \otimes \mathfrak{g}$ satisfies $\partial^* \kappa = 0$. While κ is a complete obstruction to local flatness, it is a rather complicated object. But for *normal* geometries, since $(\partial^*)^2 = 0$, we can define *harmonic curvature* $\kappa_H : \mathcal{G} \to \ker(\partial^*)/\mathrm{im}(\partial^*)$, which is a much simpler object and which is still a complete obstruction to local flatness. (As such, we say that $x \in M$ is a *non-flat point* if $\kappa_H(u) \neq 0$ for some (any) $u \in \pi^{-1}(x)$.) While κ_H is P-equivariant, P_+ acts trivially on $\ker(\partial^*)/\mathrm{im}(\partial^*)$, so κ_H descends to a G_0-equivariant function on $\mathcal{G}_0 = \mathcal{G}/P_+$. As G_0-modules $(\mathfrak{g}/\mathfrak{p})^* \cong \mathfrak{g}_-^* \cong \mathfrak{p}_+$, so consider the ($G_0$-equivariant) Lie algebra cohomology differential ∂ acting on the co-chain spaces $\bigwedge^k (\mathfrak{g}_-)^* \otimes \mathfrak{g}$, and the *Kostant Laplacian* $\Box := \partial\partial^* + \partial^*\partial$. By a lemma of Kostant, we have as G_0-modules

$$\bigwedge^k (\mathfrak{g}_-)^* \otimes \mathfrak{g} = \overbrace{\mathrm{im}(\partial^*) \oplus \rlap{$\underbrace{\phantom{\ker(\Box) \oplus \mathrm{im}(\partial)}}_{\ker(\partial)}$}\ker(\Box)}^{\ker(\partial^*)} \oplus \mathrm{im}(\partial), \qquad \ker(\Box) \cong \frac{\ker(\partial^*)}{\mathrm{im}(\partial^*)} \cong \frac{\ker(\partial)}{\mathrm{im}(\partial)} =: H^k(\mathfrak{g}_-, \mathfrak{g}).$$

By G_0-equivariancy, $\kappa_H : \mathcal{G}_0 \to H^2(\mathfrak{g}_-, \mathfrak{g})$ maps fibres of $\mathcal{G}_0 \to M$ to G_0-orbits in $H^2(\mathfrak{g}_-, \mathfrak{g})$. Kostant's theorem (Section 2.5) gives a complete description of the G_0-module structure of $H^2(\mathfrak{g}_-, \mathfrak{g})$.

The grading element Z acts on $H^2(\mathfrak{g}_-, \mathfrak{g}) \cong \ker(\Box) \subset \bigwedge^2 \mathfrak{g}_-^* \otimes \mathfrak{g}$. Let $H^2_+(\mathfrak{g}_-, \mathfrak{g})$ denote the submodules with positive homogeneity. Regularity is equivalent to $\mathrm{im}(\kappa_H) \subseteq H^2_+(\mathfrak{g}_-, \mathfrak{g})$ [10, Theorem 3.1.12]. This has the following important corollary which was mentioned in [8, Section 2.5]:

Proposition 2.4.1. *Let* $(\mathcal{G} \to M, \omega)$ *be a regular, normal parabolic geometry of type* (G, P)*. If* $\dim(\mathfrak{inf}(\mathcal{G}, \omega)) = \dim(\mathfrak{g})$*, then the geometry is locally flat.*

Proof. Given $0 \neq \phi \in H^2_+(\mathfrak{g}_-, \mathfrak{g})$, the grading element $Z \in \mathfrak{z}(\mathfrak{g}_0)$ satisfies $Z \notin \mathfrak{ann}(\phi) \subsetneq \mathfrak{g}_0$. Let $u \in \mathcal{G}$. We use two observations (see Section 4.1): (i) $\mathfrak{f}(u) := \mathrm{im}(\omega_u|_{\mathfrak{inf}(\mathcal{G},\omega)}) \subset \mathfrak{g}$ is a filtered Lie algebra with $\mathfrak{s}(u) := \mathrm{gr}(\mathfrak{f}(u))$ a graded subalgebra of $\mathfrak{g}$; and (ii) $\mathfrak{s}_0(u) \subseteq \mathfrak{ann}(\kappa_H(u))$. Hence, $\dim(\mathfrak{inf}(\mathcal{G}, \omega)) = \dim(\mathfrak{s}(u)) \leq \dim(\mathfrak{g})$. If equality holds, then $Z \in \mathfrak{s}_0(u) \subseteq \mathfrak{ann}(\kappa_H(u))$. Since $\kappa_H(u) \in H^2_+(\mathfrak{g}_-, \mathfrak{g})$, this forces $\kappa_H(u) = 0$. $\Box$

2.5. **Kostant's theorem.** Given $(\mathfrak{g}, \mathfrak{p})$ and an irreducible $\mathfrak{g}$-module $\mathbb{U}$, we define $\partial, \partial^*, \Box$, and $H^r(\mathfrak{g}_-, \mathbb{U})$ analogous to Section 2.4. Kostant's version of the Bott–Borel–Weil theorem [28], [1], [10] efficiently computes $H^r(\mathfrak{g}_-, \mathbb{U})$ as $\mathfrak{g}_0$-modules. Our main interest will be the $r = 2$ case.

Notation: Let $-\mu$ be a $\mathfrak{p}$-dominant weight. Let $\mathbb{V}_\mu$ be the irreducible $\mathfrak{g}_0$-module with *lowest* weight μ. Denote this by the Dynkin diagram notation for $-\mu$. The homogeneity of $\mathbb{V}_\mu$ is $Z(\mu)$.

Remark 2.5.1. A consequence of the seemingly perverse "minus lowest weight" convention [1] is that Kostant's algorithm below describes $H^2(\mathfrak{g}_-, \mathbb{U})$ (as opposed to $H^2(\mathfrak{p}_+, \mathbb{U})$).

If $\mathfrak{g}_0^{ss} \neq 0$ and $-\mu$ is $\mathfrak{p}$-dominant, the theorem of lowest (highest) weight says there is a unique (up to isomorphism) $\mathfrak{g}_0^{ss}$-representation with lowest weight μ_0, where μ_0 is obtained by deleting the crossed nodes from μ. (If $\mathfrak{g}_0^{ss} = 0$, take the trivial 1-dimensional representation.) We augment this to a $\mathfrak{g}_0$-module $\mathbb{V}_\mu$ by defining a $\mathfrak{z}(\mathfrak{g}_0)$-action: let Z_i act by the scalar $Z_i(\mu)$ for any $i \in I_\mathfrak{p}$. Note that if λ is $\mathfrak{g}$-dominant, then for any $w \in W^\mathfrak{p}$, $w \cdot \lambda$ is $\mathfrak{p}$-dominant (by definition of $W^\mathfrak{p}$).

Recipe 4 (**Kostant's theorem for $H^2(\mathfrak{g}_-,\mathbb{U})$**)**.** *Let $\mathfrak{g}$ be complex semisimple, $\mathfrak{p}\subset\mathfrak{g}$ a parabolic subalgebra, and $\mathbb{U}$ an irreducible $\mathfrak{g}$-module with highest weight λ. As a $\mathfrak{g}_0$-module,*

$$H^2(\mathfrak{g}_-,\mathbb{U}) \cong \bigoplus_{w\in W^{\mathfrak{p}}(2)} \mathbb{V}_{-w\cdot\lambda}.$$

For each $w=(jk)\in W^{\mathfrak{p}}(2)$, $\Phi_w=\{\alpha_j,\sigma_j(\alpha_k)\}$. Via the $\mathfrak{g}_0$-module isomorphism $H^2(\mathfrak{g}_-,\mathbb{U})\cong$ $\ker(\Box)\subset\bigwedge^2\mathfrak{g}_-^*\otimes\mathbb{U}$, the module $\mathbb{V}_{-w\cdot\lambda}$ has the unique (up to scale) lowest weight vector

$$\phi_0 := e_{\alpha_j}\wedge e_{\sigma_j(\alpha_k)}\otimes v, \tag{2.7}$$

where $e_\gamma\in\mathfrak{g}_\gamma$ are root vectors, and $v\in\mathbb{U}$ is a weight vector with weight $w(-\lambda)$.

Example 2.5.2 (G_2/P_1)**.** *In Example 2.2.2, since $\lambda_{\mathfrak{g}}=\lambda_2$ for $\mathfrak{g}=\mathrm{Lie}(G_2)$, we computed $H^2(\mathfrak{g}_-,\mathfrak{g})$ as a $\mathfrak{g}_0\cong\mathfrak{gl}_2(\mathbb{C})$ module. Since the lowest root of $\mathfrak{g}_1$ is $\alpha_1=2\lambda_1-\lambda_2$, then*

$\mathfrak{g}_1 =$ 

$\Rightarrow \quad H^2(\mathfrak{g}_-,\mathfrak{g})=\mathbb{V}_{-w\cdot\lambda_{\mathfrak{g}}} =$ 

$= \bigodot^4(\mathfrak{g}_1)=\bigodot^4(\mathfrak{g}_{-1})^*,$

where $w=(12)\in W^{\mathfrak{p}}(2)$. Since $\mathfrak{g}_1$ is the standard representation of $\mathfrak{g}_0$, then $H^2(\mathfrak{g}_-,\mathfrak{g})$ is the space of binary quartics on $\mathfrak{g}_{-1}$. This recovers Cartan's result [11] *concerning the fundamental (harmonic) curvature tensor for generic rank two distributions in dimension five. Furthermore,*

$$\Phi_w=\{\alpha_1,3\alpha_1+\alpha_2\},\qquad w(-\lambda_{\mathfrak{g}})=3\lambda_1-2\lambda_2=-\alpha_2.$$

Thus, $H^2(\mathfrak{g}_-,\mathfrak{g})\cong\ker(\Box)\subset\bigwedge^2\mathfrak{g}_-^*\otimes\mathfrak{g}$ has lowest *weight vector $\phi_0=e_{\alpha_1}\wedge e_{3\alpha_1+\alpha_2}\otimes e_{-\alpha_2}$.*

For $H^r(\mathfrak{g}_-,\mathbb{U})$, we make the obvious changes to Recipe 4, e.g. $w\in W^{\mathfrak{p}}(r)$, etc. If $\mathfrak{g}$ is simple, the highest weight $\lambda_{\mathfrak{g}}$ is also the highest root, so $\alpha=w(-\lambda_{\mathfrak{g}})\in\Delta$ and $v=e_\alpha\in\mathfrak{g}_\alpha$. If $\mathfrak{g}$ is semisimple, say with decomposition into simple ideals $\mathfrak{g}=\mathfrak{g}'\times\mathfrak{g}''$, then $H^2(\mathfrak{g}_-,\mathfrak{g})=H^2(\mathfrak{g}_-,\mathfrak{g}')\oplus H^2(\mathfrak{g}_-,\mathfrak{g}'')$, and we can apply Kostant's theorem to each factor. Note that in the semisimple case, $\mathfrak{g}_0=\mathfrak{g}_0'\times\mathfrak{g}_0''$ and writing $\mu=\mu'+\mu''$, we have $\mathbb{V}_\mu\cong\mathbb{V}_{\mu'}\boxtimes\mathbb{V}_{\mu''}$. Here, $\boxtimes$ is the "external" tensor product, so $\mathfrak{g}_0',\mathfrak{g}_0''$ act trivially on $\mathbb{V}_{\mu''},\mathbb{V}_{\mu'}$, respectively. More generally, when $\mathfrak{g}$ contains simple ideals $\mathfrak{g}'$ and $\mathfrak{g}''$, the $\mathfrak{g}_0$-irreducibles appearing $H^2(\mathfrak{g}_-,\mathfrak{g})$ come in three types (see Table 4). (The analysis for the gap problem in the general (complex or split-real) semisimple case will reduce to the case with no more than two factors – see Remark 4.3.9.)

Label	$w\in W^{\mathfrak{p}}(2)$	$w\cdot\lambda_{\mathfrak{g}'}$	Submodule of...
Type I	$(j'k')\in W^{\mathfrak{p}'}(2)$	$(j'k')\cdot\lambda_{\mathfrak{g}'}$	$H^2(\mathfrak{g}_-',\mathfrak{g}')\boxtimes\mathbb{C}$
Type II	$(j'k'')\in W^{\mathfrak{p}'}(1)\times W^{\mathfrak{p}''}(1)$	$\sigma_{j'}\cdot\lambda_{\mathfrak{g}'}+\sigma_{k''}\cdot 0''$	$H^1(\mathfrak{g}_-',\mathfrak{g}')\boxtimes\mathfrak{g}_1''$
Type III	$(j''k'')\in W^{\mathfrak{p}''}(2)$	$\lambda_{\mathfrak{g}'}+(j''k'')\cdot 0''$	$\mathfrak{g}_{-\nu'}'\boxtimes H^2(\mathfrak{g}_-'',\mathbb{C})$

Table 4. Types of $\mathfrak{g}_0$-submodules in $H^2(\mathfrak{g}_-,\mathfrak{g})$ when $\mathfrak{g}$ contains simple ideals $\mathfrak{g}'$ and $\mathfrak{g}''$

2.6. **Regularity and Yamaguchi's rigidity theorem.** Let $\mathfrak{g}$ be complex semisimple with a simple ideal having highest weight λ. Let us write the regularity condition $Z(-w\cdot\lambda)\geq 1$ in detail. Let $w=(jk)\in W^{\mathfrak{p}}(2)$, so $\Phi_w:=w\Delta^-\cap\Delta^+=\{\alpha_j,\sigma_j(\alpha_k)\}$. Using (2.5), (2.6), and writing

$r_a = \langle \lambda, \alpha_a^\vee \rangle \geq 0$ (since λ is $\mathfrak{g}$-dominant),

$$w \cdot \lambda = w(\lambda) + w \cdot 0 = w(\lambda) - \sum_{\alpha \in \Phi_w} \alpha = \sigma_j(\sigma_k(\lambda)) - \alpha_j - \sigma_j(\alpha_k) \tag{2.8}$$

$$= \sigma_j(\lambda - r_k\alpha_k) - \alpha_j - \sigma_j(\alpha_k) = \lambda - (r_j+1)\alpha_j - (r_k+1)(\alpha_k - c_{kj}\alpha_j).$$

From Recipe 3, we know $j \in I$, so $Z(w \cdot \lambda) \leq -1$ is equivalent to

$$Z(\lambda) \leq r_j + (r_k+1)(Z(\alpha_k) - c_{kj}). \tag{2.9}$$

Definition 2.6.1. *Let $\mathfrak{g}$ be complex simple with highest weight $\lambda_\mathfrak{g}$. Define*

$$W_+^\mathfrak{p}(2) := \{w \in W^\mathfrak{p}(2) \mid Z(-w \cdot \lambda_\mathfrak{g}) \geq 1\} \quad \Rightarrow \quad H_+^2(\mathfrak{g}_-, \mathfrak{g}) = \bigoplus_{w \in W_+^\mathfrak{p}(2)} \mathbb{V}_{-w \cdot \lambda_\mathfrak{g}}.$$

Example 2.6.2. *Condition* (2.9) *can be used to calculate $W_+^\mathfrak{p}(2)$. Consider $A_\ell/P_{1,2,s,t}$ for $4 \leq s < t < \ell$. Since $\lambda_\mathfrak{g} = \lambda_1 + \lambda_\ell = \alpha_1 + ... + \alpha_\ell$, then $Z(\lambda_\mathfrak{g}) = 4$. Since $r_j, r_k, Z(\alpha_k), -c_{kj} \in \{0, 1\}$, then* (2.9) *forces $r_k = Z(\alpha_k) = -c_{kj} = 1$, so $k = 1$, $j = 2$, and $W_+^\mathfrak{p}(2) = \{(21)\}$, while $|W^\mathfrak{p}(2)| = 11$.*

For regular, normal G/P geometries, $\mathrm{im}(\kappa_H) \subset H_+^2(\mathfrak{g}_-, \mathfrak{g})$. If $H_+^2(\mathfrak{g}_-, \mathfrak{g}) = 0$, then $\kappa_H \equiv 0$, so the geometry is locally flat. These geometries were classified by Yamaguchi [45], [46] so we refer to them as *Yamaguchi-rigid*.[6] For the gap problem, we need only study *Yamaguchi-nonrigid* geometries.

Theorem 2.6.3 (Yamaguchi's rigidity theorem)**.** *Let $\mathfrak{g}$ be a complex simple Lie algebra and $\mathfrak{p}$ a parabolic subalgebra. Then $H_+^2(\mathfrak{g}_-, \mathfrak{g}) \neq 0$ if and only if $(\mathfrak{g}, \mathfrak{p})$ is: (i) 1-graded, (ii) a contact gradation, or (iii) listed in Table 5.*

G	Range	2-graded	3-graded	4-graded	5-graded	6-graded
A_ℓ	$\ell \geq 4$	$P_{1,s}, P_{2,s}, P_{s,s+1}$	$P_{1,2,s}, P_{1,s,\ell}$	$P_{1,2,s,t}$	–	–
	$\ell = 3$	$P_{1,2}$	$P_{1,2,3}$	–	–	–
B_ℓ	$\ell \geq 4$	P_3, P_ℓ	$P_{1,2}$	$P_{2,3}$	–	–
	$\ell = 3$	P_3	$P_{1,2}, P_{1,3}$	$P_{2,3}$	$P_{1,2,3}$	–
	$\ell = 2$	–	$P_{1,2}$	–	–	–
C_ℓ	$\ell \geq 4$	$P_2, P_{\ell-1}$	$P_{1,\ell}, P_{2,\ell}, P_{\ell-1,\ell}$	$P_{1,2}$	$P_{1,2,\ell}$	$P_{1,2,s}$ $(s < \ell)$
	$\ell = 3$	P_2	$P_{1,3}, P_{2,3}$	$P_{1,2}$	$P_{1,2,3}$	–
D_ℓ	$\ell \geq 5$	$P_3, P_{1,\ell}$	$P_{1,2}$	$P_{2,3}, P_{1,2,\ell}$	–	–
	$\ell = 4$	$P_{1,4}$	$P_{1,2}$	$P_{1,2,4}$	–	–
G_2	–	–	P_1	–	$P_{1,2}$	–

Table 5. Yamaguchi-nonrigid geometries, excluding 1-graded and parabolic contact geometries

2.7. **Correspondence and twistor spaces.**

We review correspondence and twistor space constructions from [7, 10]. Let G be a semisimple Lie group G, and $Q \subset P \subset G$ parabolic subgroups, so there is a projection $G/Q \to G/P$. Write the decompositions of $\mathfrak{g}$ corresponding to $\mathfrak{q}, \mathfrak{p}$ as

$$\mathfrak{g} = \mathfrak{q}_- \oplus \mathfrak{q}_0 \oplus \mathfrak{q}_+ = \mathfrak{p}_- \oplus \mathfrak{p}_0 \oplus \mathfrak{p}_+, \tag{2.10}$$

where (see Lemma 4.4.1 in [10]),

$$I_\mathfrak{p} \subset I_\mathfrak{q}, \qquad \mathfrak{p}_\pm \subset \mathfrak{q}_\pm, \qquad \mathfrak{q}_0 \subset \mathfrak{p}_0, \qquad \mathfrak{z}(\mathfrak{p}_0) \subset \mathfrak{z}(\mathfrak{q}_0). \tag{2.11}$$

Given a G/P geometry $(\mathcal{G} \to N, \omega)$, the *correspondence space* for $Q \subset P$ is $\mathcal{C}N = \mathcal{G}/Q$. This carries a canonical G/Q geometry $(\mathcal{G} \to \mathcal{C}N, \omega)$, and the fibers of $\mathcal{C}N \to N$ are diffeomorphic to P/Q.

---

[6] The list in [45] contains some minor errors, which were corrected in [46].

The vertical subbundle $V\mathcal{C}N \subset T\mathcal{C}N$ corresponds to the Q-submodule $\mathfrak{p}/\mathfrak{q} \subset \mathfrak{g}/\mathfrak{q}$, and the Cartan curvature $\kappa^{\mathcal{C}N}$ of $(\mathcal{G} \to \mathcal{C}N, \omega)$ satisfies $i_\xi \kappa^{\mathcal{C}N} = 0$ for any $\xi \in \Gamma(V\mathcal{C}N)$.

Conversely, given a G/Q geometry $(\mathcal{G} \to M, \omega)$, we may ask if there exists a G/P geometry $(\mathcal{G} \to N, \omega)$ such that M is locally isomorphic to $\mathcal{C}N$. If so, we call N the (local) *twistor space* for M. From above, a necessary condition is that the subbundle $VM \subset TM$ induced from $\mathfrak{p}/\mathfrak{q} \subset \mathfrak{g}/\mathfrak{q}$ must satisfy $i_\xi \kappa = 0$ for all $\xi \in \Gamma(VM)$. Locally, this is sufficient, and moreover $(\mathcal{G} \to N, \omega)$ is unique when P/Q is assumed to be connected [7] (or Corollary 4.4.1 in [10]). More convenient still is that it suffices to only check that $\mathrm{im}(\kappa_H)$ vanishes on $\mathfrak{p} \cap \mathfrak{q}_-$, cf. Theorem 3.3 in [7]).

Two additional nice properties of correspondence spaces include:

- $\mathfrak{inf}(\mathcal{G} \to N, \omega) = \mathfrak{inf}(\mathcal{G} \to \mathcal{C}N, \omega)$, provided P/Q is connected.
- $(\mathcal{G} \to N, \omega)$ is normal iff $(\mathcal{G} \to \mathcal{C}N, \omega)$ is normal.

In contrast to normality, regularity is not preserved in passing to a correspondence space.

Proposition 2.7.1. *Let G be a complex semisimple Lie group, and $Q \subset G$ a parabolic subgroup. Consider a G/Q geometry $(\mathcal{G} \to M, \omega)$ with $\mathrm{im}(\kappa_H) \subset \mathbb{V}_\mu$, where $\mu = -w \cdot \lambda$, with λ the highest weight of a simple ideal in $\mathfrak{g}$, and $w = (jk) \in W^{\mathfrak{q}}(2)$. Let* (a) $\mathfrak{p} := \mathfrak{p}_j$ *if $c_{kj} < 0$; or,* (b) $\mathfrak{p} := \mathfrak{p}_{j,k}$ *if $c_{kj} = 0$. Then $\mathfrak{q} \subset \mathfrak{p} \subset \mathfrak{g}$ and $\mathrm{im}(\kappa_H)$ consists of maps which vanish on $\mathfrak{p} \cap \mathfrak{q}_- \subset \mathfrak{p}_0$.*

Proof. By Recipe 3, $j \in I_{\mathfrak{q}}$, and $k \in (I_{\mathfrak{q}} \cup \mathcal{N}(j)) \backslash \{j\}$. From Recipe 4, the lowest $\mathfrak{q}_0$-weight vector in the $\mathfrak{q}_0$-module $\mathbb{V}_\mu$ is $\phi_0 = e_{\alpha_j} \wedge e_{\sigma_j(\alpha_k)} \otimes e_{w(-\lambda)} \in \bigwedge^2 \mathfrak{q}_-^* \otimes \mathfrak{g}$.

If $c_{kj} < 0$, take $\mathfrak{p} := \mathfrak{p}_j$. Then $\sigma_j(\alpha_k) = \alpha_k - c_{kj}\alpha_j$ has positive $Z_{I_{\mathfrak{p}}}$-grading, so $\phi_0 \in \mathfrak{p}_1 \wedge \mathfrak{p}_+ \otimes \mathfrak{g}$. Now recall that the $\mathfrak{q}_0$-module $\mathbb{V}_\mu$ is generated by applying $\mathfrak{q}_0$-raising operators to ϕ_0. Since $\mathfrak{q}_0 \subset \mathfrak{p}_0$, and $\mathfrak{p}_1 \wedge \mathfrak{p}_+ \otimes \mathfrak{g}$ is $\mathfrak{p}_0$-invariant (hence $\mathfrak{q}_0$-invariant), then $\mathbb{V}_\mu \subset \mathfrak{p}_1 \wedge \mathfrak{p}_+ \otimes \mathfrak{g}$. Thus, any map in $\mathrm{im}(\kappa_H) \subset \mathbb{V}_\mu$ vanishes on $\mathfrak{p}_0$ and hence on $\mathfrak{p} \cap \mathfrak{q}_- \subset \mathfrak{p}_0$.

If $c_{kj} = 0$, take $\mathfrak{p} := \mathfrak{p}_{j,k}$. Then $\sigma_j(\alpha_k) = \alpha_k$ has positive $Z_{I_{\mathfrak{p}}}$-grading, and proceed as before. ☐

Corollary 2.7.2. *Under the hypothesis of Proposition 2.7.1, if there exists a parabolic subgroup P with $Q \subset P \subset G$, and Lie algebra $\mathfrak{p}$ given as in (a) or (b), then $(\mathcal{G} \to M, \omega)$ admits a twistor space N with $(\mathcal{G} \to N, \omega)$ of type (G, P).*

Remark 2.7.3. If Q is connected, then by the subalgebra–subgroup correspondence, there exists a unique connected (parabolic) subgroup P such that $Q \subset P \subset G$ with $\mathrm{Lie}(P) = \mathfrak{p}$. This guarantees the existence of a twistor space, and will play a substantial simplifying step in the calculation of submaximal symmetry dimensions – see Remark 4.3.10.

2.8. **Annihilators.** One step towards understanding the gap problem is the following question: Given a *nonzero* $\phi \in \mathbb{V}_\mu$, where $-\mu$ is $\mathfrak{p}$-dominant, what is the maximal dimension of $\mathfrak{ann}(\phi) \subset \mathfrak{g}_0$? A well-known consequence of the Borel fixed point theorem is that the G_0-action on $\mathbb{P}(\mathbb{V}_\mu)$ has a unique (Zariski) closed orbit $\mathcal{O}$, namely the orbit of the lowest weight line $[\phi_0]$. This is the unique orbit of minimal dimension, so $\max_{0 \neq \phi \in \mathbb{V}_\mu} \dim(\mathfrak{ann}(\phi))$ is realized precisely when $[\phi] \in \mathcal{O}$. Let μ_0 be the weight of $\mathfrak{g}_0^{ss}$ obtained by deleting the crossed nodes from μ. In $\mathbb{P}(\mathbb{V}_\mu)$, the stabilizer in G_0^{ss} of $[\phi_0]$ is the (*opposite*) parabolic subgroup with crosses where $-\mu_0$ has positive coefficients.

Definition 2.8.1. *Let $J_\mu = \{j \notin I \mid \langle \mu, \alpha_j^\vee \rangle \neq 0\}$ (marked by asterisks on the Dynkin diagram), and $Z_{J_\mu} := \sum_{j \in J_\mu} Z_j$. Using the Z_{J_μ}-grading, let $\mathfrak{p}_\mu^{\mathrm{op}} = (\mathfrak{g}_0^{ss})_{\leq 0}$ be the (opposite) parabolic subalgebra of $\mathfrak{g}_0^{ss}$ corresponding to J_μ. If $\mathfrak{g}$ is simple, write $J_w := J_{-w \cdot \lambda_{\mathfrak{g}}}$ and $\mathfrak{p}_w^{\mathrm{op}} := \mathfrak{p}_\mu^{\mathrm{op}}$.*

Recipe 5. *Let* $\phi_0 \in \mathbb{V}_\mu$ *be the lowest weight vector. Then* $\mathfrak{a}_0 := \mathfrak{ann}(\phi_0) \subsetneq \mathfrak{z}(\mathfrak{g}_0) \oplus \mathfrak{p}_\mu^{\mathrm{op}}$, *where*

$$\mathfrak{a}_0 = \{H \in \mathfrak{h} \mid \mu(H) = 0\} \oplus \bigoplus_{\gamma \in \Delta(\mathfrak{g}_{0,\leq 0})} \mathfrak{g}_\gamma, \tag{2.12}$$

$$\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_\mu^{\mathrm{op}}) + |I_\mathfrak{p}| - 1 = \max_{0 \neq \phi \in \mathbb{V}_\mu} \dim(\mathfrak{ann}(\phi)). \tag{2.13}$$

(Use Recipe 1 to compute the dimension of $\mathfrak{p}_\mu^{\mathrm{op}} \subset \mathfrak{g}_0^{ss}$*.)*

Example 2.8.2 (E_8/P_8)**.** *This has* $\lambda_\mathfrak{g} = \lambda_8$ *and* $W_+^\mathfrak{p}(2) = \{w := (87)\}$*. We have* $\dim(\mathfrak{g}_0) = 1 + \dim(E_7) = 134$, *so* $\dim(\mathfrak{g}_-) = \frac{1}{2}(\dim(E_8) - \dim(\mathfrak{g}_0)) = \frac{1}{2}(248 - 134) = 57$*. Moreover,*

$w \cdot \lambda_\mathfrak{g} =$  $\Rightarrow \quad J_w = \{6, 7\}.$

By Recipe 5, $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_{6,7} \subset E_7$ *with* $\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_w^{\mathrm{op}}) = \frac{1}{2}(\dim(E_7) + 2 + \dim(D_5)) = 90$.

Example 2.8.3 (B_ℓ/P_1 for $\ell \geq 4$)**.** *By Recipe 1,* $\mathfrak{g}_0 \cong B_{\ell-1} \times \mathbb{C}$, *and by Kostant (Recipe 4),*

$H^2(\mathfrak{g}_-, \mathfrak{g}) = \mathbb{V}_\mu =$  $, \qquad w = (12), \qquad Z(\mu) = +2.$

Removing the crossed node, $\mathfrak{p}_\mu^{\mathrm{op}} \cong \mathfrak{p}_2 \subset B_{\ell-1}$, *and* $(\mathfrak{g}_0^{ss})_0 = \mathbb{C} \times A_1 \times B_{\ell-3}$*. By Recipes 5 and 1,*

$$\begin{aligned} \dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_\mu^{\mathrm{op}}) &= \frac{1}{2}(\dim(B_{\ell-1}) + \dim((\mathfrak{g}_0^{ss})_0)) \\ &= \frac{1}{2}((\ell-1)(2\ell-1) + 1 + 3 + (\ell-3)(2\ell-5)) = 2\ell^2 - 7\ell + 10. \end{aligned}$$

In terms of $H^2(\mathfrak{g}_-, \mathfrak{g}) \cong \ker(\Box) \subset \bigwedge^2 \mathfrak{g}_-^* \otimes \mathfrak{g}$, *we have* $\phi_0 = e_{\alpha_1} \wedge e_{\alpha_1 + \alpha_2} \otimes e_{-\alpha_2 - 2\alpha_3 - \ldots - 2\alpha_\ell}$.

We describe $\mathfrak{a}_0$ *explicitly. Taking* $\mathbb{C}^{2\ell+1}$ *with standard anti-diagonal symmetric* $\mathbb{C}$*-bilinear form* g, $\mathfrak{so}(2\ell+1, \mathbb{C})$ *consists of* g*-skew matrices. Define* $\epsilon_i \in \mathfrak{h}^*$ *by* $\epsilon_i(\mathrm{diag}(a_1, ..., a_\ell, 0, -a_\ell, ..., -a_1)) = a_i$*. The simple roots are* $\alpha_1 = \epsilon_1 - \epsilon_2$, ..., $\alpha_{\ell-1} = \epsilon_{\ell-1} - \epsilon_\ell$, $\alpha_\ell = \epsilon_\ell$*. Hence,*

$$\mu = 4\lambda_1 - 2\lambda_3 = -2\alpha_1 + 2\alpha_3 + \ldots + 2\alpha_{k-1} + 2\alpha_k = -2(\epsilon_1 - \epsilon_2 - \epsilon_3).$$

By Recipe 5, this data completely determines $\mathfrak{a}_0$*, e.g. if* $\ell = 4$*, then*

$$\mathfrak{a}_0 = \left(\begin{array}{c|ccccccc|c} a+d & & & & & & & & 0 \\ \hline & a & b & & & & & 0 & \\ & c & d & & & & 0 & & \\ & z_1 & z_2 & z_3 & z_4 & 0 & & & \\ & z_5 & z_6 & z_7 & 0 & -z_4 & & & \\ & z_8 & z_9 & 0 & -z_7 & -z_3 & & & \\ & z_{10} & 0 & -z_9 & -z_6 & -z_2 & -d & -b & \\ & 0 & -z_{10} & -z_8 & -z_5 & -z_1 & -c & -a & \\ \hline 0 & & & & & & & & -a-d \end{array}\right).$$

(The condition $\mu(H) = 0$ *determines the diagonal, and* $\mathfrak{a}_0$ *contains* $\mathfrak{g}_\gamma \subset \mathfrak{g}_0$ *iff* $Z_{J_w}(\gamma) \leq 0$*.)*

3. PROLONGATION ANALYSIS

This section is representation-theoretic, focusing on the variant of the Tanaka prolongation introduced in Definition 2.3.3. We earlier defined $\mathfrak{a}^\phi := \mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{ann}(\phi))$, and for regular, normal parabolic geometries, we will be interested in

$$\mathfrak{U} := \max\left\{\dim(\mathfrak{a}^\phi) \mid 0 \neq \phi \in H^2_+(\mathfrak{g}_-, \mathfrak{g})\right\}. \tag{3.1}$$

From the first line in the proof of Proposition 2.4.1, $\mathfrak{U} < \dim(\mathfrak{g})$. If $H^2_+(\mathfrak{g}_-, \mathfrak{g}) = \bigoplus_i \mathbb{V}_i$ is the decomposition into $\mathfrak{g}_0$-irreducibles, and $\phi = \sum_i \phi_i$, where $\phi_i \in \mathbb{V}_i$ for each i, then $\mathfrak{ann}(\phi) \subset \mathfrak{ann}(\phi_i)$. By Lemma 2.1.5, $\mathfrak{a}^\phi \subset \mathfrak{a}^{\phi_i}$, so $\mathfrak{U} = \max_i(\max_{0\neq\phi_i\in\mathbb{V}_i} \dim(\mathfrak{a}^{\phi_i}))$. Thus, it suffices to understand $\max_{0\neq\phi\in\mathbb{V}} \dim(\mathfrak{a}^\phi)$, where $\mathbb{V}$ is an irreducible $\mathfrak{g}_0$-module. When $\mathfrak{g}$ is complex semisimple, each such submodule in $H^2_+(\mathfrak{g}_-, \mathfrak{g})$ is of the form $\mathbb{V}_\mu$, where $\mu = -w \cdot \lambda$, with λ the highest weight of a simple ideal in $\mathfrak{g}$, and $w \in W^{\mathfrak{p}}(2)$ (so $w \cdot \lambda$ is $\mathfrak{p}$-dominant). So we have

$$\mathfrak{U} = \max_{\mathbb{V}_\mu \subset H^2_+(\mathfrak{g}_-,\mathfrak{g})} \mathfrak{U}_\mu, \qquad \mathfrak{U}_\mu := \max\left\{\dim(\mathfrak{a}^\phi) \mid 0 \neq \phi \in \mathbb{V}_\mu\right\}. \tag{3.2}$$

Define $\mathfrak{a}(\mu) := \mathfrak{a}^{\phi_0}$, where ϕ_0 is the lowest weight vector of $\mathbb{V}_\mu$. (When $\mathfrak{g}$ is complex simple, we will also use the notation $\mathfrak{a}(w) := \mathfrak{a}(-w \cdot \lambda_{\mathfrak{g}})$ for $w \in W^{\mathfrak{p}}(2)$.) After proving $\mathfrak{U}_\mu = \dim(\mathfrak{a}(\mu))$, we establish a Dynkin diagram recipe to compute $\mathfrak{U}_\mu$, and then study the structure of $\mathfrak{a}(\mu)$.

3.1. Maximizing the Tanaka prolongation.

Lemma 3.1.1. *Let G be a complex semisimple Lie group, and P a parabolic subgroup. Let $\mathbb{V}_\mu$ be an irreducible G_0-module, and $\phi_0 \in \mathbb{V}_\mu$ the lowest $\mathfrak{g}_0$-weight vector. Then for any* nonzero $\phi \in \mathbb{V}_\mu$, $\dim(\mathfrak{a}^\phi_k) \leq \dim(\mathfrak{a}^{\phi_0}_k)$, $\forall k \geq 0$. *Thus,* $\mathfrak{U}_\mu = \dim(\mathfrak{a}(\mu)) = \dim(\mathfrak{a}^{\phi_0})$. *Moreover,* $\dim(\mathfrak{a}^\phi) = \dim(\mathfrak{a}^{\phi_0})$ *iff* $[\phi] \in G_0 \cdot [\phi_0] \subset \mathbb{P}(\mathbb{V}_\mu)$.

Remark 3.1.2. In general, $\mathfrak{ann}(\phi) \not\subset \mathfrak{ann}(\phi_0)$, and $\mathfrak{a}^\phi_k \not\subset \mathfrak{a}^{\phi_0}_k$.

Proof of Lemma 3.1.1. If $\mathfrak{g}_0^{ss} = 0$, then irreducibility implies $\mathbb{V}_\mu \cong \mathbb{C}$, and the result is immediate. So suppose $\mathfrak{g}_0^{ss} \neq 0$. The $k = 0$ case was established in Section 2.8, so let $k \geq 1$. Define $\Psi_k : \mathbb{P}(\mathbb{V}_\mu) \to \mathbb{Z}_{\geq 0}$ by $\Psi_k([\phi]) = \dim(\mathfrak{a}^\phi_k)$. By Lemma 2.1.5,

$$\begin{aligned}\mathfrak{a}^\phi_k &= \mathrm{pr}^{\mathfrak{g}}_k(\mathfrak{g}_-, \mathfrak{ann}(\phi)) = \{X \in \mathfrak{g}_k \mid \mathrm{ad}^k_{\mathfrak{g}_{-1}}(X) \cdot \phi = 0\}\\ &= \{X \in \mathfrak{g}_k \mid (\mathrm{ad}_{Y_{i_1}} \circ \ldots \circ \mathrm{ad}_{Y_{i_k}}(X)) \cdot \phi = 0,\ \forall Y_{i_j} \in \mathfrak{g}_{-1}\}\end{aligned}$$

The function $(\mathrm{ad}_{Y_{i_1}} \circ \ldots \circ \mathrm{ad}_{Y_{i_k}}(X)) \cdot \phi$ is a multilinear function of $X, Y_{i_1}, \ldots, Y_{i_k}, \phi$. Its vanishing is determined by evaluating all $Y_{i_1}, \ldots, Y_{i_k}$ on all tuples of basis elements $\{e_i\}$ of $\mathfrak{g}_{-1}$. So there is a bilinear function $T(X, \phi)$ such that $\mathfrak{a}^\phi_k = \{X \in \mathfrak{g}_k \mid T(X, \phi) = 0\}$. Choosing any basis of $\mathfrak{g}_k$, there is a matrix $M(\phi)$ such that $X \in \mathfrak{a}^\phi_k$ if and only if its coordinate vector is in $\ker(M(\phi))$. Since the rank of a matrix is a lower semi-continuous function of its entries, and $M(\phi)$ depends linearly on ϕ, then $\mathrm{rank}(M(\phi))$ is lower semi-continuous in ϕ (and depends only on $[\phi]$). We have $\dim(\mathfrak{a}^\phi_k) = \dim(\mathfrak{g}_k) - \mathrm{rank}(M(\phi))$, so Ψ_k is upper semi-continuous.

Since $\mathbb{V}_\mu$ is irreducible and G_0^{ss} is non-trivial, then $\mathcal{O} = G_0 \cdot [\phi_0]$ is the *unique* (Zariski) closed G_0-orbit in $\mathbb{P}(\mathbb{V}_\mu)$. Consequently, $[\phi_0]$ is in the (Zariski) closure of *every* G_0-orbit in $\mathbb{P}(\mathbb{V}_\mu)$. For any nonzero $\phi \in \mathbb{V}_\mu$, there is a sequence $\{g_n\} \subset G_0$ such that $g_n \cdot [\phi] = [g_n \cdot \phi] \to [\phi_0]$ as $n \to \infty$. For any n, $\mathfrak{ann}(g_n \cdot \phi) = \mathrm{Ad}_{g_n}(\mathfrak{ann}(\phi))$, so $\mathfrak{a}^{g_n\cdot\phi}_k = \mathrm{Ad}_{g_n}(\mathfrak{a}^\phi_k)$, and hence Ψ_k is constant on G_0-orbits. Thus, $\Psi_k([\phi]) = \Psi_k([g_n \cdot \phi]) = \Psi_k(g_n \cdot [\phi]) \leq \Psi_k([\phi_0])$, by upper semi-continuity of Ψ_k.

The final statement follows since $\dim(\mathfrak{a}^\phi_0) = \dim(\mathfrak{a}^{\phi_0}_0)$ iff $[\phi] \in G_0 \cdot [\phi_0] \subset \mathbb{P}(\mathbb{V}_\mu)$. □

Henceforth, we study $\mathfrak{a}(\mu)$. In Recipe 5, we gave a description of $\mathfrak{a}_0$. In the next section, we show that the structure of $\mathfrak{a}_+$ can be easily determined.

3.2. **A Dynkin diagram recipe.**

Let $-\mu$ be a $\mathfrak{p}$-dominant weight.

Definition 3.2.1. *Define* $I_\mu := \{j \in I \mid \langle \mu, \alpha_j^\vee \rangle = 0\}$*. If* $\mathfrak{g}$ *is simple, write* $I_w := I_{-w\cdot\lambda_\mathfrak{g}}$.

> **Notation:** We augment our Dynkin diagram notation for μ by putting a *square* around I_μ crossed nodes, and an *asterisk* on J_μ nodes. (See Definition 2.8.1.) Refer to this as $\mathcal{D}(\mathfrak{g}, \mathfrak{p}, \mu)$.

Example 3.2.2 ($C_6/P_{1,2,5}$, $w = (21)$)**.** *We have* $\lambda_\mathfrak{g} = 2\lambda_1$*, and*

$w \cdot \lambda_\mathfrak{g} =$  , *i.e.* $I_w = \{1, 5\}$, $J_w = \{4\}$.

Definition 3.2.3. *Given* $S \subset \{1, ..., \ell\}$*, let* $Z_S = \sum_{s\in S} Z_s$ *and* $\widetilde{Z}_S = (Z_s)_{s\in S}$ *denote the corresponding total grading and multi-grading elements.*

Each $\mathfrak{g}_j$ decomposes into $\mathfrak{g}_0$-irreducibles using $\widetilde{Z}_I$. In turn, these decompose into $\mathfrak{g}_{0,0}$-irreducibles using $\widetilde{Z}_{J_\mu}$. Writing $1_i \in \mathbb{Z}^I$ for the tuple which is zero except for a 1 in the i-th position, we have

$$\begin{aligned} &\mathfrak{g}_0\text{-module decomposition:} \quad && \mathfrak{g}_1 = \bigoplus_{i\in I} \mathfrak{g}_{1_i}, \quad && \Delta(\mathfrak{g}_{1_i}) = \{\alpha \in \Delta(\mathfrak{g}_1) \mid Z_i(\alpha) = 1\}; \\ &\mathfrak{g}_{0,0}\text{-module decomposition:} \quad && \mathfrak{g}_{1_i} = \bigoplus_{A\geq 0} \mathfrak{g}_{1_i,A}, \quad && \Delta(\mathfrak{g}_{1_i,A}) = \{\alpha \in \Delta(\mathfrak{g}_{1_i}) \mid \widetilde{Z}_{J_\mu}(\alpha) = A\}. \end{aligned} \tag{3.3}$$

Here, $A \in \mathbb{Z}^{J_\mu}$ is a multi-index, and "$A \geq 0$" means *all* entries are non-negative. Note the unique lowest root in $\Delta(\mathfrak{g}_{1_i,0})$ is α_i.

Definition 3.2.4. *If* $\mathcal{S}_1, \mathcal{S}_2 \subset \Delta$*, define* $\mathcal{S}_1 \dot{+} \mathcal{S}_2 := \{\alpha + \beta \mid \alpha \in \mathcal{S}_1, \beta \in \mathcal{S}_2\} \cap \Delta$.

Lemma 3.2.5. *If* $-\mu$ *is a* $\mathfrak{p}$*-dominant weight with* $I_\mu \neq \emptyset$*, and* $\mathfrak{a} = \mathfrak{a}(\mu)$*, then*

- (a) $\mathfrak{a}_+$ *is a direct sum of root spaces;*
- (b) $\Delta(\mathfrak{a}_1) = \{\alpha \in \Delta(\mathfrak{g}_1) \mid Z_{I_\mu}(\alpha) = 1,\ Z_{J_\mu}(\alpha) = 0\}$*;*
- (c) $\Delta(\mathfrak{a}_j) \subseteq \Delta(\mathfrak{a}_{j-1}) \dot{+} \Delta(\mathfrak{a}_1) \subseteq \{\alpha \in \Delta(\mathfrak{g}_j) \mid Z_{I_\mu}(\alpha) = j,\ Z_{J_\mu}(\alpha) = 0\}$, $\forall j \geq 2$.

Proof. From Recipe 5, $\mathfrak{a}_0$ is a $\mathfrak{g}_{0,0}$-module (hence, an $\mathfrak{h}$-module). Since $\mathfrak{g}_-$ is a $\mathfrak{g}_{0,0}$-module, then so is each $\mathfrak{a}_k$. By Lemma 2.1.5, $\mathfrak{a}_k \subset \mathfrak{g}_k$, and all $\mathfrak{g}_\gamma \subset \mathfrak{g}_k$ are 1-dimensional, so (a) is proven.

Prior to calculating $\mathfrak{a}_1$, we recall a few facts. For any nonzero $\gamma \in \mathfrak{h}^*$, $\exists! H_\gamma \in \mathfrak{h}$ such that $\gamma(H) = B(H, H_\gamma)$, by nondegeneracy of the Killing form B. Hence, $\alpha(H_\gamma) = \gamma(H_\alpha) = B(H_\alpha, H_\gamma) =: \langle \alpha, \gamma \rangle$. Choosing root vectors $e_\gamma \in \mathfrak{g}_\gamma$ ($\gamma \in \Delta$), we have $[e_\gamma, e_{-\gamma}] = c_\gamma H_\gamma$, where $c_\gamma = B(e_\gamma, e_{-\gamma}) \neq 0$.

Now calculate $\mathfrak{a}_1 = \{X \in \mathfrak{g}_1 \mid [X, \mathfrak{g}_{-1}] \subset \mathfrak{a}_0\}$ by testing e_α for $\alpha \in \Delta(\mathfrak{g}_1)$. Let $\beta \in \Delta(\mathfrak{g}_1)$. Then recalling $\mathfrak{a}_0$ from (2.12), we have

- (i) $\beta = \alpha$: $[e_\alpha, e_{-\alpha}] = c_\alpha H_\alpha$. Thus, if $H_\alpha \notin \mathfrak{a}_0$, i.e. $\langle \mu, \alpha \rangle \neq 0$, then $\alpha \notin \Delta(\mathfrak{a}_1)$.
- (ii) $\beta \neq \alpha$: If $\alpha - \beta \notin \Delta$, then $[e_\alpha, e_{-\beta}] = 0$ (no constraints). Otherwise, if $\alpha - \beta \in \Delta$, then $[e_\alpha, e_{-\beta}] = f_{\alpha\beta} e_{\alpha-\beta}$, where $f_{\alpha\beta} \neq 0$. Thus, $\alpha - \beta \in \Delta(\mathfrak{g}_{0,+})$ iff $e_{\alpha-\beta} \notin \mathfrak{a}_0$ iff $\alpha \notin \Delta(\mathfrak{a}_1)$.

Defining $\mathcal{T}_1 = \{\alpha \in \Delta(\mathfrak{g}_1) \mid \langle \mu, \alpha^\vee \rangle \neq 0\}$ and $\mathcal{T}_2 = \{\alpha \in \Delta(\mathfrak{g}_1) \mid \exists \beta \in \Delta(\mathfrak{g}_1) \text{ with } \alpha - \beta \in \Delta(\mathfrak{g}_{0,+})\}$, we have $\Delta(\mathfrak{a}_1) = \Delta(\mathfrak{g}_1) \backslash (\mathcal{T}_1 \cup \mathcal{T}_2)$.

Now use the $\mathfrak{g}_{0,0}$-module decomposition (3.3). By Schur's lemma, for each $\mathfrak{g}_{1_i,A}$, either $\mathfrak{g}_{1_i,A} \subset \mathfrak{a}_1$ or $\mathfrak{a}_1 \cap \mathfrak{g}_{1_i,A} = 0$. Thus, it suffices to test the lowest root in $\Delta(\mathfrak{g}_{1_i,A})$. Fix $i \in I$ and suppose $A > 0$ (i.e. *at least* one entry of A is positive), and let $\alpha \in \Delta(\mathfrak{g}_{1_i,A})$ be the lowest root. Since $A > 0$, α is not the lowest root of $\Delta(\mathfrak{g}_{1_i})$, i.e. $\alpha \neq \alpha_i$. However, since $\mathfrak{g}_{1_i}$ is an irreducible $\mathfrak{g}_0$-module, there is a sequence of simple roots in $\Delta(\mathfrak{g}_0)$ (possibly repeated) such that

$$\alpha - \alpha_{j_1} - ... - \alpha_{j_m} = \alpha_i, \qquad j_k \in \{1, ..., \ell\}\backslash I, \qquad \ell = \text{rank}(\mathfrak{g}).$$

and $\alpha - \alpha_{j_1} - \ldots - \alpha_{j_k} \in \Delta$ for $1 \leq k \leq m$. In particular, $\beta = \alpha - \alpha_{j_1} \in \Delta(\mathfrak{g}_1)$. Since α is the lowest root of $\Delta(\mathfrak{g}_{1_i,A})$, then $j_1 \in J_\mu$ and $\alpha_{j_1} \in \Delta(\mathfrak{g}_{0,+})$. Hence, $\alpha \in \mathcal{T}_2$, so $\Delta(\mathfrak{g}_{1_i,A}) \subset \mathcal{T}_2$. Thus, $\Delta(\mathfrak{a}_1) = \Delta(\mathfrak{g}_1) \backslash (\mathcal{T}_1 \cup \mathcal{T}_2) \subset \bigcup_{i \in I} \Delta(\mathfrak{g}_{1_i,0})$. For the $A = 0$ case, the lowest root of $\Delta(\mathfrak{g}_{1_i,0})$ is α_i, and clearly $\alpha_i \notin \mathcal{T}_2$. We have $i \notin I_\mu$ iff $\alpha_i \in \mathcal{T}_1$. Thus, claim (b) follows from:

$$\Delta(\mathfrak{a}_1) = \bigcup_{i \in I_\mu} \Delta(\mathfrak{g}_{1_i,0}). \tag{3.4}$$

Let $j \geq 2$ and $\gamma \in \Delta(\mathfrak{a}_j) \subset \Delta(\mathfrak{g}_j)$. Then $\gamma = \alpha + \beta$ for some $\alpha \in \Delta(\mathfrak{g}_{j-1})$ and $\beta \in \Delta(\mathfrak{g}_1)$, since $\mathfrak{g}_1$ is bracket-generating in $\mathfrak{g}_+$. By definition, $\mathfrak{a}_j = \{X \in \mathfrak{g}_j \mid [X, \mathfrak{g}_{-1}] \subset \mathfrak{a}_{j-1}\}$, so $\alpha = \gamma - \beta \in \Delta(\mathfrak{a}_{j-1})$. Since $\mathfrak{a}$ is graded, then $[\mathfrak{a}_j, \mathfrak{a}_{-(j-1)}] \subset \mathfrak{a}_1$, so $\beta = \gamma - \alpha \in \Delta(\mathfrak{a}_1)$. Using (b), we obtain (c). □

Theorem 3.2.6. *If $\mathfrak{g}$ is complex semisimple, $\mathfrak{p} \subset \mathfrak{g}$ a parabolic subalgebra, $-\mu$ is a $\mathfrak{p}$-dominant weight with $I_\mu \neq \emptyset$, and $\mathfrak{a} = \mathfrak{a}(\mu)$, then*

- (a) *$\mathfrak{a}_1$ generates all brackets $\mathfrak{a}_j$ for $j \geq 1$.*
- (b) *$\Delta(\mathfrak{a}_j) = \{\alpha \in \Delta(\mathfrak{g}_j) \mid Z_{I_\mu}(\alpha) = j,\ Z_{J_\mu}(\alpha) = 0\}$ for $j \geq 1$.*

Proof. The base case $j = 1$ is trivial for (a), while for (b) it follows from part (b) of Lemma 3.2.5. Assume the induction hypothesis for both (a) and (b) for $1 \leq j \leq r$. By part (c) of Lemma 3.2.5,

$$\Delta(\mathfrak{a}_{r+1}) \subseteq \Delta(\mathfrak{a}_r) \dot{+} \Delta(\mathfrak{a}_1) \subseteq \{\alpha \in \Delta(\mathfrak{g}_{r+1}) \mid Z_{I_\mu}(\alpha) = r+1,\ Z_{J_\mu}(\alpha) = 0\} =: \mathcal{S}_{r+1}. \tag{3.5}$$

Conversely, let $\gamma \in \mathcal{S}_{r+1}$ and $\beta \in \Delta(\mathfrak{g}_1)$. If $\alpha = \gamma - \beta \in \Delta$, then $\alpha \in \Delta(\mathfrak{g}_r)$. Since $Z_{I \backslash I_\mu}(\gamma) = 0$, then $Z_{I_\mu}(\beta) = 1$ and $Z_{J_\mu}(\beta) = 0$, so $\beta \in \Delta(\mathfrak{a}_1)$. Thus, $Z_{I_\mu}(\alpha) = r$ and $Z_{J_\mu}(\alpha) = 0$, so by the induction hypothesis, $\alpha \in \Delta(\mathfrak{a}_r)$, and $\gamma \in \Delta(\mathfrak{a}_{r+1})$. Hence, the inclusions in (3.5) are equalities. □

To the data $(\mathfrak{g}, \mathfrak{p}, \mu)$ when $I_\mu \neq \emptyset$, we associate a *reduced geometry* $(\overline{\mathfrak{g}}, \overline{\mathfrak{p}})$ via the following recipe:

Recipe 6. *On $\mathcal{D}(\mathfrak{g}, \mathfrak{p}, \mu)$, remove all nodes corresponding to $I \backslash I_\mu$ and J_μ, and any edges connected to these nodes. In the resulting diagram, remove any connected components which do not contain an I_μ node. This defines the Dynkin diagram $\mathcal{D}(\overline{\mathfrak{g}}, \overline{\mathfrak{p}})$ for the reduced geometry $(\overline{\mathfrak{g}}, \overline{\mathfrak{p}})$.*

It is clear that the data $\Delta(\mathfrak{a}_+)$ for $(\mathfrak{g}, \mathfrak{p}, \mu)$, and $\Delta(\overline{\mathfrak{g}}_+)$ for $(\overline{\mathfrak{g}}, \overline{\mathfrak{p}})$ naturally correspond. We use Recipes 1, 5, and 6 to compute $\dim(\mathfrak{g}_-), \dim(\mathfrak{a}_0)$, and $\dim(\mathfrak{a}_+)$ respectively; their sum is $\dim(\mathfrak{a})$.

Example 3.2.7 (E_8/P_8)**.** *From 2.8.2, $I_w = \emptyset$, so $\mathfrak{a}_+ = 0$. Hence,* $\dim(\mathfrak{a}) = 57 + 90 = 147$.

Example 3.2.8 ($A_\ell/P_1 \times A_\ell/P_\ell$)**.** *Let us ornament indices in the first and second factors by primes and double primes, respectively. Consider the Type II case where $w = (1', \ell'')$, $\lambda' := \lambda_{1'} + \lambda_{\ell'}$ (i.e. the highest weight in the first A_ℓ factor) and $\mu = -w \cdot \lambda'$. Then $\mathbb{V}_\mu$ is*

$\mathbb{V}_{-\sigma_{1'} \cdot \lambda'} \boxtimes \mathbb{V}_{-\sigma_{\ell''} \cdot 0''} =$  .

Since $I_\mu = \emptyset$, then $\mathfrak{a}(\mu)$ has no positive part. We have $\mathfrak{p}_\mu^{\mathrm{op}} \cong \mathfrak{p}_{1',(\ell-1)'} \times \mathfrak{p}_{(\ell-1)''} \subset A_{\ell-1} \times A_{\ell-1}$, and

$$\begin{aligned}\dim(\mathfrak{a}) &= \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0) = \dim(\mathfrak{g}_-) + 1 + \dim(\mathfrak{p}_\mu^{\mathrm{op}}) \\ &= 2\ell + 1 + \frac{1}{2}(\ell^2 - 1 + 2 + (\ell-2)^2 - 1) + \frac{1}{2}(\ell^2 - 1 + 1 + (\ell-1)^2 - 1) \\ &= 2\ell + 1 + (\ell^2 - 2\ell + 2) + (\ell^2 - \ell) = 2\ell^2 - \ell + 3.\end{aligned}$$

Example 3.2.9 ($C_6/P_{1,2,5}$, $w = (21)$)**.** *Applying Recipe 6 to Example 3.2.2, we obtain:*



i.e. $\overline{\mathfrak{g}}/\overline{\mathfrak{p}} = A_1/P_1 \times C_3/P_2$.

Since $(\bar{\mathfrak{g}}, \bar{\mathfrak{p}})$ *is 2-graded,* $\dim(\bar{\mathfrak{g}}_1) = 5$, *and* $\dim(\bar{\mathfrak{g}}_2) = 3$, *then* $\dim(\mathfrak{a}_1) = 5$ *and* $\dim(\mathfrak{a}_2) = 3$. *Indeed,*

$$\begin{aligned} \Delta(\mathfrak{a}_1):& \quad \alpha_1, \quad \alpha_5, \quad \alpha_4+\alpha_5, \quad \alpha_5+\alpha_6, \quad \alpha_4+\alpha_5+\alpha_6;\\ \Delta(\mathfrak{a}_2):& \quad 2\alpha_5+\alpha_6, \quad \alpha_4+2\alpha_5+\alpha_6, \quad 2\alpha_4+2\alpha_5+\alpha_6. \end{aligned}$$

We have $\dim(\mathfrak{g}_-) = \frac{1}{2}(\dim(C_6) - 3 - \dim(A_2 \times A_1)) = \frac{1}{2}(78 - 3 - 11) = 32$. *Since* $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_1 \times A_1 \subset A_2 \times A_1$, *then* $\dim(\mathfrak{a}_0) = 2 + \dim(\mathfrak{p}_w^{\mathrm{op}}) = 11$. *Thus,* $\dim(\mathfrak{a}) = 32 + 11 + 8 = 51$.

In fact, this last example is rather exceptional – in Theorem 3.6.1, we show that if $\mathfrak{g}$ is simple, and when $w \in W_+^{\mathfrak{p}}(2)$, then for $\mathfrak{a}(w)$, we have $\mathfrak{a}_2 = 0$ almost always.

3.3. **Prolongation-rigidity.**

Definition 3.3.1. *If* $\mathfrak{g}$ *is complex semisimple and* $-\mu$ *is* $\mathfrak{p}$*-dominant, then* $(\mathfrak{g}, \mathfrak{p}, \mu)$ *(or* $(\mathfrak{g}, \mathfrak{p}, w)$ *with* $\mu = -w \cdot \lambda_{\mathfrak{g}}$ *if* $\mathfrak{g}$ *is simple) is* prolongation-rigid *(PR) if* $\mathfrak{a}_+^{\phi} = 0$ *for any* nonzero $\phi \in \mathbb{V}_\mu$. *Here,* μ *is not assumed to have positive homogeneity. Otherwise, it is* non-prolongation-rigid *(NPR). If* $\mathfrak{g}$ *is real or complex,* $(\mathfrak{g}, \mathfrak{p})$ *is* PR_+ *if* $\mathfrak{a}_+^{\phi} = 0$ *for any* nonzero $\phi \in H_+^2(\mathfrak{g}_-, \mathfrak{g})$. *Otherwise, it is* NPR_+.

Remark 3.3.2. The following observations are immediate:
1. $(\mathfrak{g}, \mathfrak{p})$ is PR_+ if and only if $(\mathfrak{g}, \mathfrak{p}, \mu)$ is PR for all $\mathbb{V}_\mu \subset H_+^2(\mathfrak{g}_-, \mathfrak{g})$.
2. Any Yamaguchi-rigid $(\mathfrak{g}, \mathfrak{p})$ has $H_+^2(\mathfrak{g}_-, \mathfrak{g}) = 0$, so is (vacuously) PR_+.
3. If $(\mathfrak{g}, \mathfrak{p})$ is real and is NPR_+, then its complexification $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{p}_{\mathbb{C}})$ will also be NPR_+.

Let $\mathfrak{g}$ be complex semisimple. An immediate consequence of (3.4) is:

Corollary 3.3.3. *If* $-\mu$ *is* $\mathfrak{p}$*-dominant, then* $(\mathfrak{g}, \mathfrak{p}, \mu)$ *is PR iff* $I_\mu = \emptyset$.

Equivalently, in terms of our augmented marked Dynkin diagram notation,

Recipe 7. $(\mathfrak{g}, \mathfrak{p}, \mu)$ *is PR iff* $\mathcal{D}(\mathfrak{g}, \mathfrak{p}, \mu)$ *has no squares.*

Example 3.3.4 (G_2/P_1)**.** *From Example 2.2.2,* $W_+^{\mathfrak{p}}(2) = \{(12)\}$, *and* $(12) \cdot \lambda_2 =$ 

. *The coefficient over the cross is nonzero, i.e. no squares, and so* G_2/P_1 *is* PR_+.

Example 3.3.5 $(F_4/P_{1,2})$**.** *Since* $F_4/P_{1,2}$ *is Yamaguchi-rigid (see Theorem 2.6.3), it is* PR_+. *However,* $w = (21) \in W^{\mathfrak{p}}(2) \backslash W_+^{\mathfrak{p}}(2)$ *gives* $w \cdot \lambda_1 =$ 

, *so the triple* $(\mathfrak{g}, \mathfrak{p}, w)$ *is NPR. Its reduced geometry is* A_1/P_1, *so* $\mathfrak{a} = \mathfrak{a}(w)$ *has 1-dimensional* $\mathfrak{a}_+ = \mathfrak{a}_1$.

Now restrict to $\mu = -w \cdot \lambda$, where λ is the highest weight of a simple ideal in $\mathfrak{g}$ and $w = (jk) \in W^{\mathfrak{p}}(2)$. Let $r_a := \langle \lambda, \alpha_a^\vee \rangle$ and $i \in I$. Then $i \in I_\mu$ is equivalent (using (2.8) and Recipe 3) to

$$0 = r_i - (r_j + 1)c_{ji} - (r_k + 1)(c_{ki} - c_{kj}c_{ji}), \tag{3.6}$$

where $i, j \in I$, and $k \in (I \cup \mathcal{N}(j)) \backslash \{j\}$. Note $r_a \geq 0$ for all a since λ is $\mathfrak{g}$-dominant. Hence,

- If $i = j$, then $c_{ji} = 2$ and $c_{kj} \leq 0$ (since $k \neq j$). So (3.6) implies $0 = r_j + 2 - (r_k + 1)c_{kj} \geq 2$, a contradiction. So assume $i \neq j$ below, i.e. $c_{ij}, c_{ji} \leq 0$.
- If $k \in \mathcal{N}(j) \backslash I$ or $i, j, k \in I$ are distinct, then $c_{ki}, c_{kj} \leq 0$, so (3.6) implies
$$0 = \underbrace{r_i}_{\geq 0} - \underbrace{(r_j + 1)}_{\geq 1} \underbrace{c_{ji}}_{\leq 0} - \underbrace{(r_k + 1)}_{\geq 1} \underbrace{(c_{ki} - c_{kj}c_{ji})}_{\leq 0} \quad \Rightarrow \quad r_i = c_{ji} = c_{ki} = 0.$$
- If $k = i$, then (3.6) implies $(r_i + 1)(2 - c_{ij}c_{ji}) = r_i - (r_j + 1)c_{ji} \geq 0$. Thus, $0 \leq c_{ij}c_{ji} \leq 2$. This forces $c_{ij}c_{ji} = 1$, so $c_{ij} = c_{ji} = -1$, and (3.6) reduces to $r_j = 0$.

Summarizing, (3.6) is equivalent to:

$$\text{(F.1)} \qquad \begin{cases} \text{(a)}: & |I| \geq 2, \quad i,j \in I \text{ distinct}, \quad k \in \mathcal{N}(j)\backslash I, \quad 0 = r_i = c_{ji} = c_{ki}; \quad \text{or}, \\ \text{(b)}: & |I| \geq 3, \quad i,j,k \in I \text{ distinct}, \quad 0 = r_i = c_{ji} = c_{ki}; \quad \text{or}, \\ \text{(c)}: & |I| \geq 2, \quad i,j \in I \text{ distinct}, \quad k = i, \quad c_{ij} = c_{ji} = -1, \quad r_j = 0. \end{cases}$$

Proposition 3.3.6. *Let $\mathfrak{g}$ be complex semisimple and λ the highest weight of a simple ideal in $\mathfrak{g}$. Let $w = (jk) \in W^{\mathfrak{p}}(2)$ and $\mu = -w \cdot \lambda$. Then $(\mathfrak{g}, \mathfrak{p}, \mu)$ is PR if: (a) $|I_{\mathfrak{p}}| = 1$, or (b) $I_{\mathfrak{p}} = \{j,k\}$ and $c_{jk} = 0$.*

Proof. In both cases, it is impossible to satisfy (F.1). ☐

Now, impose regularity (2.9) in each case to obtain:

$$\text{(F.2)} \qquad \begin{cases} \text{(a)}: & Z(\lambda) \leq r_j - (r_k+1)c_{kj}; \\ \text{(b)}: & Z(\lambda) \leq r_j + (r_k+1)(1-c_{kj}); \\ \text{(c)}: & Z(\lambda) \leq 2(r_i+1). \end{cases}$$

Example 3.3.7. *Suppose $\mathfrak{g} = \mathfrak{g}' \times \mathfrak{g}''$ is the decomposition into simple ideals, with $I_{\mathfrak{p}'} \neq \emptyset$. If $w' = (j'k') \in W_+^{\mathfrak{p}'}(2)$, then $(\mathfrak{g}, \mathfrak{p}, -w' \cdot \lambda_{\mathfrak{g}'})$ is NPR (provided $I_{\mathfrak{p}''} \neq \emptyset$). Namely, choosing $i \in I_{\mathfrak{p}''}$, we can satisfy both* (F.1)*(a) and* (F.2)*(a). The cohomology component corresponding to $(\mathfrak{g}, \mathfrak{p}, -w' \cdot \lambda_{\mathfrak{g}'})$ lies inside $H^2_+(\mathfrak{g}'_-, \mathfrak{g}') \boxtimes \mathbb{C}$, so $\mathfrak{a}^{\phi}_+ \supset \mathfrak{g}''_+$. Thus, in the general semisimple case, $(\mathfrak{g}, \mathfrak{p})$ is always NPR$_+$.*

If $\mathfrak{g}$ is *simple*, the NPR$_+$ condition on $(\mathfrak{g}, \mathfrak{p})$ is restrictive. In particular, since the highest weight $\lambda_{\mathfrak{g}}$ is also the highest root, then $\lambda_{\mathfrak{g}} = \sum_{a=1}^{\ell} m_a \alpha_a$ with $m_a \geq 1$, so we must have $|I| \leq Z(\lambda_{\mathfrak{g}})$.

Corollary 3.3.8. *Let $\mathfrak{g}$ be complex simple. Then $(\mathfrak{g}, \mathfrak{p})$ is PR$_+$ if:* (a) *$\mathfrak{g}$ is 1-graded;* (b) *$\mathfrak{g}$ has a contact gradation; or* (c) *$\mathfrak{g}$ is a Lie algebra of exceptional type.*

Proof. All Yamaguchi-rigid $(\mathfrak{g}, \mathfrak{p})$ are PR$_+$. By Proposition 3.3.6, consider those Yamaguchi-nonrigid $(\mathfrak{g}, \mathfrak{p})$ with $\mathfrak{p}$ neither maximal nor $\mathfrak{p} \neq \mathfrak{p}_{j,k}$ for $c_{jk} = 0$. Among 1-gradings and contact gradings, this leaves $A_2/P_{1,2}$ (see Theorem 2.3.1), and among Lie algebras of exceptional type, this leaves $G_2/P_{1,2}$ (see Theorem 2.6.3). Both fail (F.1). ☐

Hence, over $\mathbb{R}$ or $\mathbb{C}$, almost all known gap results, cf. Table 2, have been for PR$_+$ geometries. (We note $C_2/P_{1,2}$ also fails (F.1).) The only exception is pairs of 2nd order ODE ($A_3/P_{1,2}$) which is NPR$_+$ (Section 5.3). In the next section, we classify all NPR$_+$ $(\mathfrak{g}, \mathfrak{p})$ when $\mathfrak{g}$ is simple.

3.4. **Classical flag varieties.**

We now analyze (F.1) and (F.2) from Section 3.3 when G/P_I is a classical complex flag variety (so G is a classical complex *simple* Lie group), and $|I| \geq 2$. Our labelling (a), (b), (c) below corresponds to the cases in (F.1) and (F.2). Note that we only list G/P up to Dynkin diagram symmetries, e.g. $D_\ell/P_{\ell-1} \cong D_\ell/P_\ell$.

3.4.1. *B_ℓ or D_ℓ case.* Here, $c_{kj} \in \{0, -1\}$ iff $\mathfrak{g} = D_\ell$. Also, $0 \leq r_j + r_k \leq 1$ (since $j \neq k$) using[7]

$$\text{(3.7)} \qquad \lambda = \lambda_2 = \begin{cases} \alpha_1 + 2\alpha_2 + \ldots + 2\alpha_\ell, & \mathfrak{g} = B_\ell \quad (\ell \geq 3); \\ \alpha_1 + 2\alpha_2 + \ldots + 2\alpha_{\ell-2} + \alpha_{\ell-1} + \alpha_\ell, & \mathfrak{g} = D_\ell \quad (\ell \geq 4). \end{cases}$$

(a) Assume $c_{kj} = -2$, so $\mathfrak{g} = B_\ell$ and $(k,j) = (\ell-1, \ell)$. From (F.1), $c_{ki} = 0$, so $\ell \geq 4$, $r_j = r_k = 0$. From (F.2), $|I| = Z(\lambda) = 2$, which contradicts (3.7). Thus, $c_{kj} \in \{0, -1\}$. Then $r_j - (r_k+1)c_{kj} \leq 2$, so from (F.2), $|I| = Z(\lambda) = 2$ and exactly one of j,k is 2. If $\mathfrak{g} = B_\ell$ or $j = 2$, then $Z(\lambda) \geq 3$, so $\mathfrak{g} = D_\ell$ and $k = 2 \notin I$. Hence, $\boxed{D_\ell/P_{1,\ell},\ \ell \geq 5;\ w = (12);\ i = \ell}$

---

[7] We treat $B_2 = C_2$ together with the general C_ℓ case.

(b) Assume $\mathfrak{g} = B_\ell$. From (F.2), $|I| \geq 3$, so $Z(\lambda) \geq 5$. But since $0 \leq r_j + r_k \leq 1$ and $c_{kj} \leq -1$, then $r_j + (r_k+1)(1-c_{kj}) \geq 5$ only if $r_j = 0$, $r_k = 1$, $c_{kj} = -2$, and hence $(k, j) = (\ell-1, \ell)$. From (F.1), $c_{ki} = 0$, so $\ell \geq 4$, $r_j = r_k = 0$. From (F.2), $Z(\lambda) = 3$, a contradiction.

Thus, $\mathfrak{g} = D_\ell$. Assuming $r_j = 1$, then $r_k = 0$ and (F.2) implies $|I| = Z(\lambda) = 3$. But $j = 2 \in I$, so by (3.7), $Z(\lambda) \geq 4$, a contradiction. Hence, $r_j = 0$, and (F.2) implies $r_k = 1$, $c_{kj} = -1$, and $3 \leq |I| \leq Z(\lambda) \leq 4$. From (3.7), $I = \{1, 2, \ell\}$, so: $\boxed{D_\ell/P_{1,2,\ell},\ \ell \geq 5;\ w = (12);\ i = \ell}$

(c) Assume $r_i = 0$, so from (F.2), $|I| = Z_I(\lambda) = 2$. Using (3.7), this is impossible given $c_{ij} = -1$ by (F.1). Thus, $r_i = 1$, and $2 \leq |I| \leq Z_I(\lambda) \leq 4$. Keeping in mind $c_{ij} = -1$, we have:

- $\boxed{B_\ell/P_{1,2},\ \ell \geq 3;\ w = (12);\ i = 2}$ and $\boxed{D_\ell/P_{1,2},\ \ell \geq 4;\ w = (12);\ i = 2}$
- $\boxed{B_\ell/P_{2,3},\ \ell \geq 4;\ w = (32);\ i = 2}$ and $\boxed{D_\ell/P_{2,3},\ \ell \geq 4;\ w = (32);\ i = 2}$
- $\boxed{D_\ell/P_{1,2,\ell},\ \ell \geq 5;\ w = (12);\ i = 2}$. Also, $\boxed{D_4/P_{1,2,4};\ w = (12) \text{ or } (42);\ i = 2}$

3.4.2. *C_ℓ case.* Here, $c_{kj} \in \{0, -1, -2\}$ with $c_{kj} = -2$ iff $(k, j) = (\ell, \ell-1)$. Also, $0 \leq r_j + r_k \leq 2$,

$$\lambda = 2\lambda_1 = 2\alpha_1 + \dots + 2\alpha_{\ell-1} + \alpha_\ell \quad (\ell \geq 2). \tag{3.8}$$

(a) Assuming $c_{kj} = -2$, then $(k, j) = (\ell, \ell-1)$. Since i, j, k are distinct, $c_{ji} = 0$, $r_i = 0$, then $\ell \geq 5$ and $r_j = r_k = 0$. From (F.2), $|I| = Z(\lambda) = 2$, but $i, j \in I\backslash\{\ell\}$ (distinct) implies $Z(\lambda) \geq 4$ (contradiction). Thus, $c_{kj} = -1$. From (F.2), (3.8), $|I| = 2$, $Z(\lambda) = 3$ with $j = 1$ or $k = 1$.
   (a) $(j, k) = (1, 2)$: Since $c_{ki} = 0$, then $\boxed{C_\ell/P_{1,\ell},\ \ell \geq 4;\ w = (12);\ i = \ell}$.
   (b) $(k, j) = (1, 2)$: Since $c_{ji} = 0$, then $\boxed{C_\ell/P_{2,\ell},\ \ell \geq 4;\ w = (21);\ i = \ell}$.

(b) Assume $c_{kj} = -2$, so $(k, j) = (\ell, \ell-1)$ and since $r_i = c_{ji} = c_{ki} = 0$, then $2 \leq i \leq \ell - 3$, and $r_j = r_k = 0$. From (F.2), $|I| = Z(\lambda) = 3$, but then $Z(\lambda) \geq 5$ by (3.8) (contradiction). Similarly, $c_{kj} = 0$ yields $|I| = Z(\lambda) = 3$ and the same contradiction. Thus,

$$c_{kj} = -1, \qquad 3 \leq |I| \leq Z(\lambda) \leq r_j + 2(r_k + 1),$$

with either $(j, k) = (1, 2)$ or $(2, 1)$. In the former case, $Z(\lambda) \leq 4$, but for $|I| \geq 3$, $Z(\lambda) \geq 5$, a contradiction. Thus, $(j, k) = (2, 1)$ and $Z(\lambda) \leq 6$, so $|I| = 3$. Keeping in mind $c_{ji} = c_{ki} = 0$, we have: $\boxed{C_\ell/P_{1,2,s}, \text{ with } 4 \leq s \leq \ell,\ w = (21),\ i = s}$.

(c) Since $c_{ij} = -1$, then $\ell \geq 3$. From (F.2) and (3.8), we cannot have $r_i = 0$. Thus, $r_i = 2$, $i = 1$, $2 \leq |I| \leq Z(\lambda) \leq 6$ and hence $|I| \leq 3$. Since $c_{ij} = c_{ji} = -1$, then $j = 2$. Two possibilities:
   (a) $\boxed{C_\ell/P_{1,2},\ \ell \geq 3;\ w = (21);\ i = 1}$
   (b) $\boxed{C_\ell/P_{1,2,s},\ 3 \leq s \leq \ell;\ w = (21);\ i = 1}$

3.4.3. *A_ℓ case.* Here, $c_{kj} \in \{0, -1\}$ and $0 \leq r_j + r_k \leq 2$ (since $j \neq k$) using

$$\lambda = \lambda_1 + \lambda_\ell = \alpha_1 + \dots + \alpha_\ell \quad (\ell \geq 1).$$

(1) We have $\ell \geq 3$ and $c_{kj} = -1$, so from (F.1), $2 \leq |I| = Z_I(\lambda) \leq r_j + r_k + 1$.
   (i) $r_j = 1$: We may take $j = 1$, so $k = 2 \notin I$, $r_k = 0$, and $|I| = Z_I(\lambda) = 2$. Since $r_i = 0$ and $c_{ji} = c_{ki} = 0$, then $4 \leq i \leq \ell - 1$. Thus, $\boxed{A_\ell/P_{1,s},\ 4 \leq i \leq \ell - 1;\ w = (12);\ i = s}$.
   (ii) $r_j = 0$: We have $2 \leq |I| = Z_I(\lambda) \leq 1 + r_k \leq 2$, so $r_k = 1$ and $|I| = 2$. We may take $k = 1 \notin I$, and $j = 2 \leq \ell - 1$. Since $c_{ji} = 0$, then: $\boxed{A_\ell/P_{2,s},\ 4 \leq s \leq \ell - 1;\ w = (21);\ i = s}$.

(2) From (F.2), $3 \leq |I| = Z_I(\lambda) \leq r_j + (r_k + 1)(1 - c_{kj})$.
   (a) $c_{kj} = 0$: We have $\ell \geq 3$ and $|I| = Z_I(\lambda) = 3$ with $r_j = r_k = 1$, so we may take $(j, k) = (1, \ell)$. Recalling $c_{ji} = c_{ki} = 0$, we have: $\boxed{A_\ell/P_{1,s,\ell},\ 3 \leq s \leq \ell - 2;\ w = (1, \ell);\ i = s}$.
   (b) $c_{kj} = -1$: We cannot have $r_j = r_k = 0$, nor $r_j = r_k = 1$ (since $\ell \geq 3$). Recalling $r_i = c_{ji} = c_{ki} = 0$,

(i) $(r_j, r_k) = (1,0)$: $|I| = 3$, $\boxed{A_\ell/P_{1,2,s},\ 4 \le s \le \ell - 1;\ w = (12);\ i = s}$

(ii) $(r_j, r_k) = (0,1)$:

(A) $|I| = 3$, $\boxed{A_\ell/P_{1,2,s},\ 4 \le s \le \ell - 1;\ w = (21);\ i = s}$.

(B) $|I| = 4$, $\boxed{A_\ell/P_{1,2,s,t},\ 4 \le s < t \le \ell;\ w = (21);\ i = s \text{ or } i = t \le \ell - 1}$.

(3) From (F.1) and (F.2), $c_{ij} = c_{ji} = -1$, $r_j = 0$, $2 \le |I| = Z_I(\lambda) \le 2(r_i + 1)$.

(a) $r_i = 0$: $|I| = 2$, $\boxed{A_\ell/P_{s,s+1},\ 2 \le s \le \ell - 2;\ w = (jk),\ i = k \text{ with } \{j,k\} = \{s, s+1\}}$.

(b) $r_i = 1$: $2 \le |I| \le 4$.

(i) $|I| = 2$: $\boxed{A_\ell/P_{1,2},\ \ell \ge 3;\ w = (21);\ i = 1}$

(ii) $|I| = 3$: $\boxed{A_\ell/P_{1,2,s},\ 3 \le s \le \ell;\ w = (21);\ i = 1}$

(iii) $|I| = 4$: $\boxed{A_\ell/P_{1,2,s,t},\ 3 \le s < t \le \ell;\ w = (21);\ i = 1}$

We summarize our classification in Table 6. (More detail is given in Appendix B.)

G	P	Range
A_ℓ	$P_{1,2}$	$\ell \ge 3$
	$P_{1,2,s}$	$3 \le s \le \ell$
	$P_{1,2,s,t}$	$3 \le s < t \le \ell$
	$P_{1,s,\ell}$	$3 \le s \le \ell - 2$
	$P_{i,i+1}$	$2 \le i \le \ell - 2$
	$P_{1,i}$	$4 \le i \le \ell - 1$
	$P_{2,i}$	$4 \le i \le \ell - 1$
B_ℓ	$P_{1,2}$	$\ell \ge 3$
	$P_{2,3}$	$\ell \ge 4$

G	P	Range
C_ℓ	$P_{1,2}$	$\ell \ge 3$
	$P_{1,\ell}$	$\ell \ge 4$
	$P_{2,\ell}$	$\ell \ge 4$
	$P_{1,2,s}$	$3 \le s \le \ell$
D_ℓ	$P_{1,2}$	$\ell \ge 4$
	$P_{1,\ell}$	$\ell \ge 5$
	$P_{2,3}$	$\ell \ge 5$
	$P_{1,2,\ell}$	$\ell \ge 4$

TABLE 6. Classification of NPR$_+$ geometries G/P with G simple

3.5. **Nested parabolic subalgebras.** We use the notation of Section 2.7. For nested parabolic subalgebras $\mathfrak{q} \subset \mathfrak{p} \subset \mathfrak{g}$ (so $I_\mathfrak{p} \subset I_\mathfrak{q}$), we have (2.10) and (2.11).

Theorem 3.5.1. *Let $\mathfrak{g}$ be complex semisimple. Let $\mathfrak{q} \subset \mathfrak{p} \subset \mathfrak{g}$ be parabolic subalgebras, and let λ be the highest weight of a simple ideal in $\mathfrak{g}$. Let $w \in W^\mathfrak{p}(2) \subset W^\mathfrak{q}(2)$ and $\mu = -w \cdot \lambda$. Choosing root vectors in $\mathfrak{g}$, define ϕ_0 as in (2.7). Let $\mathfrak{a} = \mathrm{pr}^\mathfrak{g}(\mathfrak{p}_-, \mathfrak{ann}_{\mathfrak{p}_0}(\phi_0))$ and $\mathfrak{b} = \mathrm{pr}^\mathfrak{g}(\mathfrak{q}_-, \mathfrak{ann}_{\mathfrak{q}_0}(\phi_0))$, which are respectively $Z_{I_\mathfrak{p}}$-graded and $Z_{I_\mathfrak{q}}$-graded. Then as $Z_{I_\mathfrak{p}}$-graded Lie algebras, $\mathfrak{a} = \mathfrak{b} \subset \mathfrak{g}$.*

Proof. Write $w = (jk) \in W^\mathfrak{p}(2) \subset W^\mathfrak{q}(2)$, so by Recipe 3, $k \neq j \in I_\mathfrak{p}$, and if $c_{kj} = 0$, then $k \in I_\mathfrak{p}$. We always have $\mathfrak{q} \subset \mathfrak{p} \subset \mathfrak{p}_j \subset \mathfrak{g}$, but if $c_{kj} = 0$, we also have $\mathfrak{q} \subset \mathfrak{p} \subset \mathfrak{p}_{j,k} \subset \mathfrak{g}$. Hence, it suffices to consider either $\mathfrak{p} = \mathfrak{p}_j$ if $c_{kj} < 0$, or $\mathfrak{p} = \mathfrak{p}_{j,k}$ if $c_{kj} = 0$. We assume this below.

By Proposition 3.3.6, $(\mathfrak{g}, \mathfrak{p}, \mu)$ is PR, so $\mathfrak{a} = \mathfrak{p}_- \oplus \mathfrak{a}_0$. Thus, $I^\mathfrak{p}_\mu = \emptyset$. We know $\mathfrak{p}_- \subset \mathfrak{q}_-$, and by Recipe 5, $\ker(\mu) \subset \mathfrak{a}_0 \cap \mathfrak{b}_0$. Let us show $\Delta(\mathfrak{a}_0) \subset \Delta(\mathfrak{b})$. Given $\alpha \in \Delta(\mathfrak{a}_0)$, we have $Z_{I_\mathfrak{p}}(\alpha) = 0$ and $Z_{J^\mathfrak{p}_\mu}(\alpha) \le 0$. Since $\mathfrak{q}_- \oplus \mathfrak{q}_{0,\le 0} \subset \mathfrak{b}$, we may assume $\alpha \in \Delta(\mathfrak{q}_{0,+} \oplus \mathfrak{q}_+) \subset \Delta^+$, hence $Z_{J^\mathfrak{p}_\mu}(\alpha) = 0$. (The nonzero coefficients of a root must all have the same sign.) Since $J^\mathfrak{p}_\mu = J^\mathfrak{q}_\mu \,\dot\cup\, (I_\mathfrak{q} \backslash (I_\mathfrak{p} \cup I^\mathfrak{q}_\mu))$, then:

(i) $Z_{J^\mathfrak{q}_\mu}(\alpha) = 0$. Thus, $\alpha \notin \Delta(\mathfrak{q}_{0,+})$, so $\alpha \in \Delta(\mathfrak{q}_+)$, i.e. $r = Z_{I_\mathfrak{q}}(\alpha) > 0$;

(ii) $Z_{I_\mathfrak{q} \backslash (I_\mathfrak{p} \cup I^\mathfrak{q}_\mu)}(\alpha) = 0$. Since $Z_{I_\mathfrak{p}}(\alpha) = 0$, then $Z_{I_\mathfrak{q} \backslash I^\mathfrak{q}_\mu}(\alpha) = 0$.

Therefore, $r = Z_{I^\mathfrak{q}_\mu}(\alpha) > 0$. By Theorem 3.2.6, $\alpha \in \Delta(\mathfrak{b}_r)$. Thus, $\mathfrak{a}_0 \subset \mathfrak{b}$. Hence, $\mathfrak{a} \subset \mathfrak{b}$.

Conversely, let $\alpha \in \Delta(\mathfrak{b})$. We show that $\alpha \in \Delta(\mathfrak{a})$. We have three cases:

(i) $Z_{I_\mathfrak{p}}(\alpha) < 0$: Then $\alpha \in \Delta(\mathfrak{p}_-) \subset \Delta(\mathfrak{a})$.

(ii) $Z_{I_\mathfrak{p}}(\alpha) = 0$: Assume $Z_{J_\mu^\mathfrak{p}}(\alpha) > 0$, so $Z_{J_\mu^\mathfrak{q}}(\alpha) \geq 0$ since $J_\mu^\mathfrak{q} \subset J_\mu^\mathfrak{p}$. Hence, $\alpha \in \Delta(\mathfrak{b}_{\geq 0})$, so by Theorem 3.2.6, $Z_{J_\mu^\mathfrak{q}}(\alpha) \leq 0$. Thus, $Z_{J_\mu^\mathfrak{q}}(\alpha) = 0$ and $Z_{J_\mu^\mathfrak{p} \backslash J_\mu^\mathfrak{q}}(\alpha) > 0$. Since $J_\mu^\mathfrak{p} \backslash J_\mu^\mathfrak{q} \subset I_\mathfrak{q} \backslash I_\mu^\mathfrak{q}$, then $Z_{I_\mathfrak{q} \backslash I_\mu^\mathfrak{q}}(\alpha) > 0$. But by Theorem 3.2.6, $Z_{I_\mathfrak{q} \backslash I_\mu^\mathfrak{q}}(\alpha) = 0$, a contradiction. Thus, $Z_{J_\mu^\mathfrak{p}}(\alpha) \leq 0$, so $\alpha \in \Delta(\mathfrak{a}_0)$.

(iii) $Z_{I_\mathfrak{p}}(\alpha) > 0$: Since $I_\mathfrak{p} \subset I_\mathfrak{q} \backslash I_\mu^\mathfrak{q}$, then $Z_{I_\mathfrak{q} \backslash I_\mu^\mathfrak{q}}(\alpha) > 0$ and $\alpha \in \Delta(\mathfrak{b}_+)$. But the latter implies $Z_{I_\mathfrak{q} \backslash I_\mu^\mathfrak{q}}(\alpha) = 0$ by Theorem 3.2.6, a contradiction.

Thus, $\mathfrak{b} = \mathfrak{p}_- \oplus \mathfrak{a}_0 = \mathfrak{a}$. ☐

Example 3.5.2. *Consider $w = (21)$ for A_6/P_2, $A_6/P_{1,2}$, $A_6/P_{1,2,4}$, and $A_6/P_{1,2,4,5}$. In each case, $w \in W_+^\mathfrak{p}(2)$, and the grading on $\mathfrak{a}(w)$ is given below. (Diagonal entries are suppressed.) In all four cases, we obtain the same (Z_2-graded) Lie algebra $\mathfrak{a}(w) \subset A_6 = \mathfrak{sl}_7(\mathbb{C})$.*

*	0					
0	*					
-1	-1	*				
-1	-1	0	*	0	0	
-1	-1	0	0	*	0	
-1	-1	0	0	0	*	
-1	-1	0	0	0	0	*

A_6/P_2

*	1					
-1	*					
-2	-1	*				
-2	-1	0	*	0	0	
-2	-1	0	0	*	0	
-2	-1	0	0	0	*	
-2	-1	0	0	0	0	*

$A_6/P_{1,2}$

*	1					
-1	*					
-2	-1	*				
-2	-1	0	*	1	1	
-3	-2	-1	-1	*	0	
-3	-2	-1	-1	0	*	
-3	-2	-1	-1	0	0	*

$A_6/P_{1,2,4}$

*	1					
-1	*					
-2	-1	*				
-2	-1	0	*	1	2	
-3	-2	-1	-1	*	1	
-4	-3	-2	-2	-1	*	
-4	-3	-2	-2	-1	0	*

$A_6/P_{1,2,4,5}$

3.6. **Prolongation height.** The *depth* of the Z-grading on $\mathfrak{a}$ is the maximal $\nu \geq 0$ such that $\mathfrak{a}_{-\nu} \neq 0$. The *height* of the Z-grading on $\mathfrak{a}$ is the maximal $\tilde{\nu} \geq 0$ such that $\mathfrak{a}_{\tilde{\nu}} \neq 0$.

Theorem 3.6.1. *Let $\mathfrak{g}$ be a complex simple Lie algebra, $\mathfrak{p} \subset \mathfrak{g}$ a parabolic subalgebra. If $(\mathfrak{g}, \mathfrak{p})$ is NPR$_+$, then the height of $\mathfrak{a} = \mathfrak{a}(w)$, where $w \in W_+^\mathfrak{p}(2)$, is always $\tilde{\nu} = 1$, except for*

Label	G/P	*Range*	$w \in W_+^\mathfrak{p}(2)$	J_w	I_w
NPR–A	$A_\ell/P_{1,2,s,t}$	$4 \leq s < t < \ell$	(21)	$\{3, \ell\}$	$\{1, s, t\}$
NPR–C	$C_\ell/P_{1,2,s}$	$4 \leq s < \ell$	(21)	$\{3\}$	$\{1, s\}$

In these two exceptional cases, $\tilde{\nu} = 2$ and

	$\dim(\mathfrak{a}_0)$	$\dim(\mathfrak{a}_1)$	$\dim(\mathfrak{a}_2)$
(A) :	$(s-3)(s-2) + (t-s)^2 + 2 + (\ell-t)(\ell-t+1)$	$1 + (t-s)(\ell-3-t+s)$	$(s-3)(\ell-t)$
(C) :	$2 + (s-3)(s-2) + (\ell-s)(2\ell-2s+1)$	$1 + 2(s-3)(\ell-s)$	$\binom{s-2}{2}$

Proof. Examine Tables 15 and 16. Except for NPR-A and NPR-C, $(\bar{\mathfrak{g}}, \bar{\mathfrak{p}})$ is always 1-graded, so $\mathfrak{a}_2 = 0$. For NPR-A and NPR-C, $(\bar{\mathfrak{g}}, \bar{\mathfrak{p}})$ is 2-graded, so $\mathfrak{a}_3 = 0$. The remaining data listed above is calculated using the Dynkin diagram recipes. ☐

If we remove the regularity assumption, higher prolongations can exist.

Example 3.6.2. *For $\ell \geq 5$, consider $A_\ell/B = A_\ell/P_{1,2,\ldots,\ell}$ and $w = (21)$. This yields*



, *homogeneity* $= 5 - \ell$.

Thus, $J_w = \emptyset$, $I_w = \{1, 4, 5, \ldots, \ell-1\}$, so $\bar{\mathfrak{g}}/\bar{\mathfrak{p}} \cong A_1/P_1 \times A_{\ell-4}/P_{1,\ldots,\ell-4}$. Note $\nu = \ell$, while $\tilde{\nu} = \ell-4$.

4. SUBMAXIMAL SYMMETRY DIMENSIONS

Define the *submaximal symmetry dimension* $\mathfrak{S}$ as the maximum $\dim(\mathfrak{inf}(\mathcal{G},\omega))$ among all **regular, normal** G/P geometries $(\mathcal{G} \xrightarrow{\pi} M, \omega)$ which are **not locally flat**. (Equivalently, $\mathfrak{S}$ maximizes $\dim(\mathcal{S})$, where $\mathcal{S}$ is the symmetry algebra of an underlying structure which is not locally flat.)

4.1. Filtration and embedding of the symmetry algebra.

Let us review some known results.

Čap [8], *Čap–Neusser* [9]: Let $(\mathcal{G} \xrightarrow{\pi} M, \omega)$ be a regular, normal G/P geometry. Let $u \in \mathcal{G}$, and $x = \pi(u)$. The map $\xi \mapsto \omega(\xi(u))$ defines a linear injection $\widetilde{\mathcal{S}} := \mathfrak{inf}(\mathcal{G},\omega) \hookrightarrow \mathfrak{g}$.[8] Letting $\mathfrak{f}(u) \subset \mathfrak{g}$ denote its image and κ the curvature function, the bracket on $\mathfrak{inf}(\mathcal{G},\omega)$ is mapped to the operation

$$[X,Y]_{\mathfrak{f}(u)} := [X,Y] - \kappa_u(X,Y)$$

on $\mathfrak{f}(u)$. The filtration on $\mathfrak{g}$ restricts to $\mathfrak{f}(u)$, and by regularity, $\mathfrak{f}(u)$ is a filtered Lie algebra (generally not a subalgebra of $\mathfrak{g}$). The associated-graded

$$\mathfrak{s}(u) := \mathrm{gr}(\mathfrak{f}(u)) = \bigoplus_i \mathfrak{f}^i(u)/\mathfrak{f}^{i+1}(u) \subset \mathfrak{g}$$

is a graded subalgebra. (Note that $\mathrm{gr}(\mathfrak{g}) \cong \mathfrak{g}$.) Moreover, $\mathfrak{s}_0(u) \subseteq \mathfrak{ann}(\kappa_H(u)) \subseteq \mathfrak{g}_0$ [9, Proof of (3) in Corollary 5]. Though stated for bracket-generating distributions, their argument works in general: Given $\xi \in \mathfrak{inf}(\mathcal{G},\omega)$, if $Y = \omega(\xi(u)) \in \mathfrak{p}$, then the curve $\varphi_t(u) = u \cdot \exp(tY)$ has vertical tangent vector $\xi(u) = \zeta_Y(u) = \frac{d}{dt}\big|_{t=0} \varphi_t(u)$. By P-equivariancy of κ_H,

$$0 = (\xi \cdot \kappa_H)(u) = \frac{d}{dt}\bigg|_{t=0} \kappa_H(\varphi_t(u)) = \frac{d}{dt}\bigg|_{t=0} \exp(-tY) \cdot (\kappa_H(u)) = -Y \cdot (\kappa_H(u)).$$

Complete reducibility of $H^2_+(\mathfrak{g}_-,\mathfrak{g})$ implies that this action depends only on $Y \bmod \mathfrak{p}_+ \in \mathfrak{p}/\mathfrak{p}_+ \cong \mathfrak{g}_0$. If $\kappa_H(u) \neq 0$, Kostant's theorem (see Recipe 5) yields a bound on $\dim(\mathfrak{s}_0(u))$.

Remark 4.1.1. For non-flat geometries, Čap & Neusser bound $\dim(\mathfrak{inf}(\mathcal{G},\omega))$ via the maximum dimension of any proper graded subalgebra $\mathfrak{b} \subset \mathfrak{g}$ with $\dim(\mathfrak{b}_0)$ no larger than the Kostant bound. However, this strategy lacks finer information about the annihilating subalgebra, i.e. they use (2.13), while we will additionally use (2.12). Thus, in general, their upper bounds will not be sharp.

The filtration of $\mathfrak{g}$ induces a P-invariant filtration $\{T^i\mathcal{G}\}_{i=-\nu}^{\nu}$ of $T\mathcal{G}$, and a filtration of $\widetilde{\mathcal{S}}$:

$$\widetilde{\mathcal{S}}(x)^i = \{\xi \in \widetilde{\mathcal{S}} \mid \xi(u) \in T^i_u\mathcal{G},\ \forall u \in \pi^{-1}(x)\}, \qquad -\nu \leq i \leq \nu. \tag{4.1}$$

This projects to a filtration $\{\mathcal{S}(x)^i\}_{i=-\nu}^{\nu}$ of the symmetry algebra $\mathcal{S}$ of the underlying structure on M. We have $\omega_u(\widetilde{\mathcal{S}}(x)^i) = \mathfrak{f}(u)^i$, and $\mathfrak{s}(u) = \mathrm{gr}(\mathfrak{f}(u)) \cong \mathrm{gr}(\mathcal{S}(x)) =: \mathfrak{s}(x)$ are canonically isomorphic. For bracket-generating distributions (not necessarily underlying parabolic geometries), a filtration $\{\mathcal{S}(x)^i\}$ of $\mathcal{S}$ was constructed in [29]. We summarize its definition and properties below.

Kruglikov [29]: Let $\{D^i\}_{i=-\nu}^{-1}$ be a filtration of TM, with $D := D^{-1}$ bracket-generating and suppose each D^i has constant rank.[9] Given $x \in M$, $\mathfrak{g}_i(x) := D^i_x/D^{i+1}_x$ for $i < 0$. The Tanaka prolongation $\mathfrak{g}(x) = \mathrm{pr}(\mathfrak{g}_-(x))$ has $\mathfrak{g}_i(x) \hookrightarrow \mathrm{Hom}(\otimes^{i+1}\mathfrak{g}_{-1}(x), \mathfrak{g}_{-1}(x))$ for $i \geq 0$, cf. Remark 2.1.2.

Letting $\mathrm{ev}_x : \mathcal{S} \to T_xM$ be the evaluation map, define $\mathcal{S}(x)^i := \mathrm{ev}_x^{-1}(D^i)$ for $i < 0$, and $\mathcal{S}(x)^0 := \ker(\mathrm{ev}_x)$. Inductively, for $i \geq 0$, given $\mathbf{X} \in \mathcal{S}(x)^i$, there is a map $\Psi^{i+1}_{\mathbf{X}} : \otimes^{i+1} D_x \to T_xM$ given by

$$\Psi^{i+1}_{\mathbf{X}}(Y_1,\dots,Y_{i+1}) = \big[\big[\dots\big[[\mathbf{X},\mathbf{Y}_1],\mathbf{Y}_2\big],\dots\big],\mathbf{Y}_{i+1}\big](x),$$

where $\mathbf{Y}_j \in \Gamma(D)$ and $\mathbf{Y}_j(x) = Y_j$. Moreover, $\mathrm{im}(\Psi^{i+1}_{\mathbf{X}}) \subseteq D_x$. Define $\mathcal{S}(x)^{i+1} := \{\mathbf{X} \in \mathcal{S}(x)^i \mid \Psi^{i+1}_{\mathbf{X}} = 0\}$. Then $\mathcal{S} = \mathcal{S}(x)$ is a filtered Lie algebra, and $\mathfrak{s}_i(x) \hookrightarrow \mathfrak{g}_i(x)$ via $\mathbf{X} \bmod \mathcal{S}(x)^{i+1} \mapsto \Psi^{i+1}_{\mathbf{X}}$.

---

[8] They state this for the global automorphism algebra $\mathfrak{aut}(\mathcal{G},\omega)$, but it is established for $\mathfrak{inf}(\mathcal{G},\omega)$ in their proof.

[9] These conditions hold for underlying structures of parabolic geometries.

There is a filtration[10] on $C^\infty(M)$ by ideals μ_x^i. Let $\mu_x^1 = \{f \in C^\infty(M) \mid f(x) = 0\}$ and for $i \geq 0$,

$$\mu_x^{i+1} = \{f \in C^\infty(M) \mid (\mathbf{Y}_1 \cdots \mathbf{Y}_t \cdot f)(x) = 0,\ \forall \mathbf{Y}_j \in \Gamma(D),\ 0 \leq t \leq i\}.$$

For any $\mathbf{Y} \in \Gamma(D)$, we have $\mathbf{Y} \cdot \mu_x^{i+1} \subset \mu_x^i$. Let $\{\mathbf{Z}_{jk}\}_{j=1,\ldots,\nu}$ be a local framing near x with $\mathbf{Z}_{jk} \in \Gamma(D^{-j}) \backslash \Gamma(D^{-j-1})$. Let $\mathbf{X} \in \mathcal{S}$. Then

$$i \geq 0 : \qquad \mathbf{X} = \sum_{j,k} f_{jk} \mathbf{Z}_{jk} \in \mathcal{S}(x)^i \quad \text{iff} \quad f_{jk} \in \mu_x^{i+j}, \quad \forall j, k. \tag{4.2}$$

The "only if" direction is proved in [29, Lemma 1], while an easy induction establishes the converse.

Example 4.1.2 (G_2/P_1)**.** *Consider D spanned by ∂_q and $\tilde{\partial}_x := \partial_x + p\partial_y + q\partial_p + (p^3 + q^2)\partial_z$. Then*

$$\mathcal{S}: \qquad \mathbf{X}_1 = \partial_x, \quad \mathbf{X}_2 = \partial_y, \quad \mathbf{X}_3 = \partial_z, \quad \mathbf{X}_4 = x\partial_x - y\partial_y - 2p\partial_p - 3q\partial_q - 5z\partial_z.$$

At $\mathbf{u}_0 = (x_0, y_0, p_0, q_0, z_0)$, the dimensions of $\mathfrak{s}_i = \mathfrak{s}_i(\mathbf{u}_0)$ are

	$\mathfrak{s}_{-3}$	$\mathfrak{s}_{-2}$	$\mathfrak{s}_{-1}$	$\mathfrak{s}_0$
$q_0 \neq 0$ *or* $p_0 \neq 0$	2	1	1	0
$q_0 = p_0 = 0$	2	0	1	1

When $p_0 = q_0 = 0$, $\mathcal{S}(\mathbf{u}_0)^0$ is spanned by $\mathbf{T} = \mathbf{X}_4 - x_0\mathbf{X}_1 + y_0\mathbf{X}_2 + 5z_0\mathbf{X}_3$. Writing $\mathbf{T} = \sum_{j,k} f_{jk}\mathbf{Z}_{jk}$ in the framing $(\mathbf{Z}_{11}, \mathbf{Z}_{12}, \mathbf{Z}_{21}, \mathbf{Z}_{31}, \mathbf{Z}_{32}) = (\partial_q, \tilde{\partial}_x, \partial_p + 2q\partial_z, \partial_y, \partial_z)$ shows $f_{jk} \in \mu_x^j$, so $\mathbf{T} \notin \mathcal{S}(\mathbf{u}_0)^1$. Note $[\mathbf{T}, \partial_q] = 3\partial_q$, which at $\mathbf{u}_0$ is not contained in $\mathfrak{s}_{-1}$. Thus, $[\mathfrak{s}_0, \mathfrak{g}_{-1}] \not\subset \mathfrak{s}_{-1}$ when $p_0 = q_0 = 0$.

Remark 4.1.3. The above results in Kruglikov's approach still hold if D admits additional structure. In the parabolic case, Kruglikov's filtration $\{\mathcal{S}(x)^i\}$ agrees with that of Čap–Neusser, and (4.2) gives a means to compute it explicitly. Henceforth, we will mainly rely on the Čap–Neusser presentation.

4.2. **A universal upper bound.**

Definition 4.2.1. *We say that $x \in M$ is a* regular point *if there exists a neighbourhood $N_x \subset M$ of x such that for $-\nu \leq j \leq \nu$, $\dim(\mathfrak{s}_j(y))$ is a constant function of $y \in N_x$. Otherwise, x is* irregular.

We have a tower of bundles $\mathcal{G} =: \mathcal{G}_\nu \to \ldots \to \mathcal{G}_0 \to M$, with $\mathcal{G}_i = \mathcal{G}/P_+^{i+1} = \mathcal{G}/\exp(\mathfrak{g}^{i+1})$, projections $p_i : \mathcal{G} \to \mathcal{G}_i$ and $\pi_i : \mathcal{G}_i \to M$, where $i = -1, \ldots, \nu$. (If $i = -1$, let $p_{-1} = \pi$, i.e. the projection to M.) Given any $\xi \in \mathfrak{X}(\mathcal{G})^P$, let $\xi^{(i)} = (p_i)_*\xi \in \mathfrak{X}(\mathcal{G}_i)$. By P-invariancy, $\widetilde{\mathcal{S}}$ descends to $\mathcal{S}^{(i)} \subset \mathfrak{X}(\mathcal{G}_i)^{P/P_+^{i+1}}$. Given $x \in M$, the filtration $\widetilde{\mathcal{S}}(x)$ projects to a filtration $\mathcal{S}^{(i)}(x)$, and from (4.1), $\mathcal{S}^{(i)}(x)^{i+1} = \{\xi^{(i)} \in \mathcal{S}^{(i)} \mid \xi^{(i)}(u_i) = 0,\ \forall u_i \in \pi_i^{-1}(x)\}$. Thus, for any $u_i \in \pi_i^{-1}(x)$, $\mathcal{S}^{(i)}|_{u_i} \subset T_{u_i}\mathcal{G}_i$ has dimension $\dim \mathcal{S} - \sum_{j=i+1}^{\nu} \dim \mathfrak{s}_j(x)$. If x is a regular point, then $\mathcal{S}^{(i)}$ yields a constant rank distribution on a neighbourhood of $\pi_i^{-1}(x) \subset \mathcal{G}_i$. By the Frobenius theorem, there exist local rectifying coordinates and a local foliation by maximal integral submanifolds of $\mathcal{S}^{(i)}$. We use this to prove the following "filtered" generalization of a result of Morimoto [32, Proposition 4.1].

Proposition 4.2.2. *Let $x \in M$ be a regular point. Then for any i, $[\mathfrak{s}_{i+1}(u), \mathfrak{g}_{-1}] \subset \mathfrak{s}_i(u)$. In particular, for $i > 0$, the i-th graded component of $\mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{s}_0(u))$ contains $\mathfrak{s}_i(u)$.*

If $\mathcal{S}$ is *transitive at* x, i.e. $\mathrm{ev}_x(\mathcal{S}) = T_xM$, then $[\mathfrak{s}_{i+1}(x), \mathfrak{g}_{-1}(x)] = [\mathfrak{s}_{i+1}(x), \mathfrak{s}_{-1}(x)] \subset \mathfrak{s}_i(x)$ immediately follows since $\mathfrak{s}(x)$ is a graded Lie algebra. Equivalently, $[\mathfrak{s}_{i+1}(u), \mathfrak{g}_{-1}] \subset \mathfrak{s}_i(u)$.

Example 4.2.3. *In Example 4.1.2, the regular points are precisely those with $p_0 \neq 0$ or $q_0 \neq 0$. At all irregular points, the first claim of Proposition 4.2.2 fails.*

---

[10] In the complex setting, we replace $C^\infty(M)$ by the algebra of germs of holomorphic functions about the point x.

Proof of Proposition 4.2.2. Suppose $i \geq -1$. For any $\xi \in \widetilde{\mathcal{S}} = \mathfrak{inf}(\mathcal{G},\omega)$ and $\eta \in \mathfrak{X}(\mathcal{G})$, we have $0 = (\mathcal{L}_\xi \omega)(\eta) = \xi \cdot \omega(\eta) - \omega([\xi,\eta])$. Let $\xi \in \widetilde{\mathcal{S}}(x)^{i+1}$, and $\eta \in \Gamma(T^{-1}\mathcal{G})^P$. Let $u \in \pi^{-1}(x)$, so $X = \omega(\xi(u)) \in \mathfrak{f}(u)^{i+1} \subset \mathfrak{g}^{i+1}$, and $Y = \omega(\eta(u)) \in \mathfrak{g}^{-1}$. Since $i \geq -1$, then $X \in \mathfrak{p}$, so $X = \omega(\zeta_X(u))$, where ζ_X is a fundamental vertical vector field. By P-equivariancy of $\omega(\eta)$,

$$\omega([\xi,\eta](u)) = (\xi \cdot \omega(\eta))(u) = (\zeta_X \cdot \omega(\eta))(u) = \left.\frac{d}{dt}\right|_{t=0} \omega(\eta)(u \cdot \exp(tX)) = -[X,Y].$$

In particular, since $[X,Y] \in \mathfrak{g}^i$, then $[\xi,\eta](u) \in T^i_u\mathcal{G}$.

Let $u_i = p_i(u) \in \mathcal{G}_i$. Since $\xi \in \widetilde{\mathcal{S}}(x)^{i+1}$, then $\xi^{(i)}(u_i) = 0$. Since x is a regular point, there exist local (functionally independent) functions $F_1, \dots, F_{t_i}$ on $\mathcal{G}_i$ (smooth in the real setting or holomorphic in the complex setting), where $t_i = \dim \mathcal{G}_i - \sum_{-\nu}^{i} \dim \mathfrak{s}_j(x)$, whose joint level sets define the local foliation by maximal integral submanifolds of $\mathcal{S}^{(i)}$. In particular, for our $\xi \in \widetilde{\mathcal{S}}(x)^{i+1}$, we have $\xi^{(i)} \cdot F_j = 0$ for $j = 1, \dots, t_i$, and since $\xi^{(i)}(u_i) = 0$, we have $([\xi^{(i)}, \eta^{(i)}] \cdot F_j)(u_i) = 0$. Thus, $[\xi^{(i)}, \eta^{(i)}](u_i) \in \mathcal{S}^{(i)}|_{u_i} = (p_i)_*(\widetilde{\mathcal{S}}|_u)$ is tangent to the foliation. Since any element of $\widetilde{\mathcal{S}}$ is uniquely determined by its value at u, then $[\xi^{(i)}, \eta^{(i)}](u_i) = \xi'^{(i)}(u_i)$ for some $\xi' \in \widetilde{\mathcal{S}}$. Equivalently,

$$[\xi,\eta](u) = \xi'(u) + \chi(u) \in T^i_u\mathcal{G}, \qquad \xi'(u) \in \widetilde{\mathcal{S}}|_u, \quad \chi(u) \in T^{i+1}_u\mathcal{G}.$$

Hence, $\omega([\xi,\eta](u)) \in \mathfrak{f}(u)^i + \mathfrak{g}^{i+1}$. Thus, $[\mathfrak{f}(u)^{i+1}, \mathfrak{g}^{-1}] \subset \mathfrak{f}(u)^i + \mathfrak{g}^{i+1}$, so $[\mathfrak{s}_{i+1}(u), \mathfrak{g}_{-1}] \subset \mathfrak{s}_i(u)$.

Now suppose $i \leq -2$. Since x is regular, then $D^j_{\mathcal{S}}|_y := D^j_y \cap \mathcal{S}|_y$, $j \leq -1$, define constant rank distributions near x. Let $\mathbf{X} = \sum_i f_i \mathbf{X}_i \in \Gamma(D^{i+1}_{\mathcal{S}})$ where $\{\mathbf{X}_i\}$ is basis of $\mathcal{S}$. Given $\mathbf{Y} \in \Gamma(D)$,

$$[\mathbf{X},\mathbf{Y}](x) = \sum_i f_i(x)[\mathbf{X}_i,\mathbf{Y}](x) - (\mathbf{Y}\cdot f_i)(x)\mathbf{X}_i(x) \in (D_x + \mathcal{S}|_x) \cap D^i_x.$$

Quotienting by $D^{i+1}_x \supset D_x$ implies that with respect to the Levi bracket, $[\mathfrak{s}_{i+1}(x), \mathfrak{g}_{-1}(x)] \subset \mathfrak{s}_i(x)$, so $[\mathfrak{s}_{i+1}(u), \mathfrak{g}_{-1}] \subset \mathfrak{s}_i(u)$. The second claim immediately follows from the first and Definition 2.3.3. $\square$

Lemma 4.2.4. *The set of regular points is open and dense in M.*

Proof. For $-\nu \leq i \leq \nu$, define $q_i^\pm : M \to \mathbb{Z}$ by $q_i^-(x) = \sum_{j=-\nu}^{i} \dim \mathfrak{s}_j(x)$ and $q_i^+(x) = \dim \mathcal{S} - q_i^-(x) = \dim \mathcal{S}(x)^{i+1} = \sum_{j=i+1}^{\nu} \dim \mathfrak{s}_j(x)$. Then q_i^- is lower semi-continuous because linear independence of $\xi_1^{(i)}, \dots, \xi_s^{(i)} \in \mathcal{S}^{(i)}$ at $u_i \in \mathcal{G}_i$ persists near u_i. Hence, q_i^+ is upper semi-continuous. Given $U \subset M$, define $m_i(U) = \{x \in U \mid q_i^+(x) \leq q_i^+(y), \forall y \in U\}$. Upper semi-continuity implies that if $\emptyset \neq U \subset M$ is open, then $\emptyset \neq m_i(U) \subset M$ is open.

Denote $M_0 := M$, and define $N_{j+1} = m_{-\nu} \circ \cdots \circ m_\nu(M_j)$ and $M_{j+1} = M \backslash (\bar{N}_1 \cup \cdots \cup \bar{N}_j)$ for $j \geq 1$. Each $M_j \subset M$ is open, so $N_{j+1} \subset M$ is open; when the former is non-empty, so is the latter. Each $\operatorname{im}(q_i^+) \subset [0, \dim \mathcal{S}] \cap \mathbb{Z}$ is finite, so there exists a minimal $k \geq 0$ with $M_{k+1} = M$. The open set $N = N_1 \cup \cdots \cup N_k$ is the set of all regular points, which is dense in M since $\bar{N} = M$. $\square$

From Section 3, $\mathfrak{a}^\phi := \operatorname{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{ann}(\phi))$ and $\mathfrak{U} := \max\left\{\dim(\mathfrak{a}^\phi) \mid 0 \neq \phi \in H^2_+(\mathfrak{g}_-, \mathfrak{g})\right\} < \dim(\mathfrak{g})$.

Theorem 4.2.5. *Let G be a real or complex semisimple Lie group and $P \subset G$ a parabolic subgroup. Let $(\mathcal{G} \xrightarrow{\pi} M, \omega)$ be a regular, normal G/P geometry, $x \in M$ a regular point, and $u \in \pi^{-1}(x)$. Then $\mathfrak{s}(u) \subseteq \mathfrak{a}^{\kappa_H(u)}$ is a graded subalgebra, and* $\dim(\mathfrak{inf}(\mathcal{G},\omega)) \leq \dim(\mathfrak{a}^{\kappa_H(u)})$. *Thus,* $\mathfrak{S} \leq \mathfrak{U} < \dim(\mathfrak{g})$.

Proof. Using Proposition 4.2.2 and $\mathfrak{s}_0(u) \subseteq \mathfrak{ann}(\kappa_H(u))$, we have $\mathfrak{s}(u) \subseteq \mathfrak{g}_- \oplus \mathfrak{s}_{\geq 0}(u) \subseteq \operatorname{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{s}_0(u)) \subseteq \mathfrak{a}^{\kappa_H(u)}$. Thus, $\dim(\mathfrak{inf}(\mathcal{G},\omega)) = \dim(\mathfrak{s}(u)) \leq \dim(\mathfrak{a}^{\kappa_H(u)})$.

If the geometry is not locally flat, let $N = \{x \in M \mid \kappa_H(u) \neq 0, \forall u \in \pi^{-1}(x)\}$. Then N is non-empty and open, so by Lemma 4.2.4, N contains a (non-flat) regular point x. Hence, $\dim(\mathfrak{inf}(\mathcal{G},\omega)) \leq \dim(\mathfrak{a}^{\kappa_H(u)}) \leq \mathfrak{U}$ for any $u \in \pi^{-1}(x)$. Thus, $\mathfrak{S} \leq \mathfrak{U} < \dim(\mathfrak{g})$. $\square$

Remark 4.2.6. Only $[\mathfrak{s}_{i+1}(u), \mathfrak{g}_{-1}] \subseteq \mathfrak{s}_i(u)$ for $i \geq 0$ was essential for proving Theorem 4.2.5. This relied only on local constancy of $\dim \mathfrak{s}_j(y)$ for $j > 0$, so our "regular point" definition could be weakened. The property $[\mathfrak{s}_0(u), \mathfrak{g}_{-1}] \subseteq \mathfrak{s}_{-1}(u)$ will be used in proving Theorems 5.1.3 and 5.5.2.

Remark 4.2.7. Over $\mathbb{R}$, $\mathfrak{S} \leq \mathfrak{U} \leq \mathfrak{U}_{\mathbb{C}}$, where $\mathfrak{U}_{\mathbb{C}}$ is the $\mathfrak{U}$ for the complexification $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{p}_{\mathbb{C}})$ of $(\mathfrak{g}, \mathfrak{p})$.

Corollary 4.2.8. *Let G be a real or complex semisimple Lie group and $P \subset G$ a parabolic subgroup. For regular, normal G/P geometries, suppose that $\mathfrak{S} = \mathfrak{U}$. Then any submaximally symmetric model $(\mathcal{G} \stackrel{\pi}{\to} M, \omega)$ is locally homogeneous near a non-flat regular point $x \in M$.*

Proof. Since $\mathfrak{S} = \mathfrak{U}$, any submaximally symmetric model has $\mathfrak{U} = \dim(\mathfrak{inf}(\mathcal{G}, \omega)) = \dim(\mathfrak{s}(u)) \leq \dim(\mathfrak{a}^{\kappa_H(u)}) \leq \mathfrak{U}$ for any $u \in \pi^{-1}(x)$, by Theorem 4.2.5(a). Hence, $\mathfrak{s}(u) = \mathfrak{a}^{\kappa_H(u)} \supset \mathfrak{g}_-$, so $\mathfrak{s}(x) = \mathrm{gr}(\mathcal{S}(x)) \supset \mathfrak{g}_-$, i.e. $\mathcal{S}$ is transitive at x, so the result follows by Lie's third theorem. $\square$

Remark 4.2.9. By P-equivariancy of κ_H and complete reducibility of $H^2_+(\mathfrak{g}_-, \mathfrak{g})$, fibres of $\mathcal{G} \to M$ are mapped by κ_H into G_0-orbits in $H^2_+(\mathfrak{g}_-, \mathfrak{g})$. Hence, we can further constrain κ_H in two basic ways, and since regular points are dense (Lemma 4.2.4), we get analogues of Theorem 4.2.5:

(i) Let $0 \neq \mathbb{V}_i \subset H^2_+(\mathfrak{g}_-, \mathfrak{g})$ be a G_0-submodule, and define $\mathfrak{S}_i$, $\mathfrak{U}_i$ analogous to $\mathfrak{S}$, $\mathfrak{U}$, e.g. $\mathfrak{S}_i$ requires $\mathrm{im}(\kappa_H) \subset \mathbb{V}_i$. Then $\mathfrak{S}_i \leq \mathfrak{U}_i$. In particular, if $\mathfrak{g}$ is complex semisimple, we have $\mathfrak{S}_\mu \leq \mathfrak{U}_\mu$. From Section 3, $\mathfrak{U}_\mu = \dim(\mathfrak{a}(\mu))$ is easily computable.

(ii) Let $0 \neq \mathcal{O} \subset H^2_+(\mathfrak{g}_-, \mathfrak{g})$ be G_0-invariant. Analogously define $\mathfrak{U}_{\mathcal{O}}$ and $\mathfrak{S}_{\mathcal{O}}$. Then $\mathfrak{S}_{\mathcal{O}} \leq \mathfrak{U}_{\mathcal{O}}$.

If either $\mathfrak{S}_i = \mathfrak{U}_i$ or $\mathfrak{S}_{\mathcal{O}} = \mathfrak{U}_{\mathcal{O}}$, then as in the proof of Corollary 4.2.8, any model realizing the equality must be locally homogeneous near a non-flat regular point.

We will use (ii) in Section 5.1.2 to study the Petrov types in 4-dimensional Lorentzian conformal geometry, and in Section 5.5.1 for $(2, 3, 5)$-distributions with κ_H having constant root type.

4.3. **Establishing submaximal symmetry dimensions.** In this section, we show that in the *complex* (or *split-real*) case, the upper bound $\mathfrak{S}_\mu \leq \mathfrak{U}_\mu = \dim(\mathfrak{a}(\mu))$ (for regular, normal geometries) obtained from Theorem 4.2.5 and Remark 4.2.9 is almost always sharp. Let us outline the strategy: Kostant's theorem (Recipe 4) gives an explicit description of the lowest $\mathfrak{g}_0$-weight vector $\phi_0 \in \mathbb{V}_\mu$. Using the $\mathfrak{g}_0$-module isomorphism $H^2(\mathfrak{g}_-, \mathfrak{g}) \cong \ker(\Box) \subset \bigwedge^2 \mathfrak{g}_-^* \otimes \mathfrak{g}$, we have ϕ_0 naturally realized in $\bigwedge^2 \mathfrak{g}_-^* \otimes \mathfrak{g}$. When $\mathrm{im}(\phi_0) \subset \mathfrak{a} := \mathfrak{a}^{\phi_0}$, we can regard $\phi_0 \in \bigwedge^2 \mathfrak{g}_-^* \otimes \mathfrak{a} \hookrightarrow \bigwedge^2 \mathfrak{a}^* \otimes \mathfrak{a}$. Now, we *deform the Lie bracket on $\mathfrak{a}$ by ϕ_0*. Namely, define $\mathfrak{f}$ to be the vector space $\mathfrak{a}$ with the deformed bracket

$$[\cdot, \cdot]_{\mathfrak{f}} := [\cdot, \cdot] - \phi_0(\cdot, \cdot), \tag{4.3}$$

and $[\cdot, \cdot]_{\mathfrak{f}}$ is indeed a Lie bracket. Since ϕ_0 vanishes on $\mathfrak{p} = \mathfrak{g}_{\geq 0}$, then $\mathfrak{k} := \mathfrak{a}_{\geq 0}$ is a subalgebra of both $\mathfrak{a}$ and $\mathfrak{f}$. We construct a *local* homogeneous space $M = F/K$ (i.e. F may only be a local Lie group) and P-bundle $\mathcal{G} = F \times_K P$ with an F-invariant normal Cartan connection ω whose curvature is determined by ϕ_0. Thus, $\mathfrak{S}_\mu \geq \dim(\mathfrak{inf}(\mathcal{G}, \omega)) \geq \dim(\mathfrak{f}) = \dim(\mathfrak{a})$. (In fact, this construction does not depend on regularity.)

Lemma 4.3.1. *Let λ be the highest weight of a simple ideal in $\mathfrak{g}$ and $w \in W^{\mathfrak{p}}(2)$. Suppose that $w(-\lambda) \in \Delta^-$. Let ϕ_0 be the lowest weight vector of $\mathbb{V}_{-w\cdot\lambda}$ and $\mathfrak{a} := \mathfrak{a}^{\phi_0}$. Define $\mathfrak{f}$ to be the vector space $\mathfrak{a}$ with deformed Lie bracket* (4.3). *Then $\mathfrak{f}$ is a Lie algebra.*

Proof. We have $w(-\lambda) \in \Delta^- = \Delta(\mathfrak{g}_-) \cup \Delta^-(\mathfrak{g}_0)$. All root vectors from $\Delta^-(\mathfrak{g}_0)$ annihilate the lowest $\mathfrak{g}_0$-weight vector ϕ_0, hence $e_{w(-\lambda)} \in \mathfrak{g}_- \oplus \mathfrak{ann}(\phi_0) \subset \mathfrak{a}$. Thus, $\phi_0 = e_{\alpha_j} \wedge e_{\sigma_j(\alpha_k)} \otimes e_{w(-\lambda)}$ (see (2.7)) has $\mathrm{im}(\phi_0) \subset \mathfrak{a}$, so we can regard $\phi_0 \in \bigwedge^2 \mathfrak{a}^* \otimes \mathfrak{a}$. Note $e_{\alpha_j}, e_{\sigma_j(\alpha_k)} \in \mathfrak{g}_+ \cong \mathfrak{g}_-^*$ vanish on $\mathfrak{a}_{\geq 0}$.

By Theorem 3.5.1, the Lie algebra structure of $\mathfrak{a}$ is the same if we take $\mathfrak{p} = \mathfrak{p}_j$ if $c_{kj} < 0$ or $\mathfrak{p} = \mathfrak{p}_{j,k}$ if $c_{kj} = 0$. Assuming this, then by Proposition 3.3.6, $(\mathfrak{g}, \mathfrak{p}, -w \cdot \lambda)$ is PR, i.e. $\mathfrak{a} = \mathfrak{g}_- \oplus \mathfrak{a}_0$.

It suffices to verify the Jacobi identity $\mathrm{Jac}_{\mathfrak{f}}(x,y,z)=0$ for $[\cdot,\cdot]_{\mathfrak{f}}$. For $x,y,z\in\mathfrak{f}$,

$$\begin{aligned}[][x,y]_{\mathfrak{f}},z]_{\mathfrak{f}} &= [[x,y]-\phi_0(x,y),z]_{\mathfrak{f}}\\ &= [[x,y],z]-\phi_0([x,y],z)-[\phi_0(x,y),z]+\phi_0(\phi_0(x,y),z).\end{aligned}$$

Since ϕ_0 vanishes on $\mathfrak{a}_0$, clearly $\mathrm{Jac}_{\mathfrak{f}}(x,y,z)=0$ when at least two of $x,y,z\in\mathfrak{a}_0$. Suppose $x,y\in\mathfrak{g}_-$ and $z\in\mathfrak{a}_0$. Since $\mathrm{Jac}_{\mathfrak{a}}(x,y,z)=0$ and $z\in\mathfrak{a}_0=\mathfrak{ann}(\phi_0)$, then

$$\mathrm{Jac}_{\mathfrak{f}}(x,y,z)=[z,\phi_0(x,y)]-\phi_0([z,x],y)-\phi_0([y,z],x)=(z\cdot\phi_0)(x,y)=0.$$

Finally, suppose $x,y,z\in\mathfrak{g}_-$. Since $\phi_0\in\ker(\Box)$, it is in particular ∂-closed. This means

$$\begin{aligned}0=(\partial\phi_0)(x,y,z)&=[x,\phi_0(y,z)]-[y,\phi_0(x,z)]+[z,\phi_0(x,y)]\\ &\quad-\phi_0([x,y],z)+\phi_0([x,z],y)-\phi_0([y,z],x),\qquad \forall x,y,z\in\mathfrak{g}_-.\end{aligned}\tag{4.4}$$

This equation uses the bracket on $\mathfrak{g}$, but restricts to $\mathfrak{a}$ since ϕ_0 has image in $\mathfrak{a}$, and the bracket on $\mathfrak{a}$ is inherited from that of $\mathfrak{g}$ by Lemma 2.1.5. Using (4.4) and $\mathrm{Jac}_{\mathfrak{a}}(x,y,z)=0$, we obtain

$$\mathrm{Jac}_{\mathfrak{f}}(x,y,z)=\phi_0(\phi_0(x,y),z)+\phi_0(\phi_0(y,z),x)+\phi_0(\phi_0(z,x),y).$$

Since $\mathrm{im}(\phi_0)$ is spanned by $e_{w(-\lambda)}$, it suffices to show $w(-\lambda)\notin\{-\alpha_j,-\sigma_j(\alpha_k)\}$. If $w(-\lambda)=-\alpha_j$, then $\lambda=-\sigma_k(\alpha_j)\in\Delta^-$. Otherwise, if $w(-\lambda)=-\sigma_j(\alpha_k)$, then $\lambda=-\alpha_k\in\Delta^-$. But these contradict $\lambda\in\Delta^+$. Thus, $[\cdot,\cdot]_{\mathfrak{f}}$ is a Lie bracket on $\mathfrak{f}$. ☐

Let us determine all exceptions:

Lemma 4.3.2. *Let $\mathfrak{g}$ be complex simple. Let $w\in W^{\mathfrak{p}}$ satisfy $w(-\lambda_{\mathfrak{g}})\in\Delta^+$.*
(a) If $|w|=2$, then $\mathfrak{g}/\mathfrak{p}$ is one of: $A_2/\mathfrak{p}_1$, $A_2/\mathfrak{p}_{1,2}$, $B_2/\mathfrak{p}_1$, $B_2/\mathfrak{p}_{1,2}$.
(b) If $|w|=1$, then $\mathfrak{g}/\mathfrak{p}\cong A_1/\mathfrak{p}_1$.
For $B_2/\mathfrak{p}_{1,2}$, $w(-\lambda_{\mathfrak{g}})\in\Delta^+$ if $w=(12)$, but not for $w=(21)$.

Proof. Part (b) is obvious, so we prove (a). Since $W^{\mathfrak{p}}(2)\neq\emptyset$, then $\mathrm{rank}(\mathfrak{g})\geq 2$. Write $w=(jk)$. If $w(-\lambda_{\mathfrak{g}})\in\Delta^+$, then $w(-\lambda_{\mathfrak{g}})\in w\Delta^-\cap\Delta^+=\Phi_w=\{\alpha_j,\sigma_j(\alpha_k)\}$. There are two cases:

(i) $w(-\lambda_{\mathfrak{g}})=\alpha_j$: Then $\lambda_{\mathfrak{g}}=\sigma_k(\alpha_j)=\alpha_j-c_{jk}\alpha_k$, which is the highest root, so $\mathrm{rank}(\mathfrak{g})=2$.
(ii) $w(-\lambda_{\mathfrak{g}})=\sigma_j(\alpha_k)$: Then $\lambda_{\mathfrak{g}}=\alpha_k$. Hence, $\mathrm{rank}(\mathfrak{g})=1$, a contradiction.

For B_2/P_2, $\exists! w=(21)\in W^{\mathfrak{p}}(2)$, so $w(-\lambda_{\mathfrak{g}})=w(-2\lambda_1)=2\lambda_1-2\lambda_2=-\alpha_2\notin\Delta^+$. We verify the final claim for $B_2/P_{1,2}$ directly. ☐

Remark 4.3.3 (Semisimple case). Part (a) of Lemma 4.3.2 identifies the exceptions for all Type I modules. Part (b) indicates that exceptional Type II modules must be associated with factors in $\mathfrak{g}/\mathfrak{p}$ of the form $A_1/\mathfrak{p}_1\times\mathfrak{g}'/\mathfrak{p}'$ and $w=(1,k')$, where the 1 comes from the first factor, $k'\in I_{\mathfrak{p}'}$ is arbitrary, and $\lambda=2\lambda_1$, i.e. the highest weight of A_1. (Note $\phi_0=e_{\alpha_1}\wedge e_{\alpha_{k'}}\otimes e_{\alpha_1}$, which has weight $-w\cdot\lambda$ and homogeneity $+3$.) There are no exceptions associated to Type III modules.

We first establish a result outside the parabolic context:

Lemma 4.3.4. *Let G be a real or complex Lie group, and $P\subset G$ a closed subgroup. Let $\mathfrak{f}$ be a Lie algebra, $\mathfrak{k}\subset\mathfrak{f}$ a subalgebra, and $\iota:\mathfrak{k}\to\mathfrak{p}$ a Lie algebra injection. Suppose there exists a linear map $\vartheta:\mathfrak{f}\to\mathfrak{g}$ such that: (i) $\vartheta|_{\mathfrak{k}}=\iota$; (ii) $\vartheta([z,x]_{\mathfrak{f}})=[\iota(z),\vartheta(x)]$, $\forall z\in\mathfrak{k}$ and $\forall x\in\mathfrak{f}$; (iii) ϑ induces an isomorphism from $\mathfrak{f}/\mathfrak{k}$ to $\mathfrak{g}/\mathfrak{p}$. Then there exists a G/P geometry $(\mathcal{G}\to M,\omega)$ with $\mathfrak{inf}(\mathcal{G},\omega)$ containing a subalgebra isomorphic to $\mathfrak{f}$.*

Proof. Let K and F be the unique connected, simply-connected Lie groups with Lie algebras $\mathfrak{k}$ and $\mathfrak{f}$, respectively. There are unique Lie group homomorphisms $\Phi:K\to F$ and $\Psi:K\to P$ whose differentials $\Phi':\mathfrak{k}\to\mathfrak{f}$ and $\Psi':\mathfrak{k}\to\mathfrak{p}$ are the given inclusions $\mathfrak{k}\hookrightarrow\mathfrak{f}$ and $\iota:\mathfrak{k}\hookrightarrow\mathfrak{p}$.

Case 1: Φ is injective and $\Phi(K) \subset F$ is a Lie subgroup. Define $\mathcal{G} = F \times_K P = (F \times P)/\sim_K$, where $(u, p) \sim_K (u \cdot \Phi(k), \Psi(k^{-1}) \cdot p)$ for any $k \in K$. This has a natural projection $\mathcal{G} \to M := F/\Phi(K)$ (note $\Phi(K) \subset F$ is closed). Since Φ is injective, identify K with its image $\Phi(K)$. Any F-invariant Cartan connection ω on $\mathcal{G}$ corresponds [26], [10] to a linear map $\vartheta : \mathfrak{f} \to \mathfrak{g}$ satisfying (i)–(iii). The curvature function of $\omega = \omega_\vartheta$ corresponds to

$$\kappa_\vartheta(x, y) = [\vartheta(x), \vartheta(y)] - \vartheta([x, y]_{\mathfrak{f}}) \qquad \forall x, y \in \mathfrak{f}. \tag{4.5}$$

By construction, $\mathfrak{f} \subset \mathfrak{inf}(\mathcal{G}, \omega)$.

Case 2: Φ is not injective or $\Phi(K) \subset F$ is not a Lie subgroup. Since Φ' is injective, then Φ has a discrete kernel. In place of F in the definition of the Cartan bundle $\mathcal{G}$ in Case 1, we will define a *semi-local* Lie group U, an inclusion $K \hookrightarrow U$ as a closed subgroup which acts on the right on U, and such that $\mathrm{Lie}(U) = \mathfrak{f}$. ("Semi-locality" will be clarified below.) Then $\mathfrak{f}$-invariant Cartan connections on $\mathcal{G} = U \times_K P \to M := U/K$ correspond to linear maps $\vartheta : \mathfrak{f} \to \mathfrak{g}$ as in Case 1.

Let $\mathfrak{v}$ be a vector space complementary to $\mathfrak{k}$ in $\mathfrak{f}$. Choose two norms $\|\cdot\|_{\mathfrak{v}}$, $\|\cdot\|_{\mathfrak{k}}$ on these vector spaces and define the norm on the Lie algebra $\mathfrak{f} = \mathfrak{v} \oplus \mathfrak{k}$ by the formula $\|v + k\|_{\mathfrak{f}} = \max\{\|v\|_{\mathfrak{v}}, \|k\|_{\mathfrak{k}}\}$ for $v \in \mathfrak{v}$, $k \in \mathfrak{k}$. Then the balls in these spaces satisfy: $B^{\mathfrak{f}}_\epsilon = B^{\mathfrak{v}}_\epsilon \times B^{\mathfrak{k}}_\epsilon$.

We choose $\epsilon > 0$ so small that the maps $\exp : B^{\mathfrak{k}}_{5\epsilon} \to K$, $\exp : B^{\mathfrak{f}}_{5\epsilon} \to F$ are injective, $\exp(B^{\mathfrak{k}}_{5\epsilon}) \cap \Phi^{-1}(1_F) = 1_K$, and we will further specify ϵ below. One of the requirements is that $\exp \cdot \exp : B^{\mathfrak{v}}_\epsilon \times B^{\mathfrak{k}}_\epsilon \to F$ is an embedding. This holds for small ϵ because $\log\bigl(\exp(\epsilon v)\exp(\epsilon k)\exp(-\epsilon v - \epsilon k)\bigr) = o(\epsilon)$ by the Baker–Campbell–Hausdorff formula ($\|v\|_{\mathfrak{v}} \leq 1$, $\|k\|_{\mathfrak{k}} \leq 1$), implying that for $\epsilon \ll 1$, $U_0 := \exp(B^{\mathfrak{v}}_\epsilon) \cdot \exp(B^{\mathfrak{k}}_\epsilon) \subset \exp(B^{\mathfrak{f}}_{2\epsilon})$ and $U_0^{-1} = \{g^{-1} \mid g \in U_0\} = \exp(B^{\mathfrak{k}}_\epsilon) \cdot \exp(B^{\mathfrak{v}}_\epsilon) \subset \exp(B^{\mathfrak{f}}_{2\epsilon})$.

Let $K_0 = \exp(B^{\mathfrak{k}}_\epsilon) \subset K$ and $V_0 = \Phi(K_0) \subset U_0$. Denote $U_0^2 := U_0^{-1} \cdot U_0 \subset F$ and $V_0^2 = (U_0^2 \cap \mathrm{im}(\Phi))_0$, where the last zero means the connected component of 1_F. We have $V_0^2 = \Phi(K_0^2)$ for a unique connected neighborhood K_0^2 of 1_K. Denote also $K_0^4 := K_0^2 \cdot K_0^2$. Using the Baker–Campbell–Hausdorff formula as above we get $U_0^2 \subset \exp(B^{\mathfrak{f}}_{3\epsilon})$, $K_0^2 \subset \exp(B^{\mathfrak{k}}_{3\epsilon})$ and $K_0^4 \subset \exp(B^{\mathfrak{k}}_{5\epsilon})$ for small enough ϵ. This implies that $\Phi : K_0^4 \to V_0^4 = V_0^2 \cdot V_0^2 \subset F$ is an embedding and that $V_0^4 \cap U_0^2 = V_0^2$. Indeed, from $V_0^4 \subset \exp(B^{\mathfrak{f}}_{5\epsilon})$ we get: $\log(V_0^4 \cap U_0^2) \subset \mathfrak{k} \cap \log(U_0^2) = \log(V_0^2) \subset \mathfrak{f}$. Consequently $V_0^4 \cap U_0^2 \subset V_0^2$ and the reverse inclusion is obvious.

Consider the sheaf of neighborhoods $\{U_k = U_0 \cdot \Phi(k)\}_{k \in K}$ over K. We introduce an equivalence relation on it as follows. Take $x_1 \in U_{k_1}$ and $x_2 \in U_{k_2}$, i.e. $x_1 = u_1 \cdot \Phi(k_1)$, $x_2 = u_2 \cdot \Phi(k_2)$ for some $u_1, u_2 \in U_0$. We let $x_1 \sim x_2$ iff $u_2 = u_1 \cdot \Phi(k_1 k_2^{-1})$ and $k_1 k_2^{-1} \in K_0^2$.

Let us check that it is an equivalence relation. That $\sim$ is reflexive is obvious and the symmetry follows from $(K_0^2)^{-1} = K_0^2 \Leftrightarrow (U_0^2)^{-1} = U_0^2$. Let now $x_1 \sim x_2$ and $x_2 \sim x_3$. Then $u_3 = u_2 \cdot \Phi(k_2 k_3^{-1}) = u_1 \cdot \Phi(k_1 k_2^{-1}) \cdot \Phi(k_2 k_3^{-1}) = u_1 \cdot \Phi(k_1 k_3^{-1})$ and $k_1 k_3^{-1} = (k_1 k_2^{-1})(k_2 k_3^{-1}) \in K_0^4$. On the other hand $\Phi(k_1 k_3^{-1}) = u_1^{-1} u_3 \in U_0^2$, and these two imply $k_1 k_3^{-1} \in K_0^2$. Thus we get transitivity.

Define $U = \sqcup_{k \in K} U_k/\sim$. This is a manifold with the chart centered at k given by the embedding $R_{\Phi(k)} : U_0 \to U$. Overlap maps are given by the equivalence relation. The group K is embedded into U by the formula $K \ni k \mapsto R_{\Phi(k)}(1_F) \in U_k \hookrightarrow U$. Indeed, if $k \sim k'$ i.e. $1_F \cdot \Phi(k) = 1_F \cdot \Phi(k')$ and $k' k^{-1} \in K_0^2$, then $\Phi(k' k^{-1}) = 1_F$ yields $k = k'$. This implies that topologically $U \simeq B^{\mathfrak{v}}_\epsilon \times K$.

In fact, we can make this isomorphism canonical by strengthening the condition on ϵ (taking it possibly smaller). Namely let us require that $\exp \cdot \exp : B^{\mathfrak{v}}_\epsilon \times B^{\mathfrak{k}}_{5\epsilon} \to F$ is an embedding. Every element of U can be represented in the form $U_k \ni u_0 \cdot \Phi(k) = \exp(\epsilon v) \cdot \Phi(\exp(\epsilon\kappa) \cdot k)$, where $\|v\|_{\mathfrak{v}} \leq 1$ and $\|\kappa\|_{\mathfrak{k}} \leq 1$. We map this element to $(\epsilon v, \exp(\epsilon\kappa) \cdot k)$. Independence of the representative follows from the above condition on the map $\exp \cdot \exp$.

Moreover, U is a semi-local Lie group, which means that for any two points $k_1, k_2 \in K$ there exists two neighborhoods $U'_{k_i} \subset U_{k_i}$ containing $\exp(B^{\mathfrak{k}}_\epsilon) \cdot \Phi(k_i)$ in U such that the multiplication (induced from F) acts as $U'_{k_1} \times U'_{k_2} \to U_{k_1 \cdot k_2} \subset U$ by $(u_1 \cdot \Phi(k_1)) \cdot (u_2 \cdot \Phi(k_2)) = (u_1 \cdot Ad_{\Phi(k_1)} u_2) \cdot \Phi(k_1 k_2)$. Notice that $u_1 \cdot Ad_{\Phi(k_1)} u_2 \in U_0$ if $u_1, u_2 \in U_0$ are sufficiently close to 1_F. (This condition specifies U'_{k_i}.)

Similarly, we impose locality of the inverse along K, as a map $U'_k \to U_{k^{-1}}$ for some neighborhood $U'_k \subset U_k$ of $\exp(B^{\mathfrak{k}}_\epsilon) \cdot \Phi(k)$ given by $(u \cdot \Phi(k))^{-1} = Ad_{\Phi(k^{-1})}\, u^{-1} \cdot \Phi(k^{-1})$.

The semi-local Lie group U is naturally homomorphically mapped to F (this is an immersion, not necessarily an embedding). The chart $U_{1_K} \hookrightarrow U$ is isomorphically mapped to $U_0 \subset F$. Therefore the Lie algebra of U is naturally identified with $\mathfrak{f}$.

In addition, $K \hookrightarrow U$ is a closed subgroup, which acts from the right on U. The orbit of the right K-action is given by: $K \ni k \mapsto u_0 \cdot \Phi(k) \in U_k \hookrightarrow U$, $u_0 \in U_0$. All orbits are isomorphic to K and the two orbits either do not intersect or coincide. Moreover (provided we adopt the stronger condition on ϵ above) we can represent the orbit uniquely via u_0 by claiming that $u_0 \in \exp(B^{\mathfrak{v}}_\epsilon)$. The orbit space is then $U/K = U_0/K_0 = \exp(B^{\mathfrak{v}}_\epsilon) \simeq B^{\mathfrak{v}}_\epsilon$, and this is a neighborhood of the point $K/K \equiv 0 \in B^{\mathfrak{v}}_\epsilon$. Thus we get the local homogeneous space as the desired model.

Now define the Cartan bundle $\mathcal{G} = U \times_K P = (U \times P)/\sim_\Psi$, where $(q_1, p_1) \sim_\Psi (q_2, p_2)$ iff $q_2 = q_1 \cdot k$ and $p_2 = \Psi(k)^{-1} \cdot p_1$ for some $k \in K$. This $\mathcal{G}$ is a principal P-bundle over $U/K \simeq B^{\mathfrak{v}}_\epsilon$, in particular $\mathcal{G} \simeq B^{\mathfrak{v}}_\epsilon \times P$ topologically. We conclude as in Case 1. □

Theorem 4.3.5 (Realizability). *Let G be a complex semisimple Lie group, $P \subset G$ a parabolic subgroup, λ the highest weight of a simple ideal in $\mathfrak{g}$, $w \in W^{\mathfrak{p}}(2)$, and $\mu = -w \cdot \lambda$. Suppose $w(-\lambda) \in \Delta^-$. Then there exists a normal G/P geometry $(\mathcal{G} \to M, \omega)$ such that:*

(a) $\mathrm{im}(\kappa_H) \subset \mathbb{V}_\mu$*;*
(b) $\dim(\mathfrak{inf}(\mathcal{G}, \omega)) \geq \dim(\mathfrak{a}(\mu))$*;*
(c) the geometry is regular iff μ has positive homogeneity.

Proof. Define $\mathfrak{f}$ to be the vector space $\mathfrak{a} := \mathfrak{a}(\mu)$ with deformed bracket (4.3). Since $w(-\lambda) \in \Delta^-$, then by Lemma 4.3.1, $\mathfrak{f}$ is a Lie algebra. Since ϕ_0 is trivial on $\mathfrak{k} := \mathfrak{a}_{\geq 0}$, then $\mathfrak{k} \subset \mathfrak{f}$ is a subalgebra. Also, $\mathfrak{k} \subset \mathfrak{p} = \mathfrak{g}_{\geq 0}$ is a subalgebra. Since $\mathfrak{f} = \mathfrak{a}$ as vector spaces, and $\mathfrak{a} \subset \mathfrak{g}$, take $\vartheta : \mathfrak{f} \to \mathfrak{g}$ to be the induced *vector space* inclusion. Then (i)–(iii) in Lemma 4.3.4 hold, so an $\mathfrak{f}$-invariant G/P geometry $(\mathcal{G} \to M, \omega_\vartheta)$ exists (so (b) holds) with curvature function determined by

$$\kappa_\vartheta(x, y) = [\vartheta(x), \vartheta(y)] - \vartheta([x, y]_{\mathfrak{f}}) = [x, y] - [x, y]_{\mathfrak{f}} = \phi_0(x, y), \qquad \forall x, y \in \mathfrak{f}.$$

Since $\phi_0 \in \ker(\Box)$, then $\partial^* \phi_0 = 0$, so $(\mathcal{G} \to M, \omega)$ is normal, and (a), (c) are immediate. □

Recall $\mathfrak{S}_\mu$ and $\mathfrak{U}_\mu$ from Remark 4.2.9. If $\mathfrak{g}$ is simple, we also write $\mathfrak{a}(w) := \mathfrak{a}(-w \cdot \lambda_{\mathfrak{g}})$.

Example 4.3.6. *For $B_2/P_{1,2}$, $\lambda_{\mathfrak{g}} = 2\lambda_2$. Both $w = (12)$ and $w' = (21)$ are PR with* $\dim(\mathfrak{a}(w)) = \dim(\mathfrak{a}(w')) = 5$*, so $\mathfrak{S} \leq 5$. Since $w'(-\lambda_{\mathfrak{g}}) \in \Delta^-$, then by Theorem 4.3.5, a model with 5-dimensional symmetry exists, so $\mathfrak{S} = 5$. This has twistor space type B_2/P_2 (also non-exceptional). We will see in Section 4.5 that 5 is not realizable in the $w = (12)$ branch, having twistor type B_2/P_1.*

For *split real forms*, notions of roots and weights from complex representation theory are similarly defined. All preceding statements and proofs continue to hold. Combining Theorems 4.2.5 and 4.3.5, Corollary 4.2.8 and Remark 4.2.9, together with Lemma 4.3.2, we obtain:

Theorem 4.3.7. *Let G be a complex or split-real semisimple Lie group, $P \subset G$ a parabolic subgroup, λ the highest weight of a simple ideal in $\mathfrak{g}$, $w \in W^{\mathfrak{p}}(2)$, and $\mu = -w \cdot \lambda$ satisfies $Z(\mu) > 0$. We always have $\mathfrak{S}_\mu \leq \mathfrak{U}_\mu$. If $w(-\lambda) \in \Delta^-$ or G/P does not contain the factors A_1/P_1, A_2/P_1, $A_2/P_{1,2}$, B_2/P_1, $B_2/P_{1,2}$, then $\mathfrak{S}_\mu = \mathfrak{U}_\mu =$* $\dim(\mathfrak{a}(\mu))$ *and any submaximally symmetric model is locally homogeneous about a non-flat regular point.*

From Theorem 4.2.5 and (3.2), $\mathfrak{S} \leq \mathfrak{U} = \max_\mu \mathfrak{U}_\mu$, and aside from a few exceptions, this equals $\max_\mu \mathfrak{S}_\mu$ (where all μ are of the form indicated in the theorem above), which implies $\mathfrak{S} = \mathfrak{U}$. By Example 4.3.6, this also holds for $B_2/P_{1,2}$. Thus, we have:

Theorem 4.3.8. *Let G be a complex or split-real semisimple Lie group, $P \subset G$ a parabolic subgroup. If G/P does not contain the factors A_1/P_1, A_2/P_1, $A_2/P_{1,2}$, B_2/P_1, then*

$$\mathfrak{S} = \mathfrak{U} = \max\{\mathfrak{U}_\mu \mid \mathbb{V}_\mu \subset H^2_+(\mathfrak{g}_-, \mathfrak{g})\}, \qquad \mathfrak{U}_\mu = \dim(\mathfrak{a}(\mu)),$$

and any submaximally symmetric model is locally homogeneous about a non-flat regular point.

Remark 4.3.9. In the general (complex or split-real) semisimple case, the gap problem reduces to studying each $\mathbb{V}_\mu$, which is one of the three types in Table 4. Each such $\mathbb{V}_\mu$ is essentially generated by at most two simple factors: any remaining simple factor $(\hat{\mathfrak{g}}, \hat{\mathfrak{p}})$ yields a flat factor for the geometry, so $\dim(\hat{\mathfrak{g}})$ is automatically included in the submaximal symmetry dimension count.

Remark 4.3.10. Aside from the exceptions in Theorem 4.3.7, given $(\mathfrak{g}, \mathfrak{p}, \mu)$, $\mathfrak{S}_\mu$ is the same for all choices of Lie groups (G, P) with $\mathrm{Lie}(G) = \mathfrak{g}$ and $\mathrm{Lie}(P) = \mathfrak{p}$. Consequently, for the purpose of simplifying the calculation of $\mathfrak{S}_\mu$, we may assume that P is *connected* so that we can pass to the minimal twistor space. (See Remark 2.7.3 and Corollary 2.7.2.)

4.4. **Deformations.** A graded subalgebra $\mathfrak{a} \subset \mathfrak{g}$ admits a canonical filtration, i.e. $\mathfrak{a}^i = \bigoplus_{j \geq i} \mathfrak{a}_j$.

Definition 4.4.1. *A graded subalgebra $\mathfrak{a}$ is* filtration-rigid *if any filtered Lie algebra $\mathfrak{f}$ with* $\mathrm{gr}(\mathfrak{f}) = \mathfrak{a}$ *as graded Lie algebras has $\mathfrak{f} \cong \mathfrak{a}$ as filtered Lie algebras.*

Proposition 4.4.2. *Let G be a real or complex semisimple Lie group and $P \subset G$ a parabolic subgroup, and suppose that* $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0) \cong \mathfrak{g}$. *Let $(\mathcal{G} \xrightarrow{\pi} M, \omega)$ be a regular, normal G/P geometry which is not locally flat. Given $u \in \mathcal{G}$ with $\kappa_H(u) \neq 0$, if $\mathfrak{g}_- \subset \mathfrak{s}(u)$, then $\mathfrak{s}(u)$ is not filtration-rigid.*

Proof. Since $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0) \cong \mathfrak{g}$, the geometry is completely determined by its regular infinitesimal flag structure. Assume $\mathfrak{f}(u) \cong \mathfrak{s}(u)$, so $\mathfrak{f}(u)$ and hence $\mathcal{S}(x)$ is graded, where $x = \pi(u)$. Then $\mathcal{S}(x)/\mathcal{S}(x)^0 \cong \mathfrak{g}_- =: \mathfrak{m}$ acts transitively on M at x. Thus, M can be locally identified near x with the simply-connected Lie group $M(\mathfrak{m})$ with Lie algebra $\mathfrak{m}$. This is endowed with the $M(\mathfrak{m})$-invariant distribution generated by $\mathfrak{g}_{-1}$, and there is a $M(\mathfrak{m})$-invariant principal G_0-bundle $\mathcal{G}_0 \to M(\mathfrak{m})$, where $G_0 \subseteq \mathrm{Aut}_{gr}(\mathfrak{m})$, which is a reduction of the graded frame bundle over $M(\mathfrak{m})$.

The triple $(M(\mathfrak{m}), \mathfrak{g}_{-1}, \mathfrak{g}_0)$ is the standard model in Tanaka theory, and since $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0) \cong \mathfrak{g}$ is finite-dimensional, its symmetry algebra $\mathcal{S}$ is isomorphic to $\mathfrak{g}$ [40, Section 6]. But since $\kappa_H(u) \neq 0$, then $\dim(\mathcal{S}) \leq \mathfrak{S} < \dim(\mathfrak{g})$, c.f. Proposition 2.4.1 or Theorem 4.2.5, a contradiction. ☐

In the exceptional cases, we will show that $\mathfrak{a}(\mu)$ is filtration-rigid, and hence $\mathfrak{S}_\mu < \dim(\mathfrak{a}(\mu))$.

Example 4.4.3. *Without the* $\mathrm{pr}(\mathfrak{g}_-, \mathfrak{g}_0) \cong \mathfrak{g}$ *assumption, Proposition 4.4.2 generally fails. Consider projective geometry (A_ℓ/P_1). The condition $\mathfrak{g}_- \subset \mathfrak{s}(u)$ implies local homogeneity, and since $\mathfrak{g}_-$ is abelian, then near $\pi(u)$, we have local coordinates $(t, x^1, ..., x^m)$, where $m = \ell - 1$, and symmetries $\partial_t, \partial_{x^1}, ..., \partial_{x^m}$ of the projective structure. The geodesic equations* (5.11) *admit these symmetries, so all coefficients in these equations are constants. This means that by a projective change, we may take the Γ^a_{bc} to be constants. For $m \geq 2$, this does not imply the structure is flat, e.g. taking all $\Gamma^a_{bc} = 0$ except $\Gamma^1_{01} = -\frac{1}{2}$, the Fels invariants* (5.3) *satisfy $S \equiv 0$ and $T \neq 0$.*

However, for $m = 1$, the Tresse invariants (4.7) *vanish everywhere for $y'' = F(y')$, where F is a cubic polynomial. Thus, Proposition 4.4.2 is true for 2-dimensional projective structures.*

Consider $\mathfrak{a} = \bigoplus_i \mathfrak{a}_i$ with (graded) basis $\{e^\alpha_i\}$ and structure equations $[e^\alpha_i, e^\beta_j] = \sum_\gamma c^{\alpha\beta}_\gamma e^\gamma_{i+j}$. Any filtered Lie algebra $\mathfrak{f} \subset \mathfrak{g}$ with $\mathrm{gr}(\mathfrak{f}) = \mathfrak{a}$ has a (filtered) basis $\{f^\alpha_i\}$ with

$$[f^\alpha_i, f^\beta_j]_{\mathfrak{f}} = \sum_\gamma c^{\alpha\beta}_{ij\gamma} f^\gamma_{i+j} + \sum_{k > i+j} \sum_\gamma b^{\alpha\beta k}_{ij\gamma} f^\gamma_k, \tag{4.6}$$

and we consider $\mathfrak{f}$ up to filtration-preserving automorphisms

$$f_i^\alpha \mapsto f_i^\alpha + \sum_{j>i} \sum_\beta \chi_{i\beta}^{\alpha j} f_j^\beta.$$

We refer to the terms in the double summation in (4.6) as *tails*. So establishing filtration-rigidity amounts to eliminating the presence of tails (in some basis).

Example 4.4.4. *Let $\mathfrak{g}_0$ be a semi-simple Lie algebra, and $V = \mathfrak{g}_{-1}$ an irreducible $\mathfrak{g}_0$-representation such that $\bigwedge^2 V$ does not contain V nor the adjoint representation of $\mathfrak{g}_0$ as summands. Then the graded Lie algebra $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0$ is filtration-rigid. Indeed, let $\mathfrak{f}$ be a filtered Lie algebra with $\mathrm{gr}(\mathfrak{f}) = \mathfrak{g}$. Then $\mathfrak{f}^0 \cong \mathfrak{g}_0$. By semisimplicity of $\mathfrak{g}_0$, the $\mathfrak{g}_0$-module $\mathfrak{f}$ is completely reducible, and so $\mathfrak{f} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0$ as $\mathfrak{g}_0$-modules. By hypothesis, the bracket $\bigwedge^2 \mathfrak{g}_{-1} \to \mathfrak{g}$ being $\mathfrak{g}_0$-equivariant must vanish.*

The case of $\mathfrak{g}_0 = \mathfrak{so}(n)$ and $V = \mathbb{R}^n$ has $\bigwedge^2 V \simeq \mathfrak{g}_0$, so there are many filtered deformations. By Schur's lemma, we have deformations parametrized by $R \in \mathbb{R}$, and this yields all the maximally symmetric models in Table 1. The corresponding algebras depending on the sign of R are: $\mathfrak{g} = \mathfrak{so}(n+1)$, $\mathbb{R}^n \rtimes \mathfrak{so}(n)$, and $\mathfrak{so}(n,1)$.

Proposition 4.4.5. *Let $\mathfrak{f}$ be a filtered Lie algebra, $\mathrm{gr}(\mathfrak{f}) =: \mathfrak{a} = \mathfrak{a}_{-\hat{\nu}} \oplus \ldots \oplus \mathfrak{a}_0 \oplus \ldots \oplus \mathfrak{a}_{\tilde{\nu}}$ its associated-graded, and let $\tau_j : \mathfrak{f}^j \to \mathfrak{a}_j$ denote the canonical projections. Suppose there exists $h \in \mathfrak{a}_0$ such that ad_h is diagonal in some (graded) basis $\{e_i^\alpha\}$ of $\mathfrak{a}$, and let $\mathcal{E}_i = \mathrm{Spec}(\mathrm{ad}_h|_{\mathfrak{a}_i})$. If $\mathcal{E}_i \cap \mathcal{E}_j = \emptyset$ for distinct i, j, then for any $\tilde{h} \in \mathfrak{f}^0$ with $\tau_0(\tilde{h}) = h$, there is a (filtered) basis $\{f_i^\alpha\}$ of $\mathfrak{f}$, with $\tau_i(f_i^\alpha) = e_i^\alpha$, and with respect to which $\mathrm{ad}_{\tilde{h}}$ is diagonal. Moreover, if:*[11]

(i) $\forall i \neq j$: $(\mathcal{E}_i + \mathcal{E}_j) \cap \bigcup_{k>i+j} \mathcal{E}_k = \emptyset$; and

(ii) $\forall i$: $\bigwedge^2 \mathcal{E}_i \cap \bigcup_{k>2i} \mathcal{E}_k = \emptyset$,

then $\mathfrak{a}$ is filtration-rigid.

Proof. Use (reverse) induction on r, where $-\hat{\nu} \leq r \leq \tilde{\nu}$. The $r = \tilde{\nu}$ case is trivial since $\tau_{\tilde{\nu}}|_{\mathfrak{f}^{\tilde{\nu}}} : \mathfrak{f}^{\tilde{\nu}} \to \mathfrak{a}_{\tilde{\nu}}$ is a Lie algebra isomorphism. Assume there is a basis $\{f_i^\alpha\}_{i=r+1}^{\tilde{\nu}}$ of $\mathfrak{f}^{r+1}$ with $\tau_i(f_i^\alpha) = e_i^\alpha$ such that $\mathrm{ad}_{\tilde{h}}$ is diagonal (with eigenvalues μ_i^α). We have $[\tilde{h}, f_r^\alpha] = \mu_r^\alpha f_r^\alpha + \sum_{k>r} \sum_\gamma b_\gamma^{\alpha,k} f_k^\gamma$. Since $\mathcal{E}_i \cap \mathcal{E}_j = \emptyset$ for $i \neq j$, then $\mu_r^\alpha \neq \mu_{r+1}^\beta$, so letting $\hat{f}_r^\alpha = f_r^\alpha + \sum_\beta \sigma_\beta^\alpha f_{r+1}^\beta$ with $\sigma_\beta^\alpha = (\mu_r^\alpha - \mu_{r+1}^\beta)^{-1} b_\beta^{\alpha,r+1}$, we can normalize $b_\beta^{\alpha,r+1} = 0$. Iteratively, we normalize all $b_\gamma^{\alpha,k} = 0$.

We have $[\tilde{h}, [f_i^\alpha, f_j^\beta]] = (\mu_i^\alpha + \mu_j^\beta)[f_i^\alpha, f_j^\beta]$. If $i \neq j$, then by (i), there is no tail. If $i = j$, then for $\alpha \neq \beta$, (ii) implies there is no tail. Thus, $\mathfrak{a}$ is filtration-rigid. ☐

4.5. **Exceptional cases and local homogeneity.**

We now turn to the remaining exceptional cases from Lemma 4.3.2 and Theorem 4.3.7. Our first result in this section is:

Proposition 4.5.1. *For complex or real (regular, normal) A_2/P_1, $A_2/P_{1,2}$, B_2/P_1 geometries, $\mathfrak{S} = \mathfrak{U} - 1$, given in Table 7. For complex or split-real (regular, normal) $G/P = A_1/P_1 \times G'/P'$ geometries with κ_H concentrated in the (Type II) module $\mathbb{V}_\mu$, with $\mu = -w \cdot 2\lambda_1$ corresponding to $w = (1, k') \in W^{\mathfrak{p}}(2)$, we have $\mathfrak{U}_\mu - 1 \leq \mathfrak{S}_\mu \leq \mathfrak{U}_\mu$.*

The rank two exceptions when $\mathfrak{g}$ is simple are given in Table 7. These are all PR$_+$ by Recipe 7. By Example 4.3.6, $B_2/P_{1,2}$ is not exceptional concerning $\mathfrak{S}$, so we exclude this here.

For lower bounds, we exhibit models. For upper bounds, it suffices (by Remark 4.2.7) to work over $\mathbb{C}$, and we show that $\mathfrak{a}^{\phi_0} \subset \mathfrak{g}$ is filtration-rigid using Proposition 4.4.5. By Proposition 4.4.2, this forces $\mathfrak{S} = \mathfrak{U} - 1$ for all three cases. (By Example 4.4.3, this is valid in the A_2/P_1 case.) For the $A_1/P_1 \times G'/P'$ exception, we exhibit a model with at least $\mathfrak{U}_\mu - 1$ symmetries.

---

[11] Here $T + T := \{a + b \mid a, b \in T\}$ and $\bigwedge^2 T := \{a + b \mid a, b \in T,\ a \neq b\}$.

G/P	Underlying structures	$H^2_+(\mathfrak{g}_-, \mathfrak{g})$	$\mathfrak{S}$
A_2/P_1	2-dimensional projective structures		3
$A_2/P_{1,2}$	scalar 2nd order ODE up to point transformations (3-dimensional CR structures)		3
B_2/P_1	3-dimensional conformal Lorentzian (or Riemannian) structures		4

TABLE 7. Rank two exceptions and (complex or real) underlying structures

Our second result generalizes Corollary 4.2.8 and Remark 4.2.9, which considered the general case, i.e. when $\mathfrak{S} = \mathfrak{U}$ or $\mathfrak{S}_\mu = \mathfrak{U}_\mu$. Considering each exceptional case in turn, we prove:

Theorem 4.5.2. *Among all complex or split-real (regular, normal) G/P geometries, all submaximally symmetric models are locally homogeneous near a non-flat regular point. Given a $\mathfrak{g}_0$-irreducible $\mathbb{V}_\mu \subset H^2_+(\mathfrak{g}_-, \mathfrak{g})$, the same conclusion holds with the restriction that* $\text{im}(\kappa_H) \subset \mathbb{V}_\mu$.

Given a regular point x and $u \in \pi^{-1}(x)$, this requires showing that $\mathfrak{g}_- \subset \mathfrak{s}(u)$.

Remark 4.5.3. Since x is regular, then $\mathfrak{s}(u) \subset \mathfrak{a}^{\kappa_H(u)}$. If $\kappa_H(u)$ is not in the G_0-orbit of $\phi_0 \in \mathbb{V}_\mu$, then $\dim(\mathfrak{a}^{\kappa_H(u)}) < \dim(\mathfrak{a}^{\phi_0}) = \mathfrak{U}_\mu$ by Lemma 3.1.1. In this case, if $\dim(\mathfrak{s}(u)) \geq \mathfrak{U}_\mu - 1$, then $\mathfrak{s}(u) = \mathfrak{a}^{\kappa_H(u)} \supset \mathfrak{g}_-$. Thus, it suffices to consider $\kappa_H(u)$ in the G_0-orbit of ϕ_0, and by acting on u by $G_0 \subset P$, we may assume $\kappa_H(u) = \phi_0$, and $\mathfrak{s}(u) \subsetneq \mathfrak{a}^{\phi_0}$ has codimension one.

Since $\mathfrak{g}_- \subset \mathfrak{g}$ is bracket-generating, then intransitivity ($\mathfrak{g}_- \not\subset \mathfrak{s}(u)$) would mean $\mathfrak{s}_{-1}(u) \neq \mathfrak{g}_{-1}$, and $\mathfrak{s}_0(u) = \mathfrak{a}_0$. Now recall from Proposition 4.2.2 that $[\mathfrak{s}_0(u), \mathfrak{g}_{-1}] \subset \mathfrak{s}_{-1}(u) \neq \mathfrak{g}_{-1}$. In each case, we will verify that $[\mathfrak{a}_0, \mathfrak{g}_{-1}] = \mathfrak{g}_{-1}$. This yields a contradiction, thereby establishing local homogeneity.

4.5.1. *2-dimensional projective structures.* Consider ∇ specified in coordinates $(x^0, x^1) = (x, y)$ by

$$\Gamma^0_{11} = \frac{\varepsilon}{x}, \qquad \Gamma^0_{00} = -\frac{1}{2x},$$

and zero otherwise, where $\varepsilon = \pm 1$. This corresponds by (5.11) to the ODE $xy'' = \varepsilon(y')^3 - \frac{1}{2}y'$. We have $\mathfrak{S} \geq 3$ since the projective structure $[\nabla]$ has symmetries (i.e. point symmetries of the ODE):

$$\mathbf{X}_1 = \partial_y, \qquad \mathbf{X}_2 = x\partial_x + y\partial_y, \qquad \mathbf{X}_3 = 2xy\partial_x + y^2\partial_y.$$

For A_2/P_1, $H^2_+(\mathfrak{g}_-, \mathfrak{g}) =$  $\cong \bigwedge^2 \mathfrak{g}_1 \otimes \mathfrak{g}_1$. Indeed, κ_H is the *Liouville tensor* $L = (L_1 dx + L_2 dy) \otimes (dx \wedge dy)$. Noting $5\lambda_1 - \lambda_2 = 3\alpha_1 + \alpha_2 = 3(\epsilon_1 - \epsilon_2) + \epsilon_2 - \epsilon_3 = 0$ and $\epsilon_1 + \epsilon_2 + \epsilon_3 = 0$,

$$\mathfrak{a} = \mathfrak{a}_{-1} \oplus \mathfrak{a}_0 = \left\{ \left(\begin{array}{c|cc} h & 0 & 0 \\ \hline u & 4h & 0 \\ v & s & -5h \end{array} \right) \right\} = \text{span}\{e^1_{-1}, e^2_{-1}, s_0, h_0\}.$$

We have ad_{h_0} diagonal with $\mathcal{E}_{-1} = \{3, -6\}$, $\mathcal{E}_0 = \{-9, 0\}$. Hence, $\mathfrak{a}$ is filtration-rigid, and $\mathfrak{S} \leq 3$. Clearly, $[\mathfrak{a}_0, \mathfrak{g}_{-1}] = \mathfrak{g}_{-1}$, hence local homogeneity follows.

4.5.2. *Scalar 2nd order ODE.* Given $y'' = F(x, y, y')$, there are the Tresse relative invariants [42]:

$$I_1 = \left(\frac{d}{dx}\right)^2 (F_{pp}) - F_p \frac{d}{dx}(F_{pp}) - 4\frac{d}{dx}(F_{yp}) + 4F_p F_{yp} - 3F_y F_{pp} + 6F_{yy}, \qquad I_2 = F_{pppp}, \tag{4.7}$$

where $p = y'$, and $\frac{d}{dx} = \partial_x + p\partial_y + F\partial_p$. There are two lines $L', L'' \subset H^2_+(\mathfrak{g}_-, \mathfrak{g})$ associated to $w = (12)$ and (21). Two-dimensional projective structures correspond to ODE with $I_2 = 0$.

Suppose the geometry is not locally flat. The set $U = \{z \in M \mid I_1(z) I_2(z) \neq 0\}$ is open. If $U \neq \emptyset$, it contains a regular point z, and for any $u \in \pi^{-1}(z)$, $\mathfrak{a}^{\kappa_H(u)} = \mathfrak{g}_-$, so $\dim(\mathcal{S}) \leq 3$. If $U = \emptyset$, then exactly one of the open sets $U_i = \{z \in M \mid I_i(z) \neq 0\}$ is non-empty. If $U_2 = \emptyset$, then by Section 4.5.1, $\dim(\mathcal{S}) \leq 3$. By the duality $A_\ell/P_1 \cong A_\ell/P_2$, we conclude the same if $U_1 = \emptyset$. Hence, $\mathfrak{S} = 3$. Local homogeneity follows because the same holds for 2-dimensional projective structures.

Example 4.5.4. *Given* $0 \neq a \in \mathbb{R}$, $y'' = \frac{1}{x}\left(y' - (y')^3 + a(1-(y')^2)^{3/2}\right)$ *admits point symmetries*

$$\mathbf{X}_1 = \partial_y, \qquad \mathbf{X}_2 = x\partial_x + y\partial_y, \qquad \mathbf{X}_3 = 2xy\partial_x + (x^2+y^2)\partial_y,$$

so is submaximally symmetric. Generically, $I_1 I_2 \neq 0$, *so* $\mathrm{im}(\kappa_H)$ *is not contained in only one of* L' *or* L''. *On* (x, y, p)*-space,* $\mathcal{S}$ *is obtained by prolongation – see* (5.6). *This gives* $\mathbf{X}_1^{(1)} = \mathbf{X}_1$, $\mathbf{X}_2^{(1)} = \mathbf{X}_2$, *and* $\mathbf{X}_3^{(1)} = \mathbf{X}_3 + 2x(1-p^2)\partial_p$. *Since* $x \neq 0$, $\mathcal{S}$ *is generally transitive, except along* $p = \pm 1$. *Thus, the singular set consists of: (i) points for which the right hand side of the ODE is not smooth (violating our blanket convention), and (ii) irregular points for* $\mathcal{S}$.

The other real form of $A_2/P_{1,2}$ geometries yields 3-dimensional CR structures. In [12], Cartan gave many examples of such structures having 3-dimensional symmetry. Thus, $\mathfrak{S} = 3$ here as well.

4.5.3. *3-dimensional conformal structures.* Note the models:

	Metric	Killing symmetries
Riemannian	$dx^2 + dy^2 + (dz - xdy)^2$	$\partial_y, \quad \partial_z, \quad \partial_x + y\partial_z, \quad y\partial_x - x\partial_y + \frac{1}{2}(y^2 - x^2)\partial_z$
Lorentzian	$dxdy + (dz - xdy)^2$	$\partial_y, \quad \partial_z, \quad \partial_x + y\partial_z, \quad x\partial_x - y\partial_y$

These have nonzero Cotton tensor, so are not locally conformally flat. Hence, $\mathfrak{S} \geq 4$.

For B_2/P_1, $H^2_+(\mathfrak{g}_-, \mathfrak{g}) = \overset{-5}{\times}\!\Rightarrow\!\overset{4}{\circ}$, and $5\lambda_1 - 4\lambda_2 = 3\alpha_1 + \alpha_2 = 3(\epsilon_1 - \epsilon_2) + \epsilon_2 = 3\epsilon_1 - 2\epsilon_2$:

$$\mathfrak{a} = \mathfrak{g}_{-1} \oplus \mathfrak{a}_0 = \left\{ \left(\begin{array}{c|ccc|c} 2h & 0 & 0 & 0 & 0 \\ \hline u & 3h & 0 & 0 & 0 \\ v & s & 0 & 0 & 0 \\ w & 0 & -s & -3h & 0 \\ \hline 0 & -w & -v & -u & -2h \end{array} \right) \right\} = \mathrm{span}\{e^1_{-1}, e^2_{-1}, e^3_{-1}, s_0, h_0\}.$$

We have ad_{h_0} diagonal, $\mathcal{E}_{-1} = \{1, -2, -5\}$, $\mathcal{E}_0 = \{-3, 0\}$. Hence, $\mathfrak{a}$ is filtration-rigid, and $\mathfrak{S} \leq 4$. Clearly, $[\mathfrak{a}_0, \mathfrak{g}_{-1}] = \mathfrak{g}_{-1}$, hence local homogeneity follows. This handles the Lorentzian case.

The Riemannian case is simpler. Here, $\mathfrak{g}_0 = \mathbb{R} \oplus \mathfrak{so}(3)$, and $H^2_+(\mathfrak{g}_-, \mathfrak{g})$ is isomorphic to the space of 3×3 symmetric trace-free matrices with action $A \cdot \phi = [A, \phi]$. Since any such ϕ is orthogonally diagonalizable, we get $\mathfrak{S} \leq \mathfrak{U} = 4$ from considering the two non-trivial cases:

(1) ϕ has three distinct eigenvalues: $\mathfrak{ann}(\phi) = 0$, so $\mathfrak{a}^\phi = \mathfrak{g}_-$, and $\dim(\mathfrak{a}^\phi) = 3$.
(2) ϕ has two equal nonzero eigenvalues: $\dim(\mathfrak{ann}(\phi)) = 1$, so $\dim(\mathfrak{a}^\phi) = 4$.

4.5.4. *Semisimple exceptions.* Consider $G/P = A_1/P_1 \times G'/P'$, $w = (1, k') \in W^{\mathfrak{p}}(2)$, and highest weight $2\lambda_1$ of A_1. Since we will consider only geometries with κ_H concentrated in $\mathbb{V}_\mu$ with

$$\mu = -w \cdot 2\lambda_1 = -\sigma_1 \cdot 2\lambda_1 - \sigma_{k'} \cdot 0' = 4\lambda_1 + \alpha_{k'} = 2\alpha_1 + \alpha_{k'},$$

we may assume that G' is simple. The lowest weight vector of $\mathbb{V}_\mu$ is $\phi_0 = e_{\alpha_1} \wedge e_{\alpha_{k'}} \otimes e_{\alpha_1}$, and $\mathfrak{U}_\mu = \dim(\mathfrak{a}^{\phi_0})$. We will construct a model with at least $\mathfrak{U}_\mu - 1$ symmetries.

By Theorem 3.5.1, the Lie algebra structure of $\mathfrak{a}^{\phi_0}$ is the same if we consider $\mathfrak{p}' = \mathfrak{p}_{k'}$ to be maximal parabolic. Then by Proposition 3.3.6, $(\mathfrak{g}, \mathfrak{p}, \mu)$ is PR. Take a basis H, X, Y of $\mathfrak{sl}_2(\mathbb{C})$ with the standard relations $[H, X] = X$, $[H, Y] = -Y$, $[X, Y] = H$. Identifying $e_{-\alpha_1}$ with Y, its Killing dual e_{α_1} is $\frac{X}{2}$. We have $\alpha_1(H) = 1$, i.e. H acts as the grading element with respect to P_1. Thus, we have $\mathfrak{a} = \mathfrak{a}^{\phi_0} = \mathfrak{g}_- \oplus \mathfrak{a}_0^{\phi_0}$, where $\mathfrak{g}_- = \mathrm{span}\{Y\} \oplus \mathfrak{g}'_-$, and

$$\mathfrak{a}_0 = \mathfrak{h}_0 \oplus \bigoplus_{\gamma \in \Delta(\mathfrak{g}'_{0,\leq 0})} \mathfrak{g}'_\gamma, \qquad \mathfrak{h}_0 := \left\{ -\frac{1}{2}\alpha_{k'}(H')H + H' \mid H' \in \mathfrak{h}' \right\}.$$

Let $\ker(\alpha_{k'}) = \{H' \in \mathfrak{h}' \mid \alpha_{k'}(H') = 0\}$, and define

$$\tilde{\mathfrak{a}}_0' = \ker(\alpha_{k'}) \oplus \bigoplus_{\gamma \in \Delta(\mathfrak{g}'_{0,\leq 0})} \mathfrak{g}'_\gamma, \qquad \tilde{\mathfrak{a}} = \mathfrak{g}_- \oplus \tilde{\mathfrak{a}}_0' = \mathrm{span}\{Y\} \oplus \mathfrak{g}'_- \oplus \tilde{\mathfrak{a}}_0'.$$

Similar to Section 4.3, we define $\tilde{\mathfrak{f}}$ to be the vector space $\tilde{\mathfrak{a}}$ with the same brackets, but with the single additional relation $[Y, e_{-\alpha_{k'}}]_{\tilde{\mathfrak{f}}} = Y$. Equivalently, $[Y, u]_{\tilde{\mathfrak{f}}} = \lambda(u) Y$, for all $u \in \tilde{\mathfrak{f}}$, where $\lambda \in e_{\alpha_{k'}}$ is Killing dual to $e_{-\alpha_{k'}}$. For $\tilde{\mathfrak{f}}$ to be a Lie algebra, it suffices to check $0 = \mathrm{Jac}_{\tilde{\mathfrak{f}}}(Y, u, v)$ for all $u, v \in \tilde{\mathfrak{f}}$, or equivalently, $\lambda([u, v]_{\tilde{\mathfrak{f}}}) = 0$. Clearly, this holds for $u, v \in \mathfrak{g}_-$. If $u \in \tilde{\mathfrak{a}}_0'$, then $u \cdot \phi_0 = 0$, which is equivalent to $\lambda \circ \mathrm{ad}_u = 0$ since $u \cdot e_{\alpha_1} = 0$. Thus, $\lambda([u, v]_{\tilde{\mathfrak{f}}}) = \lambda([u, v]) = 0$, so $\tilde{\mathfrak{f}}$ is a Lie algebra.

Let $\tilde{\mathfrak{k}} = \tilde{\mathfrak{a}}_0' \subset \tilde{\mathfrak{f}}$. Define the linear map $\vartheta : \tilde{\mathfrak{f}} \to \mathfrak{g}$ by $\vartheta = \mathrm{id} + H \otimes e_{\alpha_{k'}} - \frac{1}{2} X \otimes e_{\alpha_1}$. Thus, $\vartheta|_{\tilde{\mathfrak{k}}}$ is the natural inclusion, and ϑ maps all root vectors in $\mathfrak{g}_- \subset \tilde{\mathfrak{f}}$ over to $\mathfrak{g}$ naturally except for

$$\vartheta(Y) = Y - \frac{1}{2} X, \qquad \vartheta(e_{-\alpha_{k'}}) = e_{-\alpha_{k'}} + H.$$

For $u \in \tilde{\mathfrak{k}}$ and $v \in \tilde{\mathfrak{f}}$, since $Y, e_{-\alpha_{k'}} \perp \mathrm{im}(\mathrm{ad}_u)$, then $\vartheta([u, v]_{\tilde{\mathfrak{f}}}) = \vartheta([u, v]) = [u, v] = [u, \vartheta(v)] = [\vartheta(u), \vartheta(v)]$, so ϑ is $\tilde{\mathfrak{k}}$-equivariant. Also, ϑ induces a linear isomorphism $\tilde{\mathfrak{f}}/\tilde{\mathfrak{k}} \cong \mathfrak{g}/\mathfrak{p}$.

As in Section 4.3, the simplifying reduction to $\mathfrak{p}' = \mathfrak{p}_{k'}$ was mainly used to check that $\tilde{\mathfrak{f}}$ satisfies the Jacobi identity. Now for the original geometry, $\tilde{\mathfrak{a}} = \mathrm{span}\{Y\} \oplus \mathrm{pr}^{\mathfrak{g}'}(\mathfrak{g}'_-, \mathfrak{ann}(e_{\alpha_{k'}}))$ is the same $\tilde{\mathfrak{a}}$ as above (as Lie algebras), and similarly define $\tilde{\mathfrak{f}}$ and ϑ as above. Since $\tilde{\mathfrak{k}} = \tilde{\mathfrak{a}}'_{\geq 0}$ is contained in the $\tilde{\mathfrak{k}}$ stated above for $\mathfrak{p}_{k'}$, then ϑ defines an $\tilde{\mathfrak{f}}$-invariant Cartan connection ω_ϑ on some $\mathcal{G} = F \times_K P$ over $M = F/K$ (where F may only be a local Lie group). Using (4.5), κ_ϑ vanishes except for

$$\kappa_\vartheta(Y, e_{-\alpha_{k'}}) = [\vartheta(Y), \vartheta(e_{-\alpha_{k'}})] - \vartheta([Y, e_{-\alpha_{k'}}]_{\tilde{\mathfrak{f}}}) = \left[Y - \frac{1}{2} X, e_{-\alpha_{k'}} + H \right] - \vartheta(Y) = X,$$

i.e. $\kappa_\vartheta = \phi_0$. Thus, the geometry is regular and normal, and $\mathfrak{U}_\mu - 1 \leq \mathfrak{S}_\mu \leq \mathfrak{U}_\mu$. Finally, $[\mathfrak{a}_0, \mathfrak{g}_{-1}] \supset [\mathfrak{h}_0, \mathfrak{g}_{-1}] = \mathfrak{g}_{-1}$, so local homogeneity follows.

This completes the proof of both Proposition 4.5.1 and Theorem 4.5.2.

5. Results and local models for specific geometries

In this section, we illustrate the use of Theorems 4.3.7 and 3.2.6 for the computation of submaximal dimensions for specific geometries. We also refer the reader to Appendix B which summarizes our computational algorithm.

5.1. **Conformal geometry.** In dimension $n = p + q \geq 3$, conformal geometry is the prototypical underlying structure of a parabolic geometry. The model space is the (pseudo-) conformal sphere $\mathbb{S}^{p,q}$. Embedded as the (null) projective quadric in $\mathbb{P}(\mathbb{R}^{p+1,q+1})$, we have $\mathbb{S}^{p,q} \cong \mathrm{SO}_{p+1,q+1}/P_1$, where P_1 is the stabilizer of a null line. The Lie algebra $\mathfrak{g} = \mathfrak{so}_{p+1,q+1}$ is a real form of

$$\mathfrak{so}_{n+2}(\mathbb{C}) = \left\{ \begin{array}{ll} B_\ell, & n = 2\ell - 1 \text{ odd}; \\ D_\ell, & n = 2\ell - 2 \text{ even}. \end{array} \right.$$

and is 1-graded by P_1. The space $\mathbb{W} = H^2(\mathfrak{g}_-, \mathfrak{g})$ is the space of Weyl tensors (homogeneity $+2$) for $n \geq 4$ and Cotton tensors (homogeneity $+3$) for $n = 3$. The geometry is PR_+ by Corollary 3.3.8.

First work over $\mathbb{C}$, so $\mathfrak{g} = \mathfrak{so}_{n+2}(\mathbb{C})$, and $\mathfrak{g}_0^{ss}$ is either $B_{\ell-1}$ or $D_{\ell-1}$. For $H^2_+(\mathfrak{g}_-, \mathfrak{g})$, Kostant gives



Above, $W^{\mathfrak{p}}_+(2) = \{(12)\}$ always, except $W^{\mathfrak{p}}_+(2) = \{(12),(13)\}$ for D_3/P_1. The $n = 3$ case was studied in Section 4.5.3, so let $n \geq 4$. Then:

(1) $n \geq 5$: $\lambda_{\mathfrak{g}} = \lambda_2$, $w = (12)$, $J_w = \{3\}$. Look at $\mathfrak{p}_w^{\rm op} \cong \mathfrak{p}_2 \subset \mathfrak{g}_0^{ss}$. Then using Recipe 1,

$$\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_2) = \begin{cases} \frac{1}{2}\left(\dim(B_{\ell-1}) + 1 + \dim(A_1) + \dim(B_{\ell-3})\right), & n \text{ odd;} \\ \frac{1}{2}\left(\dim(D_{\ell-1}) + 1 + \dim(A_1) + \dim(D_{\ell-3})\right), & n \text{ even;} \end{cases}$$

$$= \begin{cases} 2\ell^2 - 7\ell + 10, & n \text{ odd;} \\ 2\ell^2 - 9\ell + 14, & n \text{ even} \end{cases} \quad \Rightarrow \quad \dim(\mathfrak{a}) = \dim(\mathfrak{g}_{-1}) + \dim(\mathfrak{a}_0) = \binom{n-1}{2} + 6.$$

(2) $n = 4$: D_3/P_1, $\lambda_{\mathfrak{g}} = \lambda_2 + \lambda_3$, $\mathfrak{g}_0^{ss} = A_1 \times A_1$. For $w = (12)$, $J_w = \{3\}$, $\mathfrak{p}_w^{\rm op} \cong A_1 \times \mathfrak{p}_1 \subset \mathfrak{g}_0^{ss}$, so $\dim(\mathfrak{a}_0) = 5$ and $\dim(\mathfrak{a}(w)) = 9$. By symmetry, for $w = (13)$, $\dim(\mathfrak{a}(w)) = 9$ as well.

Thus, $\mathfrak{U}_{\mathbb{C}} = \binom{n-1}{2} + 6$ for $n \geq 4$. In any signature over $\mathbb{R}$, $\mathfrak{S} \leq \mathfrak{U} \leq \mathfrak{U}_{\mathbb{C}}$ by Remark 4.2.7. For $n \geq 4$, $\mathfrak{a}_0$ is related to $\mathfrak{p}_2 \subset \mathfrak{g}_0^{ss}$, which is the Lie algebra of the stabilizer of a null 2-plane (in the standard representation of $\mathfrak{g}_0^{ss}$). These do not exist in Riemannian and Lorentzian signatures, so $\mathfrak{S} \leq \mathfrak{U} < \mathfrak{U}_{\mathbb{C}}$ in these cases. In all other signatures, we exhibit an explicit model realizing the upper bound.

5.1.1. *Non-Riemannian and non-Lorentzian signatures.* Consider the $(2,2)$ pp-wave metric:[12]

$$\mathrm{g}^{(2,2)}_{\mathbf{pp}} = y^2 dw^2 + dw dx + dy dz. \tag{5.1}$$

This has 9-dimensional conformal symmetry algebra spanned by

$$\begin{aligned} &\mathbf{X}_1 = \partial_x, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_w, \quad \mathbf{X}_4 = -y\partial_x + w\partial_z, \\ &\mathbf{X}_5 = 3(z + yw^2)\partial_x - 3w\partial_y - w^3\partial_z, \quad \mathbf{X}_6 = 2yw\partial_x - \partial_y - w^2\partial_z, \\ &\mathbf{X}_7 = -x\partial_x - y\partial_y + w\partial_w + z\partial_z, \quad \mathbf{X}_8 = 2y^3\partial_x - 3y\partial_w + 3x\partial_z, \\ &\mathbf{T} = 2x\partial_x + y\partial_y + z\partial_z. \end{aligned}$$

Note that $\mathbf{T}$ is a homothety, while $\mathbf{X}_1, \ldots, \mathbf{X}_8$ are Killing fields.

Lemma 5.1.1. *Let $n = p + q + 4$, and $\mathrm{g}^{(p,q)}_{\rm euc}$ the flat Euclidean metric of signature (p,q). Then $\mathrm{g} = \mathrm{g}^{(2,2)}_{\mathbf{pp}} + \mathrm{g}^{(p,q)}_{\rm euc}$ has conformal symmetry algebra of dimension $\binom{n-1}{2} + 6$.*

Proof. It is straightforward to check that $\mathrm{g}^{(2,2)}_{\mathbf{pp}}$ is not conformally flat, so neither is g. Writing $\epsilon_i = \pm 1$, the metric $\mathrm{g}^{(p,q)}_{\rm euc} = \sum_{i=1}^{p+q} \epsilon_i (du_i)^2$ admits the Killing fields $\mathbf{U}_i = \partial_{u_i}$ and $\mathbf{V}_{ij} = \epsilon_i u_i \partial_{u_j} - \epsilon_j u_j \partial_{u_i}$, where $i < j$. The metric g admits the Killing fields $\mathbf{U}_i$, $\mathbf{V}_{ij}$, $\mathbf{X}_k$, as well as $\mathbf{Y}_i = 2\epsilon_i u_i \partial_z - y\partial_{u_i}$, and $\mathbf{W}_i = 2\epsilon_i u_i \partial_x - w\partial_{u_i}$. The vector field $\tilde{\mathbf{T}} = \mathbf{T} + \sum_{i=1}^{p+q} u_i \partial_{u_i}$ is a homothety for g. □

Thus, we have proven:

Theorem 5.1.2. *For conformal geometry in dimension $n \geq 4$, we have $\mathfrak{S} \leq \binom{n-1}{2} + 6$. Except for Riemannian and Lorentzian signatures, this upper bound is sharp. For $n = 3$, we have $\mathfrak{S} = 4$.*

---

[12] In [30], the signature $(2,2)$ pp-wave metric was announced as having submaximal conformal symmetry dimension. No proof was given there, but using tools developed in this paper, we can confirm that this is indeed correct.

5.1.2. *4-dimensional Lorentzian.* Here, $\mathfrak{g} = \mathfrak{so}_{2,4}$ and $\mathfrak{g}_0 \cong \mathbb{R} \oplus \mathfrak{sl}_2(\mathbb{C})_\mathbb{R}$, where $\mathfrak{sl}_2(\mathbb{C})_\mathbb{R} \cong \mathfrak{so}(1,3)$ refers to the *real* Lie algebra underlying $\mathfrak{sl}_2(\mathbb{C})$. As before, $\mathbb{W} = H^2(\mathfrak{g}_-, \mathfrak{g})$ has homogeneity $+2$, which gives the action of the grading element $Z \in \mathfrak{z}(\mathfrak{g}_0) \cong \mathbb{R}$. As a $\mathfrak{sl}_2(\mathbb{C})_\mathbb{R}$-representation, $\mathbb{W} \cong \bigodot^4(\mathbb{C}^2)$, which is irreducible. Thus, Weyl tensors are identified with complex binary quartics, and their classification according to root type is precisely the well-known Petrov classification [35], [34].

To each Petrov type, there is a collection $\mathcal{O}$ of G_0-orbits. To calculate $\mathfrak{U}_\mathcal{O}$, it suffices to maximize $\mathfrak{a}_0^\phi = \mathfrak{ann}(\phi)$ among representative elements from these orbits. Then $\mathfrak{S}_\mathcal{O} \leq \mathfrak{U}_\mathcal{O}$ by Remark 4.2.9. Let $\{x, y\}$ be the standard $\mathbb{C}$-basis of $\mathbb{C}^2$, and let $H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ be the standard $\mathbb{C}$-basis for $\mathfrak{sl}_2(\mathbb{C})$, so $\mathfrak{sl}_2(\mathbb{C})_\mathbb{R}$ has $\mathbb{R}$-basis $\{H, iH, X, iX, Y, iY\}$. A simple case analysis yields Table 8 (except for the last column).

Type	Normal form ϕ	$\mathbb{R}$-basis for $\mathfrak{a}_0^\phi$	$\dim(\mathfrak{a}^\phi)$	$\mathfrak{a}^\phi$ filtration-rigid?
N	x^4	$X, iX, 2Z - H$	7	×
III	x^3y	$Z - H$	5	✓
D	x^2y^2	H, iH	6	×
II	$x^2y(x-y)$	$\cdot$	4	×
I	$xy(x-y)(x-ky)$	$\cdot$	4	×

Table 8. Petrov types and upper bounds on conformal symmetry dimensions

The following representative metrics and their conformal symmetries establish sharpness of the upper bounds (and filtration non-rigidity) in all cases except type III.

N: $\mathrm{g}^{(3,1)}_{\mathbf{pp}} = dy^2 + dz^2 + dwdx + y^2dw^2$ (signature $(3,1)$ pp-wave):

$$\mathbf{X}_1 = \partial_x, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_w, \quad \mathbf{X}_4 = -2z\partial_x + w\partial_z,$$
$$\mathbf{X}_5 = e^{-w}(2y\partial_x + \partial_y), \quad \mathbf{X}_6 = e^{w}(-2y\partial_x + \partial_y), \quad \mathbf{T} = 2x\partial_x + y\partial_y + z\partial_z.$$

III: $\mathrm{g} = \frac{3}{|\Lambda|}dz^2 + e^{4z}dx^2 + 4e^zdxdy + 2e^{-2z}(dy^2 + dudx)$ (Kaigorodov metric [38, (12.35)]):

$$\mathbf{X}_1 = \partial_u, \quad \mathbf{X}_2 = \partial_x, \quad \mathbf{X}_3 = \partial_y, \quad \mathbf{X}_4 = 2x\partial_x - y\partial_y - \partial_z - 4u\partial_u.$$

Another solution is $\mathrm{g} = \frac{r^2}{x^3}(dx^2 + dy^2) - 2dudr + \frac{3}{2}xdu^2$ (Siklos metric [38, (38.1)]):

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_u, \quad \mathbf{X}_3 = 2(x\partial_x + y\partial_y) + r\partial_r - u\partial_u, \quad \mathbf{T} = u\partial_u + r\partial_r.$$

D: $\mathrm{g} = a^2(dx^2 + \sinh^2(x)dy^2) + b^2(dz^2 - \sinh^2(z)dt^2)$ (a, b constant)

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_t, \quad \mathbf{X}_3 = e^{-t}(\partial_z + \coth(z)\partial_t), \quad \mathbf{X}_4 = e^{t}(\partial_z - \coth(z)\partial_t),$$
$$\mathbf{X}_5 = -\cos(y)\partial_x + \coth(x)\sin(y)\partial_y, \quad \mathbf{X}_6 = \sin(y)\partial_x + \coth(x)\cos(y)\partial_y.$$

This is a product of two spaces of constant curvature [38, (12.8)].

II: $\mathrm{g} = dz^2 + e^{-2z}(dy^2 + 2dxdu) - 35e^{4z}dx^2 + e^{-8z}du^2$

$$\mathbf{X}_1 = \partial_u, \quad \mathbf{X}_2 = \partial_x, \quad \mathbf{X}_3 = \partial_y, \quad \mathbf{X}_4 = 2x\partial_x - y\partial_y - \partial_z - 4u\partial_u.$$

This metric is not Einstein, has Ricci scalar a nonzero constant, and appears to be new. In [38], Table 38.3 incorrectly lists (12.29) as a type II metric with 4-dimensional isometry group, while (12.29) is in fact type D, as indicated at the bottom of p.179. Also, the type II metric (13.65) is indicated as having four Killing vectors (13.66), but the fourth listed vector field is incorrect. Professor Malcolm MacCallum has indicated to us that no type II metric with 4-dimensional isometry group appears to have been known in the literature.

Our strategy for finding such a type II metric was to obtain a deformation of the type III Kaigorodov metric while preserving its symmetries. This led to the ansatz

$$\mathrm{g} = dz^2 + a_1 e^{-2z} dy^2 + a_2 e^{-2z} dx du + a_3 e^{4z} dx^2 + a_4 e^{z} dy dx + a_5 e^{-8z} du^2,$$

where a_i are constants. Imposing the type II condition led to the metric indicated above.

I: $\mathrm{g} = dx^2 + e^{-2x}dy^2 + e^x \cos(\sqrt{3}x)(dz^2 - dt^2) - 2e^x \sin(\sqrt{3}x) dz dt$ (Petrov metric [38, (12.14)]):

$$\mathbf{X}_1 = \partial_y, \quad \mathbf{X}_2 = \partial_z, \quad \mathbf{X}_3 = \partial_t, \quad \mathbf{X}_4 = \partial_x + y\partial_y + \frac{1}{2}(\sqrt{3}t - z)\partial_z - \frac{1}{2}(t + \sqrt{3}z)\partial_t.$$

Theorem 5.1.3. *In 4-dimensional Lorentzian geometry, the maximal dimension of the conformal symmetry algebra for metrics of constant Petrov type is:*

Petrov type	*N*	*III*	*D*	*II*	*I*
max. sym. dim.	*7*	*4*	*6*	*4*	*4*

All models realizing these upper bounds are locally homogeneous near a regular point.

Proof. For type III metrics, we must show that the associated (5-dimensional) $\mathfrak{a} = \mathfrak{g}_{-1} \oplus \mathfrak{a}_0$ is filtration-rigid. As a $\mathfrak{sl}_2(\mathbb{C})_\mathbb{R}$-representation, $\mathfrak{g}_{-1}$ is the space of Hermitian 2×2-matrices $\mathcal{H}$ with the action of $A \in \mathfrak{sl}_2(\mathbb{C})_\mathbb{R}$ given by $M \mapsto AM + MA^*$. For $\mathcal{H}$, take the standard basis

$$e^1_{-1} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad e^2_{-1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad e^3_{-1} = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad e^4_{-1} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

Since H acts by $\mathrm{diag}(2, 0, 0, -2)$, and Z acts as -1 on $\mathfrak{g}_{-1}$, then $e_0 := Z - H$ acts diagonally with eigenvalues $\mathcal{E}_{-1} = \{-3, -1, -1, 1\}$, $\mathcal{E}_0 = \{0\}$. (All other brackets on $\mathfrak{a}$ are trivial.) Let $\mathfrak{f}$ be a filtered deformation of $\mathfrak{a}$. The first hypothesis of Proposition 4.4.5 is satisfied, as is (i), but (ii) is not. Following the argument there, we still have $\tilde{e}_0 \in \mathfrak{f}^0$ which acts diagonally in some basis $\{f^\alpha_i\}$ with the same eigenvalues as above, and with the only other non-trivial brackets $[f^2_{-1}, f^4_{-1}] = a\tilde{e}_0$, $[f^3_{-1}, f^4_{-1}] = b\tilde{e}_0$. By the Jacobi identity, $0 = \mathrm{Jac}_\mathfrak{f}(f^2_{-1}, f^3_{-1}, f^4_{-1}) = bf^2_{-1} - af^3_{-1}$, so $a = b = 0$. Thus, $\mathfrak{a}$ is filtration-rigid. In non-type III cases where the upper bound is realized, $\mathfrak{g}_- \subset \mathfrak{s}(u)$, so local homogeneity follows. In the type III case, $[\mathfrak{a}_0, \mathfrak{g}_{-1}] = \mathfrak{g}_{-1}$, and Remark 4.5.3 gives the result. ☐

5.1.3. *General Lorentzian and Riemannian.* For the Lorentzian case, we mimic the construction from Section 5.1.1. Consider $\mathrm{g} = \mathrm{g}^{(3,1)}_{\mathbf{pp}} + \mathrm{g}^{(p,0)}_{\mathrm{euc}}$, where $\mathrm{g}^{(3,1)}_{\mathbf{pp}}$ and its (seven) conformal symmetries were given in Section 5.1.2, and $\mathrm{g}^{(p,0)}_{\mathrm{euc}} = \sum_{i=1}^p (du^i)^2$ is the flat metric with Killing fields $\mathbf{U}_i = \partial_{u^i}$ and $\mathbf{V}_{ij} = u^i\partial_{u^j} - u^j\partial_{u^i}$. The metric g admits the Killing fields $\mathbf{X}_k$, $\mathbf{U}_i$, $\mathbf{V}_{ij}$, as well as $\mathbf{Y}_i = u^i\partial_z - z\partial_{u^i}$, and $\mathbf{W}_i = 2u^i\partial_x - w\partial_{u^i}$. In addition, there is the homothety $\widetilde{\mathbf{T}} = \mathbf{T} + \sum_{i=1}^p u_i\partial_{u_i}$, and so $\dim(\mathcal{S}) = \binom{n-1}{2} + 4$. Since the (complex) bound of $\binom{n-1}{2} + 6$ is not realizable, it remains to show that $\binom{n-1}{2} + 5$ is also not realizable. This is done in [16], from which it follows that $\mathfrak{S} = \binom{n-1}{2} + 4$.

Given a Riemannian metric g with nowhere vanishing Weyl tensor, there exists a conformally equivalent metric $\tilde{\mathrm{g}} = \lambda^2\mathrm{g}$ such that all conformal Killing fields of g are Killing fields for $\tilde{\mathrm{g}}$ [33]. Since the symmetry algebra is determined by restriction to any neighborhood, $\mathfrak{S}$ in the conformal Riemannian case is equal to the maximal dimension of the isometry algebra among metrics which are not conformally flat. According to Egorov's final result in [20] (incorporating results in his earlier article [19]), this number is exactly $\binom{n-1}{2} + 3$. This is realized by the product of spheres $\mathbb{S}^2 \times \mathbb{S}^{n-2}$ with their round metrics which is not conformally flat for $n \geq 4$, and has $\mathrm{SO}_3 \times \mathrm{SO}_{n-1}$ symmetry. However, we caution that there are omitted exceptions to Egorov's statement when $n = 4$ and $n = 6$.[13] In particular, with its Fubini–Study metric, $\mathbb{CP}^m$ is not conformally flat, has real dimension $n = 2m$, and has $\mathrm{SU}(m+1)$ symmetry group of dimension $m^2 + 2m$. This is strictly less than $\binom{n-1}{2} + 3$ for $n > 8$, equal to it when $n = 8$, and greater than it when $n = 4$ or 6.

---

[13] For $n = 4$, the submaximally symmetric structure is *unique*: $\mathbb{CP}^2$, cf. Egorov [18].

For an algebraic proof of $\mathfrak{S}$ in the conformal Lorentzian and Riemannian cases, we refer the reader to the article of B. Doubrov and D. The [16].

5.2. **Scalar 3rd order ODE.** Let $J^k = J^k(\mathbb{R}, \mathbb{R})$. Consider J^3 with its contact system $\mathcal{C}^3$. In canonical local coordinates (x, y, p, q, r), $\mathcal{C}^3$ is locally spanned by the 1-forms $\{dy - pdx, dp - qdx, dq - rdx\}$. A 3rd order ODE $y''' = F(x, y, y', y'')$ determines a hypersurface $M^4 = \{r = F(x, y, p, q)\} \subset J^3$ transverse to the projection $\pi^3_2 : J^3 \to J^2$. Pulling $\mathcal{C}^3$ back to M, we have two line fields (L, Π) on M: (i) L is spanned by $\frac{d}{dx} := \partial_x + p\partial_y + q\partial_p + F\partial_q$, and integral curves correspond to solutions of $y''' = F(x, y, y', y'')$; and (ii) Π is spanned by ∂_q. This is the fibre of $J^2 \to J^1$ (and hence $M \to J^1$), which is distinguished by Backlünd's theorem. The data (L, Π) encodes the geometry of scalar 3rd order ODE modulo contact transformations.

This is the underlying structure for a $G/P = C_2/P_{1,2}$ geometry, where $C_2 = \mathrm{Sp}(4, \mathbb{R})$. There are two well-known relative invariants due to Chern [13] and Sato–Yoshikawa [36]:

$$I_1 = -F_y - \frac{1}{3}F_pF_q - \frac{2}{27}{F_q}^3 + \frac{1}{2}\frac{d}{dx}(F_p) + \frac{1}{3}F_q\frac{d}{dx}(F_q) - \frac{1}{6}\left(\frac{d}{dx}\right)^2(F_q), \qquad I_2 = F_{qqqq}.$$

These respectively define the *conformal* branch ($I_1 = 0$) and *contact projective* branch ($I_2 = 0$). For more details, see also [24]. By Kostant, $H^2_+(\mathfrak{g}_-, \mathfrak{g})$ has two 1-dimensional components of homogeneity $+3$ and $+4$, and I_1 and I_2 correspond to $(\kappa_H)_{+3}$ and $(\kappa_H)_{+4}$ respectively. We have:

w	$\mathbb{V}_{-w\cdot\lambda_{\mathfrak{g}}}$	(Z_1, Z_2)	ϕ_0	$\dim(\mathfrak{a}(w))$	$\mathfrak{S}_w$	Twistor space type
(12)		$(+3, 0)$	$e_{\alpha_1} \wedge e_{2\alpha_1+\alpha_2} \otimes e_{-\alpha_2}$	5	5	C_2/P_1
(21)		$(+1, +3)$	$e_{\alpha_2} \wedge e_{\alpha_1+\alpha_2} \otimes e_{\alpha_2}$	5	4	C_2/P_2

The geometry is PR$_+$. If $\phi \in H^2_+(\mathfrak{g}_-, \mathfrak{g})$ has nonzero components in both submodules, then $\mathfrak{a}^\phi_0 = 0$, and $\dim(\mathfrak{a}^\phi) = \dim(\mathfrak{g}_-) = 4$. The $w = (21)$ branch is exceptional (3-dimensional conformal – see Section 4.5.3), and $\mathfrak{S}_w = \mathfrak{U}_w - 1 = \dim(\mathfrak{a}(w)) - 1 = 4$ here. For $w = (12)$, $\mathfrak{S}_w = \mathfrak{U}_w = \dim(\mathfrak{a}(w)) = 5$. Indeed, for any $a \in \mathbb{R}$, $y''' = ay' + y$ has five point (contact) symmetries:

$$\mathbf{X}_1 = \partial_x, \qquad \mathbf{X}_2 = y\partial_y, \qquad \mathbf{X}_i = \eta_i(x)\partial_y, \quad i = 3, 4, 5.$$

where $\eta_i(x)$ are the three linearly independent solutions of $y''' = ay' + y$. This proves:

Theorem 5.2.1. *For scalar 3rd order ODE modulo contact transformations, $\mathfrak{S} = 5$. Any submaximally symmetric model is in the contact projective branch ($I_2 = 0$), and is locally homogeneous near a non-flat regular point.*

Remark 5.2.2. The result $\mathfrak{S} = 5$ for 3rd order ODE is not explicitly stated in [43], but can be readily deduced as follows. Sophus Lie classified all finite-dimensional irreducible Lie algebras of *contact* vector fields. There are only three, L_6, L_7, L_{10}, with $\dim(L_i) = i$ with $L_6 \subset L_7$ and $L_6 \subset L_{10}$. The only 3rd order ODE invariant under L_6 (and hence L_7 and L_{10}) is the trivial equation $y''' = 0$. In [43], a classification of all *point* symmetry algebras for 3rd order ODE and representative equations is given. Aside from $y''' = 0$ and $y''' = \frac{3(y'')^2}{2y'}$ (which are *contact*-equivalent), having point symmetry algebras of dimensions 7 and 6, all others have dimension at most 5.

5.3. **Systems of 2nd order ODE.** Let $A_\ell = \mathrm{SL}(\ell + 1, \mathbb{R})$ and $\ell \geq 2$. If $\ell \neq 3$, the underlying structure for an $A_\ell/P_{1,2}$ geometry is a system of 2nd order ODE in $m = \ell - 1$ dependent variables,

$$\ddot{x}^i = f^i(t, x^j, \dot{x}^j), \qquad 1 \leq i \leq m, \tag{5.2}$$

with equivalence up to *point transformations.* (This is the same for $\ell = 3$ provided that $(\kappa_H)_{+1} \equiv 0$.) This is *path geometry.* The $m = 1$ case was discussed in Section 4.5.2, so here we discuss the $m \geq 2$ case, which was studied by Fels [22] and Grossman [25]. Let $J^k = J^k(\mathbb{R}, \mathbb{R}^m)$ with its contact system $\mathcal{C}^k$. Geometrically, (5.2) defines a submanifold $M \subset J^2$ transverse to $\pi^2_1 : J^2 \to J^1$, and point transformations refer to diffeomorphisms of J^2 which preserve $\mathcal{C}^2$ which are prolongations of arbitrary diffeomorphisms of J^0. On J^2, in canonical local coordinates (t, x^j, p^j, q^j), $\mathcal{C}^2$ is locally spanned by the 1-forms $\{dx^j - p^j dt, dp^j - q^j dt\}$ and $M = \{q^i = f^i(t, x^j, p^j)\}$ fibers of J^0. This is an "external" description of the geometry. Pulling back $\mathcal{C}^2$ to M yields an "internal" description: a manifold of dimension $n = 2m + 1$, with coordinates (t, x^i, p^i), and a distribution containing:

(1) a distinguished line field L spanned by $\frac{d}{dt} := \partial_t + p^i\partial_{x^i} + f^i\partial_{p^i}$ whose integral curves correspond to solutions of (5.2), and
(2) an integrable m-dimensional distribution Π spanned by $\{\partial_{p^i}\}_{i=1}^m$. This is the fibre of the submersion $M \to J^0$, which is preserved by all point transformations.

This flat model is $\ddot{x}^i = 0$, $1 \le i \le m$, which has $\dim(A_\ell) = \ell^2 + 2\ell$ dimensional symmetry algebra.

In [22], Fels studied the geometry of systems of 2nd order ODE using Cartan's method of equivalence and derived the following invariants. Given (5.2), define

$$G^i_{jkl} = \frac{\partial^3 f^i}{\partial p^j \partial p^k \partial p^l}, \qquad F^i_j = \frac{1}{2}\frac{d}{dt}\left(\frac{\partial f^i}{\partial p^j}\right) - \frac{\partial f^i}{\partial x^j} - \frac{1}{4}\frac{\partial f^i}{\partial p^k}\frac{\partial f^k}{\partial p^j}.$$

The *Fels invariants*, namely the Fels curvature and torsion, are respectively:

$$S^i_{jkl} = G^i_{jkl} - \frac{3}{m+2} G^r_{r(jk}\delta^i_{l)}, \qquad T^i_j = F^i_j - \frac{1}{m}\delta^i_j F^k_k. \tag{5.3}$$

The Tanaka prolongation of the symbol algebra $\mathfrak{g}_-$ and structure group G_0 is, by Yamaguchi, $\mathfrak{g} \cong \mathfrak{sl}(\ell + 1, \mathbb{R})$ with 2-grading given by

$$\mathfrak{g}_0 = \mathbb{R}^2 \oplus \mathfrak{sl}(\Pi_{-1}), \qquad \mathfrak{g}_{\pm 1} = L_{\pm 1} \oplus \Pi_{\pm 1}, \qquad \mathfrak{g}_{\pm 2} = L_{\pm 1} \otimes \Pi_{\pm 1}.$$

For $\ell \ge 4$, $W^{\mathfrak{p}}_+(2) = \{(21), (12)\}$, which generate homogeneity +3 and +2 modules respectively:

(5.4) $H^2_+(\mathfrak{g}_-, \mathfrak{g}) =$ 

$\oplus$ 

$$= L_1 \otimes \left(\bigodot{}^3 \Pi_1 \otimes \Pi_{-1}\right)_0 \oplus L_1^{\otimes 2} \otimes (\Pi_1 \otimes \Pi_{-1})_0 ,$$

where $(\Pi_{-1})^* \cong \Pi_1$ via the Killing form, so the "0" subscripts indicate trace-free with respect to contractions. For the tensor descriptions above, we used

$L_{-1} =$ 

, $L_1 =$ 

,

$\Pi_{-1} =$ 

, $\Pi_1 =$ 

.

(e.g. The lowest roots of Π_1 and Π_{-1} are α_2 and $-\alpha_2 - ... - \alpha_\ell$, so by the "minus lowest weight convention", we convert their negatives into weight notation using the Cartan matrix – see (2.4).)

For $\ell = 3$, there is additionally the 1-dimensional module 

$= \bigwedge^2 \Pi_1 \otimes L_{-1}$, which has homogeneity +1 and corresponds to the word $w = (23) \in W^{\mathfrak{p}}_+(2)$. For ODE, we must have $(\kappa_H)_{+1} \equiv 0$. We discuss $A_3/P_{1,2}$ geometries with $(\kappa_H)_{+1} \ne 0$ in Section 5.5.2.

The tensors S and T correspond to $(\kappa_H)_{+3}$ and $(\kappa_H)_{+2}$ respectively. By the discussion on correspondence and twistor spaces (Section 2.7), we have

- $S \equiv 0$ iff $(\kappa_H)_{+3} \equiv 0$ iff the geometry admits an A_ℓ/P_1 description, i.e. (5.2) are the equations for (unparametrized) geodesics of projective connection, cf. Section 5.4.
- $T \equiv 0$ iff $(\kappa_H)_{+2} \equiv 0$ iff the geometry admits an $A_\ell/P_2 \cong \mathrm{Gr}(2, \mathbb{R}^{\ell+1})$ description. This is a 1-graded geometry with underlying structure a manifold N^{2m}, endowed with a field $\mathcal{S} \subset \mathbb{P}(TN)$ of type $(2, m)$-Segré varieties. This is a *Segré* (or *almost Grassmannian*) *structure*.[14]

---

[14] When $m = 2$, this is equivalent to a signature $(2, 2)$ conformal structure. Note $A_3/P_2 \cong D_3/P_1$.

In [25], Grossman studies (5.2) satisfying $T \equiv 0$, which he calls *torsion-free* path geometries. If such a geometry is moreover "geodesic" (i.e. $S \equiv 0$), then $\kappa_H \equiv 0$, and so the geometry is locally flat, cf. [25, Theorem 1]. With these preparations, we state our new result:

Theorem 5.3.1. *For systems of 2nd order ODE in m dependent variables,* $\mathfrak{S} = \begin{cases} m^2+5, & m \geq 2; \\ 3, & m = 1. \end{cases}$ *When $m \geq 2$, any submaximally symmetric model satisfies $T \equiv 0$ (so is not geodesic).*

Proof. Let $m = \ell - 1 \geq 2$. Look at $A_\ell/P_{1,2}$. (For $m = 2$, we require $(\kappa_H)_{+1} \equiv 0$.) For $w = (12)$, $\mathfrak{S}_w = m^2 + 4$; see Section 5.4. For $w = (21)$, we have by (5.4), $J_w = \{3, \ell\}$ and $I_w = \{1\}$, so the geometry is NPR$_+$. By Recipe 6, the reduced geometry is $\bar{\mathfrak{g}}/\bar{\mathfrak{p}} \cong A_1/P_1$, so $\dim(\mathfrak{a}_1) = 1$ and $\dim(\mathfrak{a}_2) = 0$. From $J_w = \{3, \ell\}$ and Recipe 5, $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_{1,\ell-2} \subset A_{\ell-2}$, and $\dim(\mathfrak{a}_0) = 1 + \dim(\mathfrak{p}_w^{\mathrm{op}}) = m^2 - 2m + 3$, by Recipe 1. Hence, $\mathfrak{S}_w = \mathfrak{U}_w = \dim(\mathfrak{a}(w)) = \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0) + \dim(\mathfrak{a}_1) = 2m + 1 + (m^2 - 2m + 3) + 1 = m^2 + 5$. ☐

For any $m \geq 2$, the following is a submaximally symmetric model with its (point) symmetries:

$$\text{(5.5)} \qquad \left\{ \begin{array}{ll} \ddot{x}^1 & = 0, \\ & \vdots \\ \ddot{x}^{m-1} & = 0, \\ \ddot{x}^m & = (\dot{x}^1)^3. \end{array} \right. \quad \left| \begin{array}{l} \mathbf{T} = \partial_t, \quad \mathbf{X}_i = \partial_{x^i}, \quad \mathbf{A}_j = t\partial_{x^j}, \quad \mathbf{B}_j^k = x^k \partial_{x^j} \\ \qquad (1 \leq i \leq m, \quad 2 \leq j \leq m, \quad 1 \leq k < m) \\ \mathbf{C} = 2t\partial_{x^1} + 3(x^1)^2 \partial_{x^m}, \quad \mathbf{D}_1 = x^1 \partial_{x^1} + 3x^m \partial_{x^m}, \\ \mathbf{D}_2 = t\partial_t - x^m \partial_{x^m}, \quad \mathbf{S} = t^2 \partial_t + t \sum_{i=1}^m x^i \partial_{x^i} + \frac{1}{2}(x^1)^3 \partial_{x^m} \end{array} \right.$$

The symmetries are stated as vector fields $\mathbf{V} = \xi\partial_t + \varphi^i \partial_{x^i}$ on J^0, but these admit unique prolongations $\mathbf{V}^{(1)} = \mathbf{V} + \varphi^i_{(1)} \partial_{p^i}$ on J^1 and $\mathbf{V}^{(2)} = \mathbf{V}^{(1)} + \varphi^i_{(2)} \partial_{q^i}$ on J^2 via the formula

$$\text{(5.6)} \qquad \varphi^i_{(1)} = \frac{d\varphi^i}{dt} - p^i \frac{d\xi}{dt}, \qquad \varphi^i_{(2)} = \frac{d\varphi^i_{(1)}}{dt} - q^i \frac{d\xi}{dt}.$$

Externally, $\mathbf{V}^{(2)}$ are everywhere tangent to $M \subset J^2$. Internally, $\mathbf{V}^{(1)}$ are identified with vector fields on M which preserve the geometric data described earlier.

Remark 5.3.2. The model (5.5) was found in the $m = 2$ case by Casey et al. [14, (3.5)]. For the projective branch ($S \equiv 0$), they used Egorov's result [17] (see also our Section 5.4 below) to assert 8 as the submaximal symmetry dimension. For the conformal (Segré) branch ($T \equiv 0$), their model "*corresponds to a Ricci-flat ASD conformal structure with only one constant non-vanishing component of the ASD Weyl tensor*". They conclude submaximality in this branch and hence for pairs of ODE. While their conclusion is correct, we note that:

1. The position of the single non-vanishing component is relevant in assessing submaximal symmetry: it must correspond to a lowest (or highest) weight vector. However, this argument fails in Riemannian and Lorentzian signatures: lowest weight vectors do not exist, and the most symmetric (degenerate) Weyl tensors have many non-vanishing components [16].
2. The case where both $S \neq 0$ and $T \neq 0$ is not considered. While one expects any submaximal model to satisfy either $S \equiv 0$ or $T \equiv 0$, this needs to be proven. (See our Theorem 4.3.8.)

Since $T \equiv 0$, let us exhibit the $(2, m)$-Segré structure. The twistor space N is the quotient of M by the integral curves of $\frac{d}{dt}$, i.e. it is the solution space of the ODE. The solution of (5.5) is

$$\text{(5.7)} \qquad \begin{array}{rlll} & x^1 = a_1 t + b_1, \ \ldots, & x^{m-1} = a_{m-1} t + b_{m-1}, & x^m = \frac{1}{2}(a_1)^3 t^2 + a_m t + b_m \\ \Rightarrow & p^1 = a_1, \qquad \ \ldots, & p^{m-1} = a_{m-1}, & p^m = (a_1)^3 t + a_m, \end{array}$$

where $z = (a_1, \ldots, a_m, b_1, \ldots, b_m)$ are parameters (local coordinates on N). The map $\Psi : M \to N$ is

$$\left\{ \begin{array}{llll} a_1 = p^1, & \ldots, & a_{m-1} = p^{m-1}, & a_m = p^m - (p^1)^3 t, \\ b_1 = x^1 - p^1 t, & \ldots, & b_{m-1} = x^{m-1} - p^{m-1} t, & b_m = x^m + \frac{1}{2}(p^1)^3 t^2 - p^m t. \end{array} \right.$$

Fixing $z \in N$, $\Psi^{-1}(z)$ is a solution curve c in M given by (5.7), which induces a curve $\zeta(t) = \Psi_*(\Pi|_{c(t)})$ in $\mathrm{Gr}(m, T_z N)$. Indeed, $\zeta(t)$ is the span of $\mathbf{Z}_i(t) = \Psi_*(\partial_{p^i}|_{c(t)}) \in T_z N$, where

$$\begin{aligned} \mathbf{Z}_1(t) &= \partial_{a_1} - 3(a_1)^2 t \partial_{a_m} - t\partial_{b_1} + \tfrac{3}{2}(a_1)^2 t^2 \partial_{b_m}, \\ \mathbf{Z}_2(t) &= \partial_{a_2} - t\partial_{b_2}, \quad \dots, \quad \mathbf{Z}_m(t) = \partial_{a_m} - t\partial_{b_m}. \end{aligned} \tag{5.8}$$

Define a $2 \times m$ matrix of 1-forms $\Theta = \{\Theta_i^A\}$ on N given by

$$\Theta_i^1 = \left\{ \begin{array}{ll} da_i, & i \neq m; \\ da_m - \frac{3}{2}(a_1)^2 db_1, & i = m \end{array} \right., \qquad \Theta_i^2 = db_i, \qquad 1 \leq i \leq m. \tag{5.9}$$

The 2×2 minors of Θ give the Segré equations cutting out a Segré variety $\mathcal{S}|_z \subset \mathbb{P}(T_z N)$,

$$\Theta_i^A \Theta_j^B - \Theta_j^A \Theta_i^B = 0, \quad 1 \leq i, j \leq m, \quad 1 \leq A, B \leq 2. \tag{5.10}$$

These quadratic equations characterize the image of the Segré embedding $\sigma : \mathbb{P}^1 \times \mathbb{P}^{m-1} \hookrightarrow \mathbb{P}^{2m-1} \cong \mathbb{P}(T_z N)$, $[u] \times [v] \mapsto [u \otimes v]$. The curve $\zeta(t)$ given by (5.8) satisfies (5.10) and so $\zeta(t)$ is contained in and moreover fills out $\mathcal{S}|_z$ as t varies (regarding t as a projective parameter).

Symmetries of the Segré structure corresponding to (5.9) are easily calculated: given any symmetry $\mathbf{V}$ of the ODE (5.5), compute $\widetilde{\mathbf{V}} = \Psi_*(\mathbf{V}^{(1)})$. This yields

$$\widetilde{\mathbf{T}} = -(a_1)^3 \partial_{a_m} - \sum_{i=1}^m a_i \partial_{b_i}, \qquad \widetilde{\mathbf{X}}_i = \partial_{b_i}, \qquad \widetilde{\mathbf{A}}_j = \partial_{a_j}, \qquad \widetilde{\mathbf{B}}_j^k = a_k \partial_{a_j} + b_k \partial_{b_j},$$

$$(1 \leq i \leq m, \quad 2 \leq j \leq m, \quad 1 \leq k < m)$$

$$\widetilde{\mathbf{C}} = 2\partial_{a_1} + 6a_1 b_1 \partial_{a_m} + 3(b_1)^2 \partial_{b_m}, \qquad \widetilde{\mathbf{D}}_1 = a_1 \partial_{a_1} + 3a_m \partial_{a_m} + b_1 \partial_{b_1} + 3b_m \partial_{b_m},$$

$$\widetilde{\mathbf{D}}_2 = -\sum_{k=1}^{m-1} a_k \partial_{a_k} - 2a_m \partial_{a_m} - b_m \partial_{b_m}, \qquad \widetilde{\mathbf{S}} = \sum_{i=1}^m b_i \partial_{a_i} + \frac{3}{2} a_1 (b_1)^2 \partial_{a_m} + \frac{1}{2}(b_1)^3 \partial_{b_m}.$$

Direct calculation verifies that the ideal generated by the left hand side of (5.10) is preserved under $\mathcal{L}_{\widetilde{\mathbf{V}}}$, so these are indeed symmetries of the Segré structure.

When $m = 2$, (5.10) is a single quadratic equation $\mathrm{g} := \Theta_1^1 \Theta_2^2 - \Theta_2^1 \Theta_1^2 = 0$, so a Segré structure is the same (up to a sign) as a split signature conformal structure $[\mathrm{g}]$ in dimension four (and $\Psi_*(\Pi)$ are the null planes). Changing coordinates, g is simply the pp-wave given in (5.1).

Any Segré variety contains two distinguished *rulings*, i.e. maximal (projective) linear spaces. With respect to the Segré embedding σ, writing $\sigma_u(v) = \sigma(u, v) = \sigma^v(u)$, we have for (5.9):

- m-ruling: $\sigma_u : \mathbb{P}^{m-1} \to \mathbb{P}^{2m-1}$ for $[u] \in \mathbb{P}^1$. Each m-plane in $\mathcal{R}|_z \subset \mathrm{Gr}(m, T_z N)$ has basis
$$u_1 \partial_{a_1} + u_2 \left(\partial_{b_1} + \frac{3}{2}(a_1)^2 \partial_{a_m} \right), \quad u_1 \partial_{a_2} + u_2 \partial_{b_2}, \quad \dots, \quad u_1 \partial_{a_m} + u_2 \partial_{b_m} \quad \in T_z N.$$
- 2-ruling: $\sigma^v : \mathbb{P}^1 \to \mathbb{P}^{2m-1}$ for $[v] \in \mathbb{P}^{m-1}$. Each 2-plane in $\mathcal{P}|_z \subset \mathrm{Gr}(2, T_z N)$ has basis
$$v_1 \partial_{a_1} + \dots + v_m \partial_{a_m}, \quad v_1 \left(\frac{3}{2}(a_1)^2 \partial_{a_m} + \partial_{b_1} \right) + v_2 \partial_{b_2} \dots + v_m \partial_{b_m} \quad \in T_z N.$$

This gives submanifolds $\mathcal{R} \subset \mathrm{Gr}(m, TN)$ and $\mathcal{P} \subset \mathrm{Gr}(2, TN)$. From (5.8), we see that $\Psi_*(\Pi) \subset \mathcal{R}$. The image under Ψ of each fibre of $M \to J^0$ is an m-dimensional submanifold $\Sigma \subset N$, with $T_z \Sigma \in \mathcal{R}|_z \subset T_z N$ for any $z \in \Sigma$. Conversely, every element of $\mathcal{R}$ is tangent to such a submanifold, so $\mathcal{R}$ is *integrable.* In fact, Grossman [25] shows that integrability of the m-ruling $\mathcal{R}$ is a general feature of Segré structures arising from 2nd order ODE systems by establishing an isomorphism of the bundles $M \to N$ and the $\mathbb{P}^1$-bundle $P_\Delta \to N$ whose fibres parametrize the m-ruling. The 2-ruling $\mathcal{P}$ is never integrable for such structures, except for the flat model.

The $T \equiv 0$ branch for ODE and associated Segré structures have κ_H concentrated in the respective modules corresponding to $w = (21)$. For Segré structures, there is an additional (23)-branch. For structures in this branch, $\mathcal{R}$ is non-integrable while $\mathcal{P}$ is integrable.

Theorem 5.3.3. *Let $m \geq 2$. For $(2,m)$-Segré structures,*

$$\mathfrak{S} = \mathfrak{S}_{(21)} = m^2 + 5, \qquad \mathfrak{S}_{(23)} = \begin{cases} m^2 - m + 8, & m \geq 3; \\ 9, & m = 2. \end{cases}$$

Proof. Consider A_ℓ/P_2 with $\ell = m+1$. We have established $\mathfrak{S}_{(21)} = m^2 + 5$. If $m = 2$, then $A_3/P_2 \cong D_3/P_1$, which we have already considered. Letting $m \geq 3$ and $w = (23)$, $J_w = \{1, 4, \ell\}$ and $I_w = \emptyset$, so PR. By Recipes 5 and 1, $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_1 \times \mathfrak{p}_{2,\ell-2} \subset A_1 \times A_{\ell-2}$, and $\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_w^{\mathrm{op}}) = m^2 - 3m + 8$. Hence, $\mathfrak{S}_w = \mathfrak{U}_w = \dim(\mathfrak{a}(w)) = \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0) = m^2 - m + 8$. ☐

Thus, the Segré structure (5.9) in the (21)-branch is indeed submaximally symmetric. When $m = 2$ or 3 there is also a submaximally symmetric model in the (23)-branch.

5.4. **Projective structures.** Two torsion-free affine connections ∇, ∇' are equivalent if their *unparametrized* geodesics are the same, and a *projective connection* is such an equivalence class $[\nabla]$. These are the underlying structures for a (regular, normal) A_ℓ/P_1 geometry, or equivalently (see Section 5.3) an $A_\ell/P_{1,2}$ geometry with vanishing Fels curvature ($S \equiv 0$).

Locally, take coordinates $(x^a)_{a=0}^{\ell-1}$ on an ℓ-manifold N and define ∇ via its Christoffel symbols

$$\nabla_{\partial_{x^a}} \partial_{x^b} = \Gamma^c_{ab} \partial_{x^c}, \qquad \Gamma^c_{ab} = \Gamma^c_{(ab)}, \qquad 0 \leq a, b, c \leq \ell - 1 =: m.$$

The geodesic equation $\nabla_{\sigma'}\sigma' = 0$ for a curve $\sigma = \sigma(t)$ takes the local form $(x^a)'' + \Gamma^a_{bc}(x^b)'(x^c)' = 0$. Locally, this always has solutions, so relabelling coordinates if necessary we may assume $(x^0)' \neq 0$, and we can solve for $t = t(x^0)$. We eliminate dependence on the parameter t by regarding x^i, $i > 0$, as functions of x^0. By the chain rule, the geodesic equations become

$$\begin{aligned} \ddot{x}^i &= \Gamma^0_{ab}\dot{x}^i\dot{x}^a\dot{x}^b - \Gamma^i_{ab}\dot{x}^a\dot{x}^b, \\ &= \Gamma^0_{jk}\dot{x}^i\dot{x}^j\dot{x}^k + 2\Gamma^0_{0j}\dot{x}^i\dot{x}^j - \Gamma^i_{jk}\dot{x}^j\dot{x}^k + \Gamma^0_{00}\dot{x}^i - 2\Gamma^i_{0j}\dot{x}^j - \Gamma^i_{00}, \qquad 1 \leq i, j, k \leq m, \end{aligned} \tag{5.11}$$

and we consider this system up to point transformations. Internally, the ODE structure is defined on a $(2m+1)$-manifold M with local coordinates (t, x^j, p^j). Projective equivalence $\nabla \mapsto \nabla'$ is equivalently given operationally by $\Gamma^a_{bc} \mapsto \Gamma^a_{bc} + \delta^a_b \Upsilon_c + \delta^a_c \Upsilon_b$, where Υ is an arbitrary 1-form. As expected, (5.11) is invariant under these changes.

Egorov [17] studied the gap problem for projective structures by analyzing the integrability conditions for symmetries. Let us reprove his results using our Lie algebraic methods. Observe that the model A_ℓ/P_1 gives a 1-grading $\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1$, where $\dim(\mathfrak{g}_{-1}) = \ell$, and $\mathfrak{g}_0 \cong \mathbb{C} \times A_m$. By Kostant, we have $W^{\mathfrak{p}}_+(2) = \{(12)\}$, and

$$H^2_+(\mathfrak{g}_-, \mathfrak{g}) =$$



$$= \left(\bigwedge^2 \mathfrak{g}_1 \otimes \mathfrak{sl}(\mathfrak{g}_{-1}) \right)_0, \qquad \ell \geq 3.$$

Theorem 5.4.1. *Let $\ell \geq 2$. For projective structures on ℓ-manifolds,* $\mathfrak{S} = \begin{cases} (\ell-1)^2 + 4, & \ell \geq 3; \\ 3, & \ell = 2. \end{cases}$

Proof. The $\ell = 2$ case is exceptional (see Section 4.5), so let $\ell \geq 3$. For $w = (12)$, $J_w = \{2, 3, \ell\}$, $I_w = \emptyset$, and $\mathfrak{a}(w) = \mathfrak{g}_- \oplus \mathfrak{a}_0$. Then $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_{1,2,\ell-1} \subset A_{\ell-1}$, and $\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_w^{\mathrm{op}}) = \ell^2 - 3\ell + 5$. Then $\mathfrak{S} = \mathfrak{U}_w = \dim(\mathfrak{a}(w)) = \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0) = \ell^2 - 2\ell + 5 = (\ell - 1)^2 + 4$. ☐

Egorov [17] also gave the following submaximally symmetric model for $m = \ell - 1 \geq 2$:

$$\Gamma^0_{12} = \Gamma^0_{21} = x^1, \qquad \Gamma^c_{ab} = 0 \text{ otherwise.} \tag{5.12}$$

The Weyl curvature of this connection has components

$$W_{121}{}^0 = -W_{211}{}^0 = 1, \qquad W_{abc}{}^d = 0 \text{ otherwise.}$$

The associated 2nd order ODE system is

$$\ddot{x}^i = 2x^1\dot{x}^1\dot{x}^2\dot{x}^i \qquad (1 \le i \le m). \tag{5.13}$$

Its m^2+4 point symmetries are (writing $t := x^0$):

$$\begin{cases} \partial_t, \quad \partial_{x^2}, \ \ldots, \ \partial_{x^m}, \quad x^i\partial_t, \quad x^i\partial_{x^j} \quad (i \ge 1, \ j \ge 3), \\ 2t\partial_t + x^1\partial_{x^1}, \quad t\partial_t + x^2\partial_{x^2}, \quad x^1x^2\partial_t - \partial_{x^1}, \quad (x^1)^3\partial_t - 3x^1\partial_{x^2}. \end{cases} \tag{5.14}$$

where, as in Section 5.3, the actual symmetries on (t, x^j, p^j)-space M are the prolongation of these vector fields. Since $\{\partial_{p^i}\}_{i=1}^m$ spans the vertical subspace for the projection $M \to N$, then (5.14) are also the symmetries of the projective connection determined by (5.12). The Fels torsion (5.3) $T = (T^i_j)$ of (5.13) has matrix rank 1 and components

$$T^i_1 = -p^1p^2p^i, \qquad T^i_2 = (p^1)^2p^i, \qquad T^i_j = 0, \quad \text{for} \quad j \ge 3.$$

5.5. **Bracket-generating distributions.**

5.5.1. G_2/P_1 *geometry.* We work over $\mathbb{C}$. (Over $\mathbb{R}$, use the split-real form of G_2.) The underlying geometry for a (regular, normal) G_2/P_1 geometry is a $(2,3,5)$-distribution. The equivalence problem for such structures was studied by Cartan in his well-known five-variables paper [11]. The flat model, in Monge form (see Example 2.1.1), is the Hilbert–Cartan equation $z' = (y'')^2$.

Writing $\mathfrak{g} = \text{Lie}(G_2)$, $\mathfrak{p}_1$ induces a 3-grading on $\mathfrak{g}$, with $\mathfrak{g}_0 \cong \mathfrak{gl}_2(\mathbb{C})$. In Example 2.5.2, we saw $H^2_+(\mathfrak{g}_-, \mathfrak{g}) \cong \overset{-8}{\times}\!\Lleftarrow\!\overset{4}{*} \cong \bigodot^4(\mathfrak{g}_{-1})^* \cong \bigodot^4(\mathbb{C}^2)^*$, i.e. binary quartics.

Theorem 5.5.1. *For* $(2,3,5)$*-distributions,* $\mathfrak{S} = 7$.

Proof. For $w = (12)$, $J_w = \{2\}$, $I_w = \emptyset$, so G_2/P_1 is PR$_+$, and $\mathfrak{a}(w) = \mathfrak{g}_- \oplus \mathfrak{a}_0$. Hence, $\mathfrak{p}_w^{\text{op}} \cong \mathfrak{p}_1 \subset A_1$, so $\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_w^{\text{op}}) = 2$. Thus, $\mathfrak{S} = \dim(\mathfrak{a}(w)) = \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0) = 5 + 2 = 7$. ☐

Using the classification of $G_0 \cong \text{GL}_2(\mathbb{C})$ orbits in $\bigodot^4(\mathbb{C}^2)^*$, we analyze each root type (similar to the Petrov analysis in Section 5.1.2). Let $\{e_1, e_2\}$ and $\{\omega_1, \omega_2\}$ be (Killing) dual bases in $\mathfrak{g}_{-1}$ and $\mathfrak{g}_1$ respectively. Any element of $\mathfrak{g}_0 = \mathfrak{gl}(\mathfrak{g}_{-1})$ acts on $\bigodot^4(\mathfrak{g}_{-1})^*$, e.g. $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \omega_1^4 = -4(a\omega_1 + b\omega_2)\omega_1^3$. For a $\mathfrak{gl}_2(\mathbb{C})$ basis, take $X = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $Y = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. (Note $Z = -I$.) Symmetry bounds for each root type are given in Table 9. Some models, found either by Cartan [11] or Strazzullo [39, Section 6.10], are given in Table 10. We refer to [39] for the symmetry algebras.

Type	Normal form ϕ	Basis for $\mathfrak{a}_0^\phi$	$\dim(\mathfrak{a}^\phi)$	$\mathfrak{a}^\phi$ filtration-rigid?
(4)	ω_1^4	$X, H-I$	7	×
$(3,1)$	$\omega_1^3\omega_2$	$2H-I$	6	✓
$(2,2)$	$\omega_1^2\omega_2^2$	H	6	×
$(2,1,1)$	$\omega_1^2\omega_2(\omega_1-\omega_2)$	$\cdot$	5	×
$(1,1,1,1)$	$\omega_1\omega_2(\omega_1-\omega_2)(\omega_1-k\omega_2)$	$\cdot$	5	×

TABLE 9. Root types and upper bounds on symmetry dimensions for G_2/P_1 geometries

Theorem 5.5.2. *Among* $(2,3,5)$*-distributions with constant root type:*

Root type	(4)	$(3,1)$	$(2,2)$	$(2,1,1)$	$(1,1,1,1)$
max. $\dim(\mathcal{S})$	7	4 *or* 5	6	5	5

Root type	$F(x,y,p,q,z)$	$\dim(\mathcal{S})$	Reference in [39]
(4)	q^3	7	6.1.1
$(3,1)$	p^3+q^2	4	6.1.3
$(2,2)$	$y+\ln(q)$	6	6.3.4
$(2,1,1)$	pq^2	5	6.1.6
$(1,1,1,1)$	pq^3	5	6.1.7

Table 10. Some model $(2,3,5)$-distributions for each root type

Except for type $(3,1)$*, all models realizing these upper bounds are locally homogeneous near a regular point.*[15]

Proof. We show that in type $(3,1)$, $\mathfrak{a}=\mathfrak{g}_-\oplus\mathfrak{a}_0$ is filtration-rigid. Here, $e_0:=2H-I$ spans $\mathfrak{a}_0$. Take a basis of $\mathfrak{g}_-$ as in (2.1). Then ad_{e_0} is diagonal with

$$\mathcal{E}_0=\{0\},\quad \mathcal{E}_{-1}=\{1,-3\},\quad \mathcal{E}_{-2}=\{-2\},\quad \mathcal{E}_{-3}=\{-1,-5\}.$$

Let $\mathfrak{f}$ be a filtered deformation of $\mathfrak{a}$. The first hypothesis of Proposition 4.4.5 is satisfied, as is (i), but (ii) is not. Similar to the Petrov type III case, there is $\tilde{e}_0\in\mathfrak{f}^0$ which acts diagonally in some basis $\{f_i^\alpha\}$ with the same eigenvalues as above, and with the only other non-trivial brackets

$$[f_{-1}^1,f_{-1}^2]=f_{-2},\quad [f_{-1}^1,f_{-2}]=f_{-3}^1,\quad [f_{-1}^2,f_{-2}]=f_{-3}^2,\quad [f_{-1}^1,f_{-3}^1]=a\tilde{e}_0,\quad [f_{-2},f_{-3}^1]=bf_{-1}^2.$$

Since $0=\mathrm{Jac}_{\mathfrak{f}}(f_{-1}^1,f_{-3}^1,f_{-3}^2)=-5af_{-3}^2$ and $0=\mathrm{Jac}_{\mathfrak{f}}(f_{-1}^1,f_{-2},f_{-3}^1)=(2a-b)f_{-2}$, then $\mathfrak{a}$ is filtration-rigid. Aside from type $(3,1)$, when the upper bound is realized, $\mathfrak{g}_-\subset\mathfrak{s}(u)$. □

Theorem 5.5.2 recovers the bounds found by Cartan [11] using his method of equivalence.

5.5.2. *$A_3/P_{1,2}$ geometries.* As in Section 5.3, the underlying structure for regular, normal $A_3/P_{1,2}$ geometries is still a distribution D on a 5-manifold with a splitting $D=L\oplus\Pi$ into rank 1 and 2 subbundles, but now Π is not necessarily integrable; instead $[\Pi,\Pi]\subseteq D$ [10, 6]. Having $(\kappa_H)_{+1}$ nowhere vanishing is equivalent to $[\Pi,\Pi]=D$, in which case Π becomes a $(2,3,5)$-distribution.

Example 5.5.3. *On a 5-manifold* (x,y,p,q,z)*, consider* Π *spanned by* $\{\partial_q,\partial_x+p\partial_y+q\partial_p+q^2\partial_z\}$ *(i.e. the flat model for* $(2,3,5)$*-distributions) and* $L=\{\partial_p+2q\partial_z\}$*. Then* (Π,L) *has* $\mathcal{S}$ *given by*

$$\mathbf{X}_1=\partial_x,\quad \mathbf{X}_2=\partial_y,\quad \mathbf{X}_3=\partial_z,\quad \mathbf{X}_4=x\partial_y+\partial_p,\quad \mathbf{X}_5=x^2\partial_y+2x\partial_p+2\partial_q+4p\partial_z,$$

$$\mathbf{X}_6=q\partial_x+\left(pq-\frac{z}{2}\right)\partial_y+\frac{q^2}{2}\partial_p+\frac{q^3}{3}\partial_z,\quad \mathbf{X}_7=\frac{x^3}{6}\partial_y+\frac{x^2}{2}\partial_p+x\partial_q+(2px-2y)\partial_z,$$

$$\mathbf{X}_8=x\partial_x+y\partial_y-q\partial_q-z\partial_z,\quad \mathbf{X}_9=y\partial_y+p\partial_p+q\partial_q+2z\partial_z.$$

Abstractly, in the flat G_2/P_1 *model,* Π *and* L *correspond to the subspaces of* $\mathrm{Lie}(G_2)/\mathfrak{p}_1$ *spanned by* $\{e_{-\alpha_1},e_{-\alpha_1-\alpha_2}\}$ *and* $\{e_{-2\alpha_1-\alpha_2}\}$ *(modulo* $\mathfrak{p}_1$*). Only* $\mathfrak{gl}_2(\mathbb{C})\cong\mathfrak{g}_0\subset\mathfrak{p}_1$ *stabilizes both* Π *and* L*. Hence,* $\mathcal{S}\cong\mathfrak{p}_1^{\mathrm{op}}$ *(via a Lie algebra anti-automorphism).*

Proposition 5.5.4. *Consider a* $(2,3,5)$*-distribution* Π *and a line field* L *such that* $[\Pi,\Pi]=L\oplus\Pi$*. The infinitesimal symmetry algebra of* (Π,L) *is at most 9-dimensional, and this bound is sharp.*

Proof. For $A_3/P_{1,2}$, $\mathfrak{g}_0\cong\mathbb{C}^2\times A_1$, and $H^2(\mathfrak{g}_-,\mathfrak{g})_{+1}=\mathbb{V}_{-w\cdot\lambda_{\mathfrak{g}}}=\overset{4}{\times}\!\!-\!\!\!-\!\!\overset{-4}{\times}\!\!-\!\!\!-\!\!\overset{0}{\circ}$, where $w=(23)\in W_+^{\mathfrak{p}}(2)$. We have $J_w=I_w=\emptyset$, so $\mathfrak{a}(w)=\mathfrak{g}_-\oplus\mathfrak{a}_0$, $\mathfrak{p}_w^{\mathrm{op}}\cong A_1$, and $\dim(\mathfrak{a}_0)=|I_{\mathfrak{p}}|-1+\dim(\mathfrak{p}_w^{\mathrm{op}})=4$. Then $\mathfrak{S}_w=\dim(\mathfrak{a}(w))=\dim(\mathfrak{g}_-)+\dim(\mathfrak{a}_0)=5+4=9$. □

Remark 5.5.5. As indicated above, the structure is in fact not just a bracket-generating distribution since D has a splitting on it. However, it fits in naturally after discussing G_2/P_1 geometry.

---

[15] R.L. Bryant [5] analyzed the $(3,1)$ case by continuing Cartan's method of equivalence [11] and showed that no models with 5-dimensional symmetry exist. Thus, 4 is the maximal symmetry dimension for type $(3,1)$.

5.5.3. B_ℓ/P_ℓ *geometry,* $\ell \geq 3$. Let $\mathfrak{g} \cong B_\ell \cong \mathrm{SO}_{2\ell+1}(\mathbb{C})$, but all statements here still hold when considering $\mathrm{SO}_{\ell+1,\ell}$. The underlying structure for a (regular, normal) B_ℓ/P_ℓ geometry is a rank ℓ distribution on a $\binom{\ell+1}{2}$-manifold, with generic growth $(\ell, \binom{\ell+1}{2})$. This is a 2-graded geometry with $\mathfrak{g}_0 \cong \mathbb{C} \times A_{\ell-1} \cong \mathfrak{gl}_\ell(\mathbb{C})$, and Levi bracket inducing the $\mathfrak{g}_0$-module isomorphism $\bigwedge^2 \mathfrak{g}_{-1} \cong \mathfrak{g}_{-2}$. The $\ell = 3$ case was studied by R.L. Bryant [3], [4].

Theorem 5.5.6. *Let* $\ell \geq 3$. *For* $(\ell, \binom{\ell+1}{2})$*-distributions,* $\mathfrak{S} = \begin{cases} \frac{\ell(3\ell-7)}{2} + 10, & \ell \geq 4; \\ 11, & \ell = 3. \end{cases}$

Proof. We have $W^{\mathfrak{p}}_+(2) = \{(\ell, \ell-1)\}$. For $w = (\ell, \ell-1)$, $I_w = \emptyset$, so $\mathfrak{a}(w) = \mathfrak{g}_- \oplus \mathfrak{a}_0$, and

1. $\ell \geq 4$: $J_w = \{2, \ell-2, \ell-1\}$, so $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_{2,\ell-2,\ell-1} \subset A_{\ell-1}$, and $\dim(\mathfrak{a}_0) = \frac{\ell(3\ell-7)}{2} + 10 - \binom{\ell+1}{2}$;
2. $\ell = 3$: $J_w = \{1, 2\}$, so $\mathfrak{p}_w^{\mathrm{op}} \cong \mathfrak{p}_{1,2} \subset A_2$, and $\dim(\mathfrak{a}_0) = 5$;

since $\dim(\mathfrak{a}_0) = \dim(\mathfrak{p}_w^{\mathrm{op}})$. Then $\mathfrak{S} = \dim(\mathfrak{a}(w)) = \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0)$ yields the result. □

Let us exhibit some models in the $\ell = 3$ case. On a 6-manifold $(x_1, x_2, x_3, y_1, y_2, y_3)$, we specify 3-plane distributions by the vanishing of three 1-forms. The flat model is

$$\theta_1 = dy_1 - x_3 dx_2, \qquad \theta_2 = dy_2 - x_1 dx_3, \qquad \theta_3 = dy_3 - x_2 dx_1,$$

which has B_3 symmetry algebra (21-dimensional). Two new models are

$$\begin{array}{llll} \text{(a)}: & \theta_1 = dy_1 - x_3 dx_2, & \theta_2 = dy_2 - x_1 dx_3, & \theta_3 = dy_3 - x_2 dx_1 - {x_1}^2 {x_3}^2 dx_3; \\ \text{(b)}: & \theta_1 = dy_1 - x_3 dx_2, & \theta_2 = dy_2 - x_1 dx_3, & \theta_3 = dy_3 - (x_2 + y_3) dx_1. \end{array}$$

The respective 11 and 10-dimensional symmetry algebras (found using the `DifferentialGeometry` package in `Maple`) are given in Table 11.

	Symmetries
(a)	$\mathbf{X}_1 = \partial_{y_1}, \quad \mathbf{X}_2 = \partial_{y_2}, \quad \mathbf{X}_3 = \partial_{y_3}, \quad \mathbf{X}_4 = \partial_{x_2} + x_1 \partial_{y_3},$ $\mathbf{X}_5 = 2x_1 \partial_{x_1} - x_2 \partial_{x_2} - x_3 \partial_{x_3} - 2y_1 \partial_{y_1} + y_2 \partial_{y_2} + y_3 \partial_{y_3},$ $\mathbf{X}_6 = -x_1 \partial_{x_1} + 2x_2 \partial_{x_2} + x_3 \partial_{x_3} + 3y_1 \partial_{y_1} + y_3 \partial_{y_3},$ $\mathbf{X}_7 = x_1 \partial_{x_2} + (x_1 x_3 - y_2) \partial_{y_1} + \frac{1}{2} {x_1}^2 \partial_{y_3}, \quad \mathbf{X}_8 = x_3 \partial_{x_2} + \frac{1}{2} {x_3}^2 \partial_{y_1} + (x_1 x_3 - y_2) \partial_{y_3},$ $\mathbf{X}_9 = \frac{1}{2} \partial_{x_1} + \frac{1}{3} {x_3}^3 \partial_{x_2} + \frac{1}{4} {x_3}^4 \partial_{y_1} + \frac{1}{2} x_3 \partial_{y_2} + \frac{1}{3} x_1 {x_3}^3 \partial_{y_3},$ $\mathbf{X}_{10} = x_3 \partial_{x_1} + \frac{1}{2} {x_3}^4 \partial_{x_2} + \frac{2}{5} {x_3}^5 \partial_{y_1} + \frac{1}{2} {x_3}^2 \partial_{y_2} + (\frac{1}{2} x_1 {x_3}^4 - y_1 + x_2 x_3) \partial_{y_3},$ $\mathbf{X}_{11} = (2x_1 {x_3}^2 + 4x_3 y_2) \partial_{x_2} + 3\partial_{x_3} + (3x_2 + 2x_1 {x_3}^3 + 2{x_3}^2 y_2) \partial_{y_1}$ $\qquad + ({x_1}^2 {x_3}^2 + 4x_1 x_3 y_2 - 2{y_2}^2) \partial_{y_3}$
(b)	$\mathbf{X}_1 = \partial_{y_1}, \quad \mathbf{X}_2 = \partial_{y_2}, \quad \mathbf{X}_3 = \partial_{x_2} - \partial_{y_3}, \quad \mathbf{X}_4 = e^{x_1} \partial_{y_3}, \quad \mathbf{X}_5 = \partial_{x_3} + x_2 \partial_{y_1},$ $\mathbf{X}_6 = \partial_{x_1} + x_3 \partial_{y_2}, \quad \mathbf{X}_7 = x_2 \partial_{x_2} + y_1 \partial_{y_1} + y_3 \partial_{y_3}, \quad \mathbf{X}_8 = x_3 \partial_{x_3} + y_1 \partial_{y_1} + y_2 \partial_{y_2},$ $\mathbf{X}_9 = e^{-x_1} (\partial_{x_3} + (y_3 + x_2) \partial_{y_1} + (1 + x_1) \partial_{y_2}), \quad \mathbf{X}_{10} = x_1 \partial_{x_2} + (x_1 x_3 - y_2) \partial_{y_1} - (1 + x_1) \partial_{y_3}$

TABLE 11. Symmetries of new models for generic rank 3 distributions on 6-manifolds

Model (a) is submaximally symmetric and realizes the abstract model $\mathfrak{f}/\mathfrak{k}$ in Theorem 4.3.5. We now outline its construction. Take $\mathbb{C}^7$ with anti-diagonal symmetric $\mathbb{C}$-bilinear form, $\epsilon_i : \mathfrak{h} \to \mathbb{C}$, and simple roots as in Example 2.8.3. By Kostant's theorem,

$H^2(\mathfrak{g}_-, \mathfrak{g}) \cong$  $= \mathbb{V}_{-w\cdot\lambda_{\mathfrak{g}}}, \qquad w = (32),$

with lowest weight $-w\cdot\lambda_2 = -\alpha_1 + 3\alpha_3 = -\epsilon_1 + \epsilon_2 + 3\epsilon_3$ (homogeneity $+3$). Since $\Phi_w = \{\alpha_2 + 2\alpha_3, \alpha_3\}$, and $w(-\lambda_2) = -\lambda_1 - \lambda_2 + 2\lambda_3 = -\alpha_1 - \alpha_2$, then $\phi_0 = e_{\alpha_2+2\alpha_3} \wedge e_{\alpha_3} \otimes e_{-\alpha_1-\alpha_2}$. Letting $\mathfrak{a}_0 = \mathfrak{ann}(\phi_0)$,

$$\mathfrak{a} = \mathfrak{g}_- \oplus \mathfrak{a}_0 = \left(\begin{array}{ccc|c|ccc} a_{22}+3a_{33} & 0 & 0 & 0 & 0 & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 & 0 & 0 & 0 \\ a_{31} & a_{32} & a_{33} & 0 & 0 & 0 & 0 \\ \hline a_{41} & a_{42} & a_{43} & 0 & 0 & 0 & 0 \\ \hline a_{51} & a_{52} & 0 & -a_{43} & -a_{33} & 0 & 0 \\ a_{61} & 0 & -a_{52} & -a_{42} & -a_{32} & -a_{22} & 0 \\ 0 & -a_{61} & -a_{51} & -a_{41} & -a_{31} & -a_{21} & -(a_{22}+3a_{33}) \end{array}\right),$$

and let E_{ij} denote the matrix with $a_{ij} = 1$ and 0's elsewhere. Calculating matrix commutators determines the Lie algebra structure on $\mathfrak{a}$. If we take the following as root vectors

$$e_{-\alpha_2-2\alpha_3} = E_{52} \in \mathfrak{g}_{-2}; \qquad e_{-\alpha_3} = E_{43} \in \mathfrak{g}_{-1}; \qquad e_{-\alpha_1-\alpha_2} = E_{31} \in \mathfrak{g}_0,$$

then ϕ_0 yields the deformation $[E_{52}, E_{43}] = E_{31}$ of $\mathfrak{a}$. (Note that $[E_{52}, E_{43}] = 0$ in $\mathfrak{a}$.) This yields a filtered deformation $\mathfrak{f}$ of $\mathfrak{a}$, and model (a) is a local model for $\mathfrak{f}/\mathfrak{k}$, where $\mathfrak{k} = \mathfrak{a}_0$.

Appendix A. Representation-theoretic data for complex simple Lie algebras

We use the Bourbaki [2] / `LiE` [31] ordering of simple roots. Notation such as $0_r 1_s 2_t 1$ means $\alpha = \alpha_{r+1} + ... + \alpha_{r+s} + 2\alpha_{r+s+1} + ... + 2\alpha_{r+s+t} + \alpha_{r+s+t+1}$, where $\ell = \mathrm{rank}(\mathfrak{g}) = r + s + t + 1$.

$\mathfrak{g}$	$\dim(\mathfrak{g})$	$\lambda_{\mathfrak{g}}$	Dynkin diagram	Positive roots
A_ℓ $(\ell \geq 1)$ $\mathfrak{sl}(\ell+1,\mathbb{C})$	$\ell(\ell+2)$	$\lambda_1 + \lambda_\ell$	 α_1 α_2 α_3 $\alpha_{\ell-2}$ $\alpha_{\ell-1}$ α_ℓ	$0_r 1_s 0_t$ $(s \geq 1)$
B_ℓ $(\ell \geq 3)$ $\mathfrak{so}(2\ell+1,\mathbb{C})$	$\ell(2\ell+1)$	λ_2	 α_1 α_2 α_3 $\alpha_{\ell-2}$ $\alpha_{\ell-1}$ α_ℓ	$0_r 1_s 0_t$ $(s \geq 1)$ $0_r 1_s 2_t$ $(s,t \geq 1)$
C_ℓ $(\ell \geq 2)$ $\mathfrak{sp}(2\ell,\mathbb{C})$	$\ell(2\ell+1)$	$2\lambda_1$	 α_1 α_2 α_3 $\alpha_{\ell-2}$ $\alpha_{\ell-1}$ α_ℓ	$0_r 1_s 0_t$ $(s \geq 1)$ $0_r 1_s 2_t 1$ $(t \geq 1)$
D_ℓ $(\ell \geq 4)$ $\mathfrak{so}(2\ell,\mathbb{C})$	$\ell(2\ell-1)$	λ_2	 $\alpha_{\ell-1}$ α_1 α_2 α_3 $\alpha_{\ell-3}$ $\alpha_{\ell-2}$ α_ℓ	$0_r 1_s 0_t$ $(s,t \geq 1)$ $0_r 1_s 01$ $(s \geq 1)$ $0_r 1_s 2_t 11$ $(s \geq 1)$
G_2	14	λ_2		$10, 01, 11, 21, 31, 32$
F_4	52	λ_1		(Use `LiE`)
E_6 E_7 E_8	78 133 248	λ_2 λ_1 λ_8	 α_1 α_3 α_4 α_5 α_ℓ α_2	(Use `LiE`)

Appendix B. Submaximal symmetry dimensions

The tables to follow summarize data associated to the gap problem for all *regular, normal* parabolic geometries of *complex* or *split-real* type (G, P) with G *simple*. Let $\mathfrak{S}$ be the submaximal symmetry dimension, and define $\mathfrak{S}_w$ similarly but require $\mathrm{im}(\kappa_H) \subset \mathbb{V}_{-w\cdot\lambda_{\mathfrak{g}}}$. Summarizing:

1. Calculation of $W^{\mathfrak{p}}_+(2)$: Given $w \in W^{\mathfrak{p}}(2)$ (use Recipe 3), $w \in W^{\mathfrak{p}}_+(2)$ iff $Z(-w\cdot\lambda_{\mathfrak{g}}) \geq +1$.
2. Maximal non-flat Tanaka prolongation $\mathfrak{a}(w)$ for each $w \in W^{\mathfrak{p}}_+(2)$:
    - (a) Let $\phi_0 \in \mathbb{V}_{-w\cdot\lambda_{\mathfrak{g}}}$ be a lowest weight vector. Define $\mathfrak{a}(w) = \mathrm{pr}^{\mathfrak{g}}(\mathfrak{g}_-, \mathfrak{ann}(\phi_0))$.
    - (b) Find $J_w = \{j \notin I_{\mathfrak{p}} \mid \langle w\cdot\lambda_{\mathfrak{g}}, \alpha_j^\vee\rangle \neq 0\}$ and $I_w = \{i \in I_{\mathfrak{p}} \mid \langle w\cdot\lambda_{\mathfrak{g}}, \alpha_i^\vee\rangle = 0\}$.
    - (c) $(\mathfrak{g}, \mathfrak{p}, w)$ is PR iff $I_w = \emptyset$. If $I_w \neq \emptyset$, i.e. $(\mathfrak{g}, \mathfrak{p}, w)$ is NPR, find $(\bar{\mathfrak{g}}, \bar{\mathfrak{p}})$ using Recipe 6.
3. Dimensions: $\mathfrak{S} = \max_{w\in W^{\mathfrak{p}}_+(2)} \mathfrak{S}_w$, where $\mathfrak{S}_w$ are calculated as follows:
    - (a) Find $\dim(G/P) = \dim(\mathfrak{g}_-)$ and $\dim(\mathfrak{a}_0)$ by Recipes 1 and 5.
    - (b) Using $(\bar{\mathfrak{g}}, \bar{\mathfrak{p}})$, find $\dim(\mathfrak{a}_+)$ and hence $\dim(\mathfrak{a}) = \dim(\mathfrak{g}_-) + \dim(\mathfrak{a}_0) + \dim(\mathfrak{a}_+)$.
    - (c) Other than exceptional A_2/P and B_2/P, we always have $\mathfrak{S}_w = \dim(\mathfrak{a}(w))$. For the exceptions, see Table 12.

Note: $\mathfrak{a}_3 = 0$ always, and $\mathfrak{a}_2 \neq 0$ only for NPR-A/C (Section 3.6). In Tables 12, 13, 14, $\mathfrak{a}_1 = 0$.

By Remark 4.3.10, we may assume that P is *connected*, so that we can substantially simplify the calculation of $\mathfrak{S}_w$ by passing to the minimal twistor space (despite possible non-existence of such a twistor space when P is not connected).

G	P	$\dim(G/P)$	w	J_w	Twistor space type	$\dim \mathfrak{a}(w)$	$\mathfrak{S}_w$
A_2	P_1	2	(12)	$\{2\}$	–	4	3
	$P_{1,2}$	3	(12)	$\emptyset$	A_2/P_1	4	3
			(21)	$\emptyset$	A_2/P_2	4	3
B_2	P_1	3	(12)	$\{2\}$	–	5	4
	P_2	3	(21)	$\{1\}$	–	5	5
	$P_{1,2}$	4	(12)	$\emptyset$	B_2/P_1	5	4
			(21)	$\emptyset$	B_2/P_2	5	5

Table 12. Submaximal symmetry dimensions for geometries of type A_2/P and B_2/P

G	P	Range	$\dim(G/P)$	w	J_w	$\mathfrak{S}_w = \dim \mathfrak{a}(w)$
A_ℓ	P_1	$\ell \geq 3$	ℓ	(12)	$\{2,3,\ell\}$	$(\ell-1)^2+4$
	P_2	$\ell \geq 3$	$2(\ell-1)$	(21)	$\{3,\ell\}$	$(\ell-1)^2+5$
				(23)	$\begin{cases} \{1,4,\ell\}, & \ell \geq 4; \\ \{1\}, & \ell = 3 \end{cases}$	$\begin{cases} \ell^2-3\ell+10, & \ell \geq 4; \\ 9, & \ell = 3 \end{cases}$
	P_k	$3 \leq k \leq \lceil \frac{\ell}{2} \rceil$	$k(\ell+1-k)$	$(k,k+1)$	$\{1,k-1,k+2,\ell\}$	$\ell(\ell-1)-k(\ell-k)+6$
				$(k,k-1)$	$\{1,k-2,k+1,\ell\}$	$\ell(\ell-k)+(k-1)^2+6$
B_ℓ	P_1	$\ell \geq 3$	$2\ell-1$	(12)	$\{3\}$	$2\ell^2-5\ell+9$
C_ℓ	P_ℓ	$\ell \geq 3$	$\binom{\ell+1}{2}$	$(\ell,\ell-1)$	$\{1,\ell-2,\ell-1\}$	$\frac{\ell(3\ell-5)}{2}+5$
D_ℓ	P_1	$\ell \geq 4$	$2\ell-2$	(12)	$\begin{cases} \{3\}, & \ell \geq 5; \\ \{3,4\}, & \ell = 4 \end{cases}$	$2\ell^2-7\ell+12$
	P_ℓ	$\ell \geq 5$	$\binom{\ell}{2}$	$(\ell,\ell-2)$	$\{2,\ell-3,\ell-1\}$	$\frac{\ell(3\ell-11)}{2}+16$
E_6	P_6	–	16	(65)	$\{2,4\}$	45
E_7	P_7	–	27	(76)	$\{1,5\}$	76

Table 13. Submaximal symmetry dimensions for 1-graded geometries

G	P	Range	$\dim(G/P)$	w	J_w	$\mathfrak{S}_w = \dim \mathfrak{a}(w)$
A_ℓ	$P_{1,\ell}$	$\ell \geq 3$	$2\ell-1$	$(1,\ell)$	$\{2,\ell-1\}$	$(\ell-1)^2+4$
				(12)	$\begin{cases} \{2,3\}, & \ell \geq 4; \\ \{2\}, & \ell = 3 \end{cases}$	$(\ell-1)^2+4$
				$(\ell,\ell-1)$	$\begin{cases} \{\ell-2,\ell-1\}, & \ell \geq 4; \\ \{2\}, & \ell = 3 \end{cases}$	$(\ell-1)^2+4$
B_ℓ	P_2	$\ell \geq 3$	$4\ell-5$	(21)	$\{1,3\}$	$2\ell^2-5\ell+8$
				(23)	$\begin{cases} \{1,3,4\}, & \ell \geq 4; \\ \{1,3\}, & \ell = 3 \end{cases}$	$\begin{cases} 2\ell^2-7\ell+15, & \ell \geq 4; \\ 11, & \ell = 3 \end{cases}$
C_ℓ	P_1	$\ell \geq 2$	$2\ell-1$	(12)	$\begin{cases} \{2,3\}, & \ell \geq 3; \\ \{2\}, & \ell = 2 \end{cases}$	$\begin{cases} 2\ell^2-5\ell+8, & \ell \geq 3; \\ 5, & \ell = 2 \end{cases}$
D_ℓ	P_2	$\ell \geq 4$	$4\ell-7$	(21)	$\begin{cases} \{1,3\}, & \ell \geq 5; \\ \{1,3,4\}, & \ell = 4 \end{cases}$	$2\ell^2-7\ell+11$
				(23)	$\begin{cases} \{1,3,4\}, & \ell \neq 5; \\ \{1,3,4,5\}, & \ell = 5 \end{cases}$	$2\ell^2-9\ell+19$
		$\ell = 4$		(24)	$\{1,3,4\}$	15
G_2	P_2	–	5	(21)	$\{1\}$	7
F_4	P_1	–	15	(12)	$\{2,3\}$	28
E_6	P_2	–	21	(24)	$\{3,4,5\}$	43
E_7	P_1	–	33	(13)	$\{3,4\}$	76
E_8	P_8	–	57	(87)	$\{6,7\}$	147

Table 14. Submaximal symmetry dimensions for parabolic contact geometries

G	P	Range	$\dim(G/P)$	w	J_w	I_w	$(\overline{\mathfrak{g}}, \overline{\mathfrak{p}})$	$\mathfrak{a}_1$	$\mathfrak{a}_2$	Twistor space type	$\mathfrak{S}_w = \dim \mathfrak{a}(w)$
A_ℓ	$P_{1,2}$	$\ell \geq 3$	$2\ell - 1$	(21)	$\{3, \ell\}$	$\{1\}$	A_1/P_1	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
				(12)	$\{3, \ell\}$	$\emptyset$	–	–	–	A_ℓ/P_1	$(\ell-1)^2+4$
		$\ell = 3$		(23)	$\emptyset$	$\emptyset$	–	–	–	A_3/P_2	9
	$P_{1,3}$	$\ell \geq 4$	$3\ell - 4$	(31)	$\{2, 4, \ell\}$	$\emptyset$	–	–	–	–	$\ell^2 - 3\ell + 9$
				(12)	$\{2, \ell\}$	$\emptyset$	–	–	–	A_ℓ/P_1	$(\ell-1)^2+4$
		$\ell = 4$		(34)	$\{2\}$	$\emptyset$	–	–	–	A_4/P_3	14
	$P_{1,s}$	$4 \leq s \leq \ell - 1$	$\ell s - (s-1)^2$	(12)	$\{2, 3, \ell\}$	$\{s\}$	$A_{\ell-4}/P_{s-3}$	✓	–	A_ℓ/P_1	$(\ell-1)^2+4$
				$(1, s)$	$\{2, s-1, s+1, \ell\}$	$\emptyset$	–	–	–	–	$(\ell-2)s + (\ell-s)^2 + 6$
		$4 \leq s = \ell - 1$		$(\ell-1, \ell)$	$\{\ell - 2\}$	$\emptyset$	–	–	–	$A_\ell/P_{\ell-1}$	$(\ell-1)^2+5$
	$P_{2,s}$	$4 \leq s \leq \ell - 1$	$s(\ell+3-s) - 4$	(21)	$\{3, \ell\}$	$\{s\}$	$A_{\ell-4}/P_{s-3}$	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
		$4 \leq s = \ell - 1$		$(\ell-1, \ell)$	$\{1, \ell-2\}$	$\{2\}$	$A_{\ell-4}/P_1$	✓	–	$A_\ell/P_{\ell-1}$	$(\ell-1)^2+5$
	$P_{s,s+1}$	$2 \leq s \leq \ell - 2$	$(\ell-s)(s+1)+s$	$(s, s+1)$	$\{1, s-1, s+2, \ell\}$	$\{s+1\}$	A_1/P_1	✓	–	A_ℓ/P_s	$\ell(\ell-1) - s(\ell-s) + 6$
				$(s+1, s)$	$\{1, s-1, s+2, \ell\}$	$\{s\}$	A_1/P_1	✓	–	A_ℓ/P_{s+1}	$\ell(\ell-1) - s(\ell-s) + 6$
		$s = 2 \leq \ell - 2$		(21)	$\{\ell\}$	$\emptyset$	–	–	–	A_ℓ/P_2	$(\ell-1)^2+5$
	$P_{1,s,\ell}$	$3 \leq s \leq \ell - 2$	$\ell - 1 + (\ell+1-s)s$	$(1, \ell)$	$\{2, \ell-1\}$	$\{s\}$	$A_{\ell-4}/P_{s-2}$	✓	–	$A_\ell/P_{1,\ell}$	$(\ell-1)^2+4$
	$P_{1,2,3}$	$\ell \geq 3$	$3(\ell-1)$	(21)	$\{\ell\}\backslash\{3\}$	$\{1\}$	A_1/P_1	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
				(12)	$\{\ell\}\backslash\{3\}$	$\emptyset$	–	–	–	A_ℓ/P_1	$(\ell-1)^2+4$
		$\ell = 3$	6	(23)	$\emptyset$	$\{3\}$	A_1/P_1	✓	–	A_3/P_2	9
				(32)	$\emptyset$	$\emptyset$	–	–	–	A_3/P_3	8
				(13)	$\emptyset$	$\emptyset$	–	–	–	$A_3/P_{1,3}$	8
	$P_{1,2,s}$	$4 \leq s < \ell$	$s(\ell+3-s) - 3$	(21)	$\{3, \ell\}$	$\{1, s\}$	$A_1/P_1 \times A_{\ell-4}/P_{s-3}$	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
				(12)	$\{3, \ell\}$	$\{s\}$	$A_{\ell-4}/P_{s-3}$	✓	–	A_ℓ/P_1	$(\ell-1)^2+4$
	$P_{1,2,\ell}$	$\ell \geq 4$	$3(\ell-1)$	(21)	$\{3\}$	$\{1\}$	A_1/P_1	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
				(12)	$\{3\}$	$\emptyset$	–	–	–	A_ℓ/P_1	$(\ell-1)^2+4$
				$(1, \ell)$	$\{\ell - 1\}$	$\emptyset$	–	–	–	$A_\ell/P_{1,\ell}$	$(\ell-1)^2+4$
	$P_{1,2,3,s}$	$4 \leq s < \ell$	$3(\ell-1) + (\ell+1-s)(s-3)$	(21)	$\{\ell\}$	$\{1, s\}$	$A_1/P_1 \times A_{\ell-4}/P_{s-3}$	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
	$P_{1,2,3,\ell}$	$\ell \geq 4$	$4\ell - 6$	(21)	$\emptyset$	$\{1\}$	A_1/P_1	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
		$\ell = 4$		(34)	$\emptyset$	$\{4\}$	A_1/P_1	✓	–	A_ℓ/P_3	14
	$P_{1,2,s,t}$	$4 \leq s < t < \ell$	$2\ell - 1 + (\ell+1-s)(s-2)$ $+(\ell+1-t)(t-s)$	(21)	$\{3, \ell\}$	$\{1, s, t\}$	$A_1/P_1 \times A_{\ell-4}/P_{s-3,t-3}$	✓	✓	A_ℓ/P_2	$(\ell-1)^2+5$
	$P_{1,2,s,\ell}$	$4 \leq s < \ell$	$3(\ell-1) + (\ell-s)(s-2)$	(21)	$\{3\}$	$\{1, s\}$	$A_1/P_1 \times A_{\ell-4}/P_{s-3}$	✓	–	A_ℓ/P_2	$(\ell-1)^2+5$
		$4 \leq s = \ell - 1$		$(\ell-1, \ell)$	$\{\ell - 2\}$	$\{2, \ell\}$	$A_{\ell-4}/P_1 \times A_1/P_1$	✓	–	$A_\ell/P_{\ell-1}$	$(\ell-1)^2+5$
B_ℓ	P_3	$\ell \geq 4$	$6\ell - 12$	(32)	$\{1, 4\}$	$\emptyset$	–	–	–	–	$2\ell^2 - 7\ell + 16$
		$\ell = 3$		(32)	$\{1, 2\}$	$\emptyset$	–	–	–	–	11
	P_ℓ	$\ell \geq 4$	$\binom{\ell+1}{2}$	$(\ell, \ell-1)$	$\{2, \ell-2, \ell-1\}$	$\emptyset$	–	–	–	–	$\frac{\ell(3\ell-7)}{2} + 10$
	$P_{1,2}$	$\ell \geq 3$	$4\ell - 4$	(12)	$\{3\}$	$\{2\}$	A_1/P_1	✓	–	B_ℓ/P_1	$2\ell^2 - 5\ell + 9$
				(21)	$\{3\}$	$\emptyset$	–	–	–	B_ℓ/P_2	$2\ell^2 - 5\ell + 8$
	$P_{2,3}$	$\ell \geq 4$	$6\ell - 10$	(32)	$\{1, 4\}$	$\{2\}$	A_1/P_1	✓	–	B_ℓ/P_3	$2\ell^2 - 7\ell + 16$
		$\ell = 3$		(32)	$\{1\}$	$\emptyset$	–	–	–	B_3/P_3	11
	$P_{1,3}$	$\ell = 3$	8	(32)	$\{2\}$	$\emptyset$	–	–	–	B_3/P_3	11
	$P_{1,2,3}$	$\ell = 3$	9	(32)	$\emptyset$	$\emptyset$	–	–	–	B_3/P_3	11

TABLE 15. Submaximal symmetry dimensions for Yamaguchi-nonrigid geometries in type A and B, excluding 1-graded & parabolic contact geometries

G	P	Range	$\dim(G/P)$	w	J_w	I_w	$(\overline{\mathfrak{g}}, \overline{\mathfrak{p}})$	$\mathfrak{a}_1$	$\mathfrak{a}_2$	Twistor space type	$\mathfrak{S}_w = \dim \mathfrak{a}(w)$
C_ℓ	P_2	$\ell \geq 3$	$4\ell - 5$	(21)	$\{3\}$	$\emptyset$	–	–	–	–	$2\ell^2 - 5\ell + 9$
		$\ell = 3$		(23)	$\{1, 3\}$	$\emptyset$	–	–	–	–	11
	$P_{\ell-1}$	$\ell \geq 4$	$\frac{(\ell+4)(\ell-1)}{2}$	$(\ell - 1, \ell)$	$\{1, \ell - 2, \ell\}$	$\emptyset$	–	–	–	–	$\frac{\ell(3\ell-5)}{2} + 5$
	$P_{1,2}$	$\ell \geq 3$	$4\ell - 4$	(21)	$\{3\}$	$\{1\}$	A_1/P_1	✓	–	C_ℓ/P_2	$2\ell^2 - 5\ell + 9$
				(12)	$\{3\}$	$\emptyset$	–	–	–	C_ℓ/P_1	$2\ell^2 - 5\ell + 8$
	$P_{1,\ell}$	$\ell \geq 3$	$\frac{\ell^2+3\ell-2}{2}$	$(1, \ell)$	$\{2, \ell - 1\}$	$\emptyset$	–	–	–	–	$\frac{\ell(3\ell-5)}{2} + 5$
		$\ell \geq 4$		(12)	$\{2, 3\}$	$\{\ell\}$	$C_{\ell-3}/P_{\ell-3}$	✓	–	C_ℓ/P_1	$2\ell^2 - 5\ell + 8$
		$\ell = 3$		(12)	$\{2\}$	$\emptyset$	–	–	–	C_3/P_1	11
	$P_{2,\ell}$	$\ell \geq 4$	$\frac{\ell^2+5\ell-8}{2}$	(21)	$\{3\}$	$\{\ell\}$	$C_{\ell-3}/P_{\ell-3}$	✓	–	C_ℓ/P_2	$2\ell^2 - 5\ell + 9$
		$\ell = 3$		(21)	$\emptyset$	$\emptyset$	–	–	–	C_3/P_2	12
		$\ell = 3$		(23)	$\{1\}$	$\emptyset$	–	–	–	C_3/P_2	11
	$P_{\ell-1,\ell}$	$\ell \geq 4$	$\frac{\ell^2+3\ell-2}{2}$	$(\ell - 1, \ell)$	$\{1, \ell - 2\}$	$\emptyset$	–	–	–	$C_\ell/P_{\ell-1}$	$\frac{\ell(3\ell-5)}{2} + 5$
	$P_{1,2,3}$	$\ell \geq 3$	$6\ell - 9$	(21)	$\emptyset$	$\{1\}$	A_1/P_1	✓	–	C_ℓ/P_2	$2\ell^2 - 5\ell + 9$
	$P_{1,2,s}$	$4 \leq s < \ell$	$\frac{-6-3s^2+5s+4s\ell}{2}$	(21)	$\{3\}$	$\{1, s\}$	$A_1/P_1 \times C_{\ell-3}/P_{s-3}$	✓	✓	C_ℓ/P_2	$2\ell^2 - 5\ell + 9$
	$P_{1,2,\ell}$	$\ell \geq 4$	$\frac{\ell^2+5\ell-6}{2}$	(21)	$\{3\}$	$\{1, \ell\}$	$A_1/P_1 \times C_{\ell-3}/P_{\ell-3}$	✓	–	C_ℓ/P_2	$2\ell^2 - 5\ell + 9$
D_ℓ	P_3	$\ell \geq 5$	$6\ell - 15$	(32)	$\begin{cases} \{1, 4\}, & \ell \geq 6; \\ \{1, 4, 5\}, & \ell = 5 \end{cases}$	$\emptyset$	–	–	–	–	$2\ell^2 - 9\ell + 20$
	$P_{1,2}$	$\ell \geq 4$	$4\ell - 6$	(12)	$\begin{cases} \{3\}, & \ell \geq 5; \\ \{3, 4\}, & \ell = 4 \end{cases}$	$\{2\}$	A_1/P_1	✓	–	D_ℓ/P_1	$2\ell^2 - 7\ell + 12$
				(21)	$\begin{cases} \{3\}, & \ell \geq 5; \\ \{3, 4\}, & \ell = 4 \end{cases}$	$\emptyset$	–	–	–	D_ℓ/P_2	$2\ell^2 - 7\ell + 11$
	$P_{1,\ell}$	$\ell \geq 5$	$\frac{(\ell+2)(\ell-1)}{2}$	(12)	$\{3\}$	$\{\ell\}$	$D_{\ell-3}/P_{\ell-3}$	✓	–	D_ℓ/P_1	$2\ell^2 - 7\ell + 12$
		$\ell = 4$		(12)	$\{3\}$	$\emptyset$	–	–	–	D_4/P_1	16
				(42)	$\{3\}$	$\emptyset$	–	–	–	D_4/P_4	16
	$P_{2,3}$	$\ell \geq 5$	$6\ell - 13$	(32)	$\begin{cases} \{1, 4\}, & \ell \geq 6; \\ \{1, 4, 5\}, & \ell = 5 \end{cases}$	$\{2\}$	A_1/P_1	✓	–	D_ℓ/P_3	$2\ell^2 - 9\ell + 20$
	$P_{1,2,\ell}$	$\ell \geq 5$	$\frac{\ell^2+3\ell-6}{2}$	(12)	$\{3\}$	$\{2, \ell\}$	$A_1/P_1 \times D_{\ell-3}/P_{\ell-3}$	✓	–	D_ℓ/P_1	$2\ell^2 - 7\ell + 12$
		$\ell = 4$		(12)	$\{3\}$	$\{2\}$	A_1/P_1	✓	–	D_4/P_1	16
				(42)	$\{3\}$	$\{2\}$	A_1/P_1	✓	–	D_4/P_4	16
G_2	P_1	–	5	(12)	$\{2\}$	$\emptyset$	–	–	–	–	7
	$P_{1,2}$	–	6	(12)	$\emptyset$	$\emptyset$	–	–	–	G_2/P_1	7

TABLE 16. Submaximal symmetry dimensions for Yamaguchi-nonrigid geometries in type C, D, G, excluding 1-graded & parabolic contact geometries

APPENDIX C. DIRECT PROOF OF PROLONGATION-RIGIDITY FOR CONFORMAL GEOMETRY

Consider $\mathbb{C}^{n+2}$ with an anti-diagonal symmetric $\mathbb{C}$-bilinear form g, i.e. $g_{ij} = \delta_{i,j'}$, where $j' := n - j + 1$. Then

$$\mathfrak{so}(n+2,\mathbb{C}) = \left\{ \left(\begin{array}{c|c|c} a & \omega & 0 \\ \hline v & A & -\omega' \\ \hline 0 & -v' & -a \end{array} \right) : a \in \mathbb{C}, A \in \mathfrak{so}(n,\mathbb{C}), \omega \in (\mathbb{C}^n)^*, v \in \mathbb{C}^n \right\}$$

consists of matrices skew with respect to the anti-diagonal. We have $\mathfrak{g}_0 = \mathfrak{z}(\mathfrak{g}_0) \oplus \mathfrak{g}_0^{ss} = \mathbb{C} \oplus \mathfrak{so}(n,\mathbb{C})$. Symbolically, the brackets are:

1. $\mathfrak{g}_0 \times \mathfrak{g}_0 \to \mathfrak{g}_0$: $[(a,A),(b,B)] = (0,[A,B])$.
2. $\mathfrak{g}_0 \times \mathfrak{g}_1 \to \mathfrak{g}_1$: $[(a,A),\omega] = \omega(a - A)$.
3. $\mathfrak{g}_0 \times \mathfrak{g}_{-1} \to \mathfrak{g}_{-1}$: $[(a,A),v] = (A - a)v$.
4. $\mathfrak{g}_1 \times \mathfrak{g}_{-1} \to \mathfrak{g}_0$: $[\omega, v] = (\omega v, \omega' v' - v\omega)$.

The $\mathfrak{g}_0$-module decomposition of $H^2(\mathfrak{g}_-,\mathfrak{g})$ is given by

$$H^2(\mathfrak{g}_-,\mathfrak{g}) \cong \begin{cases} \left(\bigwedge^2 \mathfrak{g}_1 \otimes \mathfrak{g}_0^{ss}\right)_0, & n \geq 4 \quad \text{(Weyl tensors);} \\ \begin{array}{|c|c|}\hline & \\ \hline \\ \cline{1-1} \end{array}(\mathfrak{g}_1) & n = 3 \quad \text{(Cotton tensors).} \end{cases}$$

If $n \neq 4$, $H^2(\mathfrak{g}_-,\mathfrak{g})$ is $\mathfrak{g}_0$-irreducible; if $n = 4$, it splits into self-dual and anti-self-dual submodules.

Let $\{e_i\}$ and $\{\omega^j\}$ be the standard (dual) bases of $\mathbb{C}^n$ and $(\mathbb{C}^n)^*$ respectively. Define $E_i{}^j = e_i \otimes \omega^j$ and $R_\ell{}^k = E_\ell{}^k - E_{k'}{}^{\ell'} \in \mathfrak{g}_0^{ss}$. Note $R_\ell{}^k = -R_{k'}{}^{\ell'}$, and we require $k \neq \ell'$. Let $\omega^{ijk}{}_\ell = \omega^i \wedge \omega^j \otimes R_\ell{}^k$. An element $W = W_{ijk}{}^\ell \omega^{ijk}{}_\ell \in \bigwedge^2 \mathfrak{g}_1 \otimes \mathfrak{g}_0^{ss}$ has the index symmetries $W_{ijk}{}^\ell = -W_{jik}{}^\ell$, $W_{ijk}{}^\ell = -W_{ij\ell'}{}^{k'}$.

Letting $A = \sum_{i<j'} A^i_j R_i{}^j$ and using the bracket relations above, the $\mathfrak{g}_0$-action on $\bigwedge^2 \mathfrak{g}_1 \otimes \mathfrak{g}_0^{ss}$ is

$$\widetilde{W} := (a,A) \cdot W = \left(2aW_{ijk}{}^\ell - W_{rjk}{}^\ell A^r_i - W_{irk}{}^\ell A^r_j - W_{ijr}{}^\ell A^r_k + W_{ijk}{}^r A^\ell_r\right) \omega^{ijk}{}_\ell$$

$$\widetilde{W}_{ijk}{}^\ell = 2aW_{ijk}{}^\ell - W_{rjk}{}^\ell A^r_i - W_{irk}{}^\ell A^r_j + \frac{1}{2}\left(-W_{ijr}{}^\ell A^r_k + W_{ijr}{}^{k'} A^r_{\ell'} + W_{ijk}{}^r A^\ell_r - W_{ij\ell'}{}^r A^{k'}_r\right),$$

where we have accounted for "anti-diagonal skew-symmetry" in the last line above.

Similarly when $n = 3$, given $C = C_{ijk}\omega^i \wedge \omega^j \otimes \omega^k$, and $\widetilde{C} = (a,A) \cdot C$, we have

$$\widetilde{C}_{ijk} = 3aC_{ijk} - C_{rjk}A^r_i - C_{irk}A^r_j - C_{ijr}A^r_k.$$

Lemma C.0.7. *Conformal geometry in dimension ≥ 3 is prolongation-rigid.*

Proof. Let $\omega \in \mathfrak{g}_1$ and $\mathfrak{k} = [\omega, \mathfrak{g}_{-1}]$. Use the $G_0 = CO(\mathfrak{g}_{-1})$ action to normalize ω (over $\mathbb{C}$) to one of two values, namely a null vector or a non-null vector. Suppose first that $n \geq 4$.

Case 1: ω is null. Take $\omega = \omega^1$. Then $\mathfrak{k} = \{(v^1, -v^i R_i{}^1) : v \in \mathfrak{g}_{-1}\}$ and $\dim(\mathfrak{k}) = n - 1$. (Note $R_n{}^1 = 0$.) Letting $(a,A) = (1, E^n_n - E^1_1)$, we obtain

$$0 = \widetilde{W}_{ijk}{}^\ell = W_{ijk}{}^\ell \left(2 - A^i_i - A^j_j + \frac{1}{2}\left(-A^k_k + A^\ell_\ell - A^{\ell'}_{\ell'} + A^{k'}_{k'}\right)\right).$$

Since $A^n_n - A^1_1 = 2$, then the only possible nonzero components are $W_{njn}{}^\ell = -W_{nj\ell'}{}^1 = -W_{jnn}{}^\ell$ for $1 < j, \ell < n$, i.e. $W = W_{njn}{}^\ell \omega^{njn}{}_\ell$. Now let $(a,A) = (0, R_c{}^1)$, $1 < c < n$. Note $A^c_1 = 1 = -A^n_{c'}$. Then for $1 < j, \ell < n$ with $j \neq c'$, we have $0 = \tilde{W}_{c'jn}{}^\ell = -W_{rjn}{}^\ell A^r_{c'} = W_{njn}{}^\ell$. Thus, $W = 0$.

Case 2: ω is non-null. Take $\omega = \omega^1 + \omega^n$. Then $\mathfrak{k} = \{(v^1 + v^n, v^{i'}(R_1{}^i + R_n{}^i)) : v \in \mathfrak{g}_{-1}\}$ and $\dim(\mathfrak{k}) = n$. In particular, note that $(1, 0_{n\times n}) \in \mathfrak{k}$, and $0 = \widetilde{W} = (1, 0_{n\times n}) \cdot W = 2W$.

Now consider $n = 3$. If ω is null, then letting $(a,A) = (1, E_n{}^n - E_1{}^1)$ yields $0 = \widetilde{C}_{ijk} = C_{ijk}(3 - A^i_i - A^j_j - A^k_k)$, so $C_{ijk} = 0$ since $i \neq j$. If ω is not null, then as above $(1, 0_{n\times n}) \in \mathfrak{k}$, so again $C = 0$. ☐

References

[1] R.J. Baston and M.G. Eastwood, The Penrose transform: its interaction with representation theory, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1989.

[2] N. Bourbaki, Groupes et algèbres de Lie, Chap. 4, 5 et 6, Hermann, Paris, 1968.

[3] R.L. Bryant, Some aspects of the local and global theory of Pfaffian systems, Ph.D. thesis, University of North Carolina at Chapel Hill (1979).

[4] R.L. Bryant, Conformal geometry and 3-plane fields on 6-manifolds, Developments of Cartan Geometry and Related Mathematical Problems, RIMS Symposium Proceedings, Kyoto University, vol. **1502** (July, 2006), 1–15.

[5] R.L. Bryant, private communication, 2013.

[6] R.L. Bryant, M.G. Eastwood, A.R. Gover, and K. Neusser, Some differential complexes within and beyond parabolic geometry, arXiv:1112.2142v3 (2012).

[7] A. Čap, Correspondence spaces and twistor spaces for parabolic geometries, J. Reine Angew. Math **582** (2005), 143–172.

[8] A. Čap, Automorphism groups of parabolic geometries, in: Proceedings of the 24th Winter School on Geometry and Physics, Srni, 2004, Rend. Circ. Mat. Palermo (2) Suppl. **75** (2005), 233–239.

[9] A. Čap and K. Neusser, On automorphism groups of some types of generic distributions, Differential Geometry and its Applications, vol. **27**, no. 6 (2009), 769–779.

[10] A. Čap and J. Slovák, Parabolic Geometries I: Background and General Theory, Mathematical Surveys and Monographs, vol. 154, American Mathematical Society, 2009.

[11] E. Cartan, Les systèmes de Pfaff à cinq variables et les équations aux dérivées partielles du second ordre, Ann. Sci. Ecole Norm. Sup (3) **27** (1910), 109–192.

[12] E. Cartan, Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes. I, Ann. Mat. Pura Appl. **11** (1932), 17–90.

[13] S.S. Chern, The geometry of the differential equation $y''' = F(x, y, y', y'')$, Sci. Rep. Nat. Tsing Hua Univ. (A) **4** (1940), 97–111.

[14] S. Casey, M. Dunajski, and P. Tod, Twistor geometry of a pair of second order ODEs, arXiv:math.DG/1203.4158v2 (2012)

[15] G. Darboux, Leçons sur la théorie générale des surfaces et les applications géométriques du calcul infinitésimal, II partie (1887), IV partie (1896), Gauthier-Villar, Paris.

[16] B. Doubrov and D. The, Maximally degenerate Weyl tensors in Riemannian and Lorentzian signatures, arXiv:1305.3499 [math.DG] (2013).

[17] I.P. Egorov, Collineations of projectively connected spaces, Doklady Akad. Nauk SSSR **80** (1951), 709–712.

[18] I.P. Egorov, Maximally mobile Riemannian spaces V_4 of nonconstant curvature, Dokl. Akad. Nauk SSSR **103** (1955), 9–12.

[19] I.P. Egorov, Riemannian spaces of second lacunarity. (Russian) Dokl. Akad. Nauk SSSR **111**, 276–279 (1956).

[20] I.P. Egorov, Maximally mobile Einstein spaces of nonconstant curvature. (Russian) Dokl. Akad. Nauk SSSR **145**, 975–978 (1962). English transl.: Sov. Math., Dokl. **3**, 1124–1127 (1962).

[21] I.P. Egorov, Automorphisms in generalized spaces, VINITI, Moscow **10** (1978), 147–191; English: Journal of Soviet Mathematics **14**, no. 3 (1980), 1260–1287.

[22] M.E. Fels, The equivalence problem for systems of second-order ordinary differential equations, Proc. London Math. Soc. **71** (1995), 221–240.

[23] G. Fubini, Sugli spazi che ammettono un gruppo continuo di movimenti, Ann. di Mat. pura e applicata **8** (1903), 39–81.

[24] M. Godlinski and P. Nurowski, Geometry of third-order ODEs, arXiv:0902.4129v1 (2009).

[25] D.A. Grossman, Torsion-free path geometries and integrable second order ODE systems, Selecta Math. **6**, no.4 (2000), 399–342.

[26] M. Hammerl, Homogeneous Cartan geometries, Archivum Mathematicum **43**, no. 5 (2007), 431–442.

[27] S. Kobayashi, Transformation groups in differential geometry, Springer Verlag (1972).

[28] B. Kostant, Lie algebra cohomology and the generalized Borel–Weil theorem, Ann. Math. **74**, no. 2 (1961), 329–387.

[29] B. Kruglikov, Finite-dimensionality in Tanaka theory. Ann. Inst. H. Poincaré Anal. Non Linéaire **28**, no.1, (2011), 75–90.

[30] B. Kruglikov, The gap phenomenon in the dimension study of finite type systems, Cent. Eur. J. Math. **10**, no. 5 (2012), 1605–1618.

[31] `LiE`, available at `http://wwwmathlabo.univ-poitiers.fr/~maavl/LiE/`

[32] T. Morimoto, On the intransitive Lie algebras whose transitive parts are infinite and primitive, J. Math. Soc. Japan, **29**, no. 1 (1977), 35–65.

[33] T. Nagano, On conformal transformations of Riemannian spaces, J. Math. Soc. Japan **10**, no. 1, (1958).

[34] R. Penrose, A spinor approach to general relativity, Ann. Phys. **10**, no. 2 (1960), 171–201.

[35] A.Z. Petrov, Classification of spaces defining gravitational fields. (Russian), Kazan. Gos. Univ. Uč. Zap. **114**, no. 8 (1954), 55–69. English translation: A.Z. Petrov, Classification of spaces defining gravitational fields, Gen. Rel. Grav. **32**, no.8 (2000), 1665–1685.

[36] H. Sato, A.Y. Yoshikawa, Third order ordinary differential equations and Legendre connections, J. Math. Soc. Japan **50**, no. 4 (1998), 993–1013.

[37] J. Šilhan, Online `LiE` implementation of Kostant's algorithm at `http://bart.math.muni.cz/~silhan/lie/`. WARNING: We use *lowest* weights to describe components of $H^2(\mathfrak{g}_-, \mathfrak{g})$, while Šilhan's implementation is in terms of *highest* weights. To get our weights, use Šilhan's routines for $H^2(\mathfrak{p}_+, \mathfrak{g})$ (but then insert a negative sign on his homogeneities).

[38] H. Stephani, D. Kramer, M. Maccallum, C. Hoenselaers, E. Herlt, Exact Solutions of Einstein's Field Equations, 2nd ed., Cambridge University Press, 2003.

[39] F. Strazzullo, Symmetry analysis of general rank-3 Pfaffian systems in five variables, Ph.D. thesis, Utah State University (2009).

[40] N. Tanaka, On differential systems, graded Lie algebras and pseudogroups, J. Math. Kyoto. Univ. **10** (1970), 1–82.

[41] N. Tanaka, On the equivalence problems associated with simple graded Lie algebras, Hokkaido Math. J. **8** (1979), 23–84.

[42] M.A. Tresse, Déterminations des invariants ponctuels de l'équation différentielle ordinaire du second ordre $y'' = \omega(x, y, y')$, Leipzig: Hirzel (1896).

[43] C. Wafo Soh, F.M. Mahomed, and C. Qu, Contact symmetry algebras of scalar ordinary differential equations, Nonlinear Dynamics **28** (2002), 213–230.

[44] H.C. Wang, On Finsler spaces with completely integrable equations of Killing, J. Lond. Math. Soc. **22** (1947), 5–9.

[45] K. Yamaguchi, Differential systems associated with simple graded Lie algebras, Advanced Studies in Pure Mathematics **22**, Progress in Differential Geometry (1993), 413–494.

[46] K. Yamaguchi, G_2-geometry of overdetermined systems of second order, Analysis and geometry in several complex variables (Katata, 1997), 289–314, Trends Math., Birkhäuser Boston, Boston, MA, 1999.

[47] I. Zelenko, On Tanaka's prolongation procedure for filtered structures of constant type, SIGMA **094**, vol. 5 (2009), 21 pp.

Department of Mathematics and Statistics, University of Tromsø, 90-37, Norway
E-mail address: `boris.kruglikov@uit.no`

Mathematical Sciences Institute, Australian National University, ACT 0200, Australia
E-mail address: `dennis.the@anu.edu.au`